#### ERGA Assembly Report

v24.04.03 beta

Tags: ERGA-BGE

TxID	488418	
ToLID	wsMarCava1	
Species	Marifugia cavatica	
Class	Polychaeta	
Order	Sabellida	

Genome Traits	Expected	Observed
Haploid size (bp)	1,215,561,629	1,324,532,507
Haploid Number	14 (source: ancestor)	11
Ploidy	2 (source: ancestor)	2
Sample Sex	Unknown	Unknown

#### EBP metrics summary and curation notes

Obtained EBP quality metric for pri: 5.8.Q55

The following metrics were automatically flagged as below EBP recommended standards or different from expected:

- . Observed Haploid Number is different from Expected
- . BUSCO single copy value is less than 90% for pri
- . BUSCO duplicated value is more than 5% for pri
- . More than 1000 gaps/Gbp for pri

#### Curator notes

```
. Interventions/Gb: 557
. Contamination notes: "Total length of scaffolds removed: 2,641,141 (0.2
%);Scaffolds removed: 24 (3.3 %);Largest scaffold removed: (1,646,899);FCS-GX
contaminant species (number of scaffolds; total length of scaffolds):;Terrimonas sp.,
CFB group bacteria (1; 1,646,899); Sediminibacterium salmoneum, CFB group bacteria (1;
188,069); Candidatus Methylophosphatis roskildensis, b-proteobacteria (1;
121,725); Pedobacter quisquiliarum, CFB group bacteria (1; 89,660); Reyranella sp.,
a-proteobacteria (1; 74,734); Solimonas fluminis, g-proteobacteria (1;
72,394); Paludibacter sp. UBA5753, CFB group bacteria (1; 64,682); Chitinophaga
chungangae, CFB group bacteria (1; 38,422); Pedobacter ureilyticus, CFB group bacteria
(1; 34,861); Lewinella cohaerens, CFB group bacteria (1; 32,076); Limnohabitans parvus,
b-proteobacteria (1; 31,546); Iodobacter ciconiae, b-proteobacteria (1;
30,257); Gelidibacter maritimus, CFB group bacteria (1; 28,085); Cupriavidus lacunae,
b-proteobacteria (1; 24,928); Dyadobacter beijingensis, CFB group bacteria (1;
21,585); Hoeflea marina, a-proteobacteria (1; 20,549); Methyloversatilis sp.,
b-proteobacteria (1; 19,482); Rhodoferax sediminis, b-proteobacteria (1;
```


- 18,802);Haliscomenobacter hydrossis, CFB group bacteria (1; 18,768);Limnovirga soli, CFB group bacteria (1; 18,687);Ottowia beijingensis, b-proteobacteria (1; 13,239);Dyadobacter jejuensis, CFB group bacteria (1; 2,000);Niastella soli, CFB group bacteria (1; 1,000)"
- . Other observations: "PacBio is from a ULI library, so lower contig N50 is expected; Hi-C from a different individual (wsMarCava2), so assembly is not Hi-C phased; Expected haploid number from GoaT is an ancestral estimate, likely incorrect in this case."

# Quality metrics table

Metrics	Pre-curation pri	Curated pri
Total bp	1,364,684,544	1,324,532,507
GC %	36.41	36.41
Gaps/Gbp	2,720.04	2,829.68
Total gap bp	742,400	749,600
Scaffolds	728	581
Scaffold N50	91,216,656	144,721,733
Scaffold L50	6	5
Scaffold L90	14	10
Contigs	4,440	4,329
Contig N50	569,563	563,563
Contig L50	707	700
Contig L90	2,476	2,432
QV	55.2	55.2
Kmer compl.	97.46	98.68
BUSCO sing.	81.2%	80.7%
BUSCO dupl.	7.3%	8.1%
BUSCO frag.	6.6%	6.2%
BUSCO miss.	4.9%	5.0%


Warning: BUSCO versions or lineage datasets are not the same across results

# HiC contact map of curated assembly




pri [LINK]


# K-mer spectra of curated assembly



Distribution of k-mer counts per copy numbers found in asm




Distribution of k-mer counts coloured by their presence in reads/assemblies



Distribution of k-mer counts per copy numbers found in asm

### Post-curation contamination screening



pri. Bubble plot circles are scaled by sequence length, positioned by coverage and GC
proportion, and coloured by taxonomy. Histograms show total assembly length
distribution on each axis.

## Data profile

Data	PacBio HiFi	Arima v2
Coverage	32x	83x

### Assembly pipeline

# Curation pipeline

Submitter: Thomas Mathers

Affiliation: WSI

Date and time: 2024-09-23 12:29:12 CEST