

ERGA Assembly Report

v24.10.15

Tags: ERGA-BGE

TxID	243007
ToLID	iqAioThal3
Species	Aiolopus thalassinus
Class	Insecta
Order	Orthoptera

Genome Traits	Expected	Observed
Haploid size (bp)	6,979,785,760	7,281,159,242
Haploid Number	11 (source: direct)	12
Ploidy	2 (source: ancestor)	2
Sample Sex	X0	X0

EBP metrics summary and curation notes

The following metrics were automatically flagged as below EBP recommended standards or different from expected:

- . Observed Haploid Number is different from Expected

Curator notes


- . Interventions/Gb: 39
- . Contamination notes: "Contaminations were identified and removed with FCS-GX (removed: 4 contigs, 150Kb). Mitochondrial genome was removed from the assembly."
- . Other observations: "PacBio reads were assembled with hifiiasm. FCS-GX detected a few contaminations which where removed. Purge_dups was run to remove haplotypic duplications. But it removed to much sequence and even the Busco score dropped significantly. Therefore the Busco gene set was used to guide the purging. Afterwards Tiara was used to remove archaea|bacteria|organelle hits. Due to time issues and compute cluster down-time the scaffolding was done manually within the curation step. This worked well as the input assembly consisted of just 230 contigs. The curation revealed more contig misjoins than usual (76 contigs cuts were necessary). This is probably caused by the large heterozygosity (1.6%). Several large-scale structural haplotypic variations can be seen in the HiC-map. The X-chromosome was identified by synteny analysis with other Acrididae (e.g. GCF_021461395.2_iqSchAmer2) assemblies, but also the PacBio read coverage showed a 1n coverage pattern."

Quality metrics table

Metrics	Pre-curation pri	Curated pri
Total bp	7,497,130,708	7,281,159,242
GC %	41.47	41.47
Gaps/Gbp	0	28.02
Total gap bp	0	40,800
Scaffolds	227	48
Scaffold N50	63,728,264	765,756,381
Scaffold L50	33	4
Scaffold L90	115	9
Contigs	227	252
Contig N50	63,728,264	52,464,910
Contig L50	33	36
Contig L90	115	127
QV	72.0441	
Kmer compl.	77.349	
BUSCO sing.	93.4%	96.1%
BUSCO dupl.	4.9%	2.5%
BUSCO frag.	0.6%	0.4%
BUSCO miss.	1.1%	1.1%

BUSCO: 6.0.0 (euk_genome_min, miniprot) / Lineage: insecta_odb12 (genomes:79, BUSCOs:3114)


HiC contact map of curated assembly



pri [\[LINK\]](#)

K-mer spectra of curated assembly

Post-curation contamination screening



pri. Bubble plot circles are scaled by sequence length, positioned by coverage and GC proportion, and coloured by taxonomy. Histograms show total assembly length distribution on each axis.

Data profile

Data	HiFi	HiC
Coverage	53x	210x

Assembly pipeline

- **Hifiasm**
 - |_ *ver*: 0.25.0-r726
 - |_ *key param*: 13
- **purge_dups**
 - |_ *ver*: 1.2.6
 - |_ *key param*: NA

Curation pipeline

- **GRIT_Rapid**
 - |_ *ver*: 1a3d79a8
 - |_ *key param*: NA

Submitter: Guilherme Dias Martin Pippel

Affiliation: SciLifeLab

Date and time: 2026-02-14 17:21:27 CET