

Midterm 2 Standard 16 - Analyzing Recurrences II: Tree method

Due Date May 3rd
Name **Your Name**
Student ID **Your Student ID**

Contents

1	Instructions	1
2	Standard 16 - Analyzing Recurrences II: Tree method	2
2.1	Problem 1	2

1 Instructions

- The solutions **should be typed**, using proper mathematical notation. We cannot accept hand-written solutions. Here's a short intro to L^AT_EX.
- You should submit your work through the **class Canvas page** only. Please submit one PDF file, compiled using this L^AT_EX template.
- You may not need a full page for your solutions; pagebreaks are there to help Gradescope automatically find where each problem is. Even if you do not attempt every problem, please submit this document with no fewer pages than the blank template (or Gradescope has issues with it).
- You **may not collaborate with other students. Copying from any source is an Honor Code violation. Furthermore, all submissions must be in your own words and reflect your understanding of the material.** If there is any confusion about this policy, it is your responsibility to clarify before the due date.
- Posting to **any** service including, but not limited to Chegg, Discord, Reddit, StackExchange, etc., for help on an assignment is a violation of the Honor Code.

2 Standard 16 - Analyzing Recurrences II: Tree method

2.1 Problem 1

Problem 1. Using the tree method, find a suitable function $f(n)$ such that $T(n) \in \Theta(f(n))$. Show all work.

$$T(n) = \begin{cases} 1 & : n < 3, \\ T(n/3) + T(2n/3) & : n \geq 3. \end{cases}$$

Answer. We have two base cases in this problem. The first is when $n = 3^k < 3$ and the other is when $n = \frac{2^k}{3}$. Solving for both of these we get base cases at $k = \log_3(n)$ and $k = \log_{\frac{3}{2}}(n)$.

At each level i we have 2^i nodes. The non recursive work done at each node is $(\frac{1}{3} + \frac{2}{3})$. So the total work done at each level is $n(\frac{1}{3} + \frac{2}{3})^i$. Thus:

$$T(n) = \sum_{i=0}^{\log_{\frac{3}{2}}(n) - \log_3(n)} n \left(\frac{1}{3} + \frac{2}{3}\right)^i \quad (1)$$

$$= n \sum_{i=0}^{\log_{\frac{3}{2}}(n) - \log_3(n)} \left(\frac{1}{3} + \frac{2}{3}\right)^i \quad (2)$$

$$= n * (3)^{\log_{\frac{3}{2}}(n) - \log_3(n)} \quad (3)$$

$$(4)$$

So $T(n) \in \Theta(f(n))$

□