

Quiz 15 - Solve Recurrence by Unrolling

Due Date March 11
Name **Your Name**
Student ID **Your Student ID**

Contents

1	Instructions	1
2	Standard 15 - Unrolling	2

1 Instructions

- The solutions **should be typed**, using proper mathematical notation. We cannot accept hand-written solutions. Here's a short intro to \LaTeX .
- You should submit your work through the **class Canvas page** only. Please submit one PDF file, compiled using this \LaTeX template.
- You may not need a full page for your solutions; pagebreaks are there to help Gradescope automatically find where each problem is. Even if you do not attempt every problem, please submit this document with no fewer pages than the blank template (or Gradescope has issues with it).
- You **may not collaborate with other students. Copying from any source is an Honor Code violation. Furthermore, all submissions must be in your own words and reflect your understanding of the material.** If there is any confusion about this policy, it is your responsibility to clarify before the due date.
- Posting to **any** service including, but not limited to Chegg, Discord, Reddit, StackExchange, etc., for help on an assignment is a violation of the Honor Code.

2 Standard 15 - Unrolling

Problem 1. Using the unrolling method, find a suitable function $f(n)$ such that $T(n) \in \Theta(f(n))$. Show all work.

$$T(n) = \begin{cases} 5 & : n < 2, \\ 7T(n-2) + 9 & : n \geq 2. \end{cases}$$

Answer. First we determine the number of times that we need to unroll. We hit a base case when $n - 2k < 2$. Solving for k we get that.

$$n < 2 + 2k$$

$$n - 2 < 2k$$

$$\frac{n-2}{2} < k$$

So we need to unroll until $k = \lceil \frac{n-2}{2} \rceil$

When we unroll $T(n - 2i)$, we obtain $7T(n - 2(i + 1)) + 9$. So

$$T(n - 2i) = 7 * \lceil 2T(n - 2(i + 2)) + 9 \rceil + 9 = 7^2 T(n - 2(i + 2)) + 9 * 9 + 9^0 * 9.$$

Thus $T(n) = (\text{cost of base case}) * (\text{number of times the base case is reached}) + \sum_{i=0}^{\lceil \frac{n-2}{2} \rceil} 7^i * 9$

$$T(n) = 1 * 7^{\lceil \frac{n-2}{2} \rceil} + 9 * \sum_{i=0}^{\lceil \frac{n-2}{2} \rceil} 7^i$$

$$T(n) = 7^{\lceil \frac{n-2}{2} \rceil} + 9 * \frac{7^{\lceil \frac{n-2}{2} \rceil + 1} - 1}{2 - 1}$$

$$T(n) = 7^{\lceil \frac{n-2}{2} \rceil} + 9 * 7^{\lceil \frac{n-2}{2} \rceil + 1}$$

Taking the high-order term, we obtain that $T(n) \in \Theta 7^{\lceil \frac{n-2}{2} \rceil} = \Theta(7^{n/2})$

□