
MATHEMATICAL PHYSICS: A COMPLETE GUIDE

PHYS 435

SEPTEMBER 8, 2021

ELIJAH THOMPSON,
PHYSICS AND MATH HONORS

Solo Pursuit of Learning



Contents

I	Complex Analysis	3
1	Properties of Complex Functions	4
1.1	Elementary Definitions	4
1.2	Multivalued Functions and Branch Cuts	4
1.3	Analytic Functions and the Cauchy-Riemann Equations	4
1.4	Singularities and Zeros	4
1.5	Conformal Mappings	4
2	Power Series and Laurent Series	5
2.1	Power Series Fundamentals	5
2.2	Laurent Series	5
2.3	Series Operations	5
3	Complex Integration	6
3.1	Complex Integral	6
3.2	Cauchy's Theorems	6
3.3	Residue Theorem	6
3.4	Contour Integrals	6
4	Applications of Complex Functions	7
4.1	Complex Potentials	7
4.2	Finding Zeros	7
4.3	Inverse Laplace	7
4.4	Stokes' Equations and Airy Integrals	7
4.5	WKB Methods, and Integral Approximations	7
II	PDEs	8
5	General and Particular Solutions	9
5.1	Important Examples and Motivation	9
5.2	General Forms of Solutions	9
5.3	Wave and Diffusion Equations	9
5.4	Existence and Uniqueness	9
6	Fourier Series	10
6.1	Initial Definitions and Dirichlet Conditions	10
	Symmetry Conditions	11
6.2	Discontinuous and Non-Periodic Functions	12
	Discontinuities	12

Non-Periodic Functions	13
6.3 Integration and Differentiation	13
6.4 Complex Fourier Series	14
Parseval's Theorem	15
7 Integral Transforms	17
7.1 Fourier Transform	17
Construction and Definitions	17
The Uncertainty Principle	18
Dirac δ -function	20
Properties of Fourier Transforms	22
Odd and Even Functions	23
Convolution and Deconvolution	25
Correlation Functions and Energy Spectra	26
Parseval's Theorem	28
7.2 Multidimensional Fourier Transform	28
7.3 Laplace Transform	29
General Transform	31
8 Separation of Variables and Other Methods	33
8.1 Separation of Variables	33
8.2 Applying Integral Transforms	33
8.3 Inhomogeneous Problems	33
Appendices	34

Part I

Complex Analysis

Chapter 1

Properties of Complex Functions

1.1.0 Elementary Definitions

1.2.0 Multivalued Functions and Branch Cuts

1.3.0 Analytic Functions and the Cauchy-Riemann Equations

1.4.0 Singularities and Zeros

1.5.0 Conformal Mappings

Chapter 2

Power Series and Laurent Series

2.1.0 Power Series Fundamentals

2.2.0 Laurent Series

2.3.0 Series Operations

Chapter 3

Complex Integration

3.1.0 Complex Integral

3.2.0 Cauchy's Theorems

3.3.0 Residue Theorem

3.4.0 Contour Integrals

Chapter 4

Applications of Complex Functions

4.1.0 Complex Potentials

4.2.0 Finding Zeros

4.3.0 Inverse Laplace

4.4.0 Stokes' Equations and Airy Integrals

4.5.0 WKB Methods, and Integral Approximations

Part II

PDEs

Chapter 5

General and Particular Solutions

5.1.0 Important Examples and Motivation

5.2.0 General Forms of Solutions

5.3.0 Wave and Diffusion Equations

5.4.0 Existence and Uniqueness

Chapter 6

Fourier Series

6.1.0 Initial Definitions and Dirichlet Conditions

Definition 6.1.1. *Sufficient conditions for which a function $f(x)$ to have its Fourier series to converge to it are known as the Dirichlet conditions:*

- (i) $f(x)$ must be periodic; i.e. there exists $p \in \mathbb{R}$ such that $f(x + p) = f(x)$ for all $x \in \mathbb{R}$.
- (ii) $f(x)$ must be continuous, except possibly at a finite number of jump (i.e. finite) discontinuities in any bounded interval.
- (iii) $f(x)$ must be of **bounded variation** on any bounded interval, which is to say its total variation is finite; if f is differentiable and its derivative is Riemann-integrable on the interval, then the total variation is the absolute integral of the derivative over the interval:

$$V_a^b(f) = \int_a^b |f'(x)| dx$$

An alternative formulation is to require that any bounded interval contains only a finite number of extrema of f .

- (iv) $f(x)$ is absolutely integrable over a period, so

$$\int_0^p |f(x)| dx < \infty$$

If these criteria hold, the Fourier series converges to $f(x)$ at all points where the function is continuous.

Recall that a function $f(x)$ can be split into an even and odd part:

$$f(x) = \frac{1}{2}[f(x) + f(-x)] + \frac{1}{2}[f(x) - f(-x)] = f_{\text{even}}(x) + f_{\text{odd}}(x)$$

Then, we can write the even component as a cosine series and the odd component as a sine series.

Proposition 6.1.1. For any $L \in \mathbb{R}$, the set of functions

$$\left\{ 1, \cos \frac{2\pi x}{L}, \sin \frac{2\pi x}{L}, \cos \frac{2\pi 2x}{L}, \sin \frac{2\pi 2x}{L}, \dots, \cos \frac{2\pi nx}{L}, \sin \frac{2\pi nx}{L}, \dots \right\}$$

form an orthogonal set with respect to the inner product

$$\langle f, g \rangle = \int_{x_0}^{x_0+L} f(x)g(x)dx$$

for $x_0 \in \mathbb{R}$ fixed. In particular, we have

$$\begin{aligned} \int_{x_0}^{x_0+L} \sin \frac{2\pi nx}{L} \cos \frac{2\pi mx}{L} dx &= 0, \quad \forall n, m \in \mathbb{N} \cup \{0\} \\ \int_{x_0}^{x_0+L} \cos \frac{2\pi nx}{L} \cos \frac{2\pi mx}{L} dx &= \begin{cases} L & \text{if } n = m = 0, \\ \frac{L}{2} & \text{if } n = m > 0, \\ 0 & \text{if } n \neq m \end{cases} \\ \int_{x_0}^{x_0+L} \sin \frac{2\pi nx}{L} \sin \frac{2\pi mx}{L} dx &= \begin{cases} 0 & \text{if } n = m = 0, \\ \frac{L}{2} & \text{if } n = m > 0, \\ 0 & \text{if } n \neq m \end{cases} \end{aligned}$$

Definition 6.1.2. The classical Fourier series expansion of a function $f(x)$ is

$$f(x) \sim \frac{a_0}{2} + \sum_{n=1}^{\infty} \left[a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right] \quad (6.1.1)$$

where a_0, a_n, b_n , for $n \geq 1$, are called the Fourier coefficients

For a periodic function $f(x)$ of period L , we use the orthogonality conditions to find the Fourier coefficients as follows:

$$a_0 = \frac{2}{L} \int_{x_0}^{x_0+L} f(x) \cdot 1 dx \quad (6.1.2)$$

$$a_n = \frac{2}{L} \int_{x_0}^{x_0+L} f(x) \cdot \cos \frac{2\pi nx}{L} dx \quad (6.1.3)$$

$$b_n = \frac{2}{L} \int_{x_0}^{x_0+L} f(x) \cdot \sin \frac{2\pi nx}{L} dx \quad (6.1.4)$$

where x_0 is arbitrary, but fixed, and $n \geq 1$.

Symmetry Conditions

From these coefficient equations we observe that if $f(x)$ is even with respect to the origin then all sine terms, b_n , are zero. Conversely, if $f(x)$ is odd with respect to the origin then all cosine terms, a_n , are zero. We now consider a more subtle symmetry about $L/4$, where L is a period of f , so $f(x+L) = f(x)$ for all $x \in \mathbb{R}$.

Definition 6.1.3. We say that $f(x)$ has even symmetry about $L/4$ if $f(L/4 - x) = f(x - L/4)$ for all $x \in \mathbb{R}$. We say that $f(x)$ has odd symmetry about $L/4$ if $f(L/4 - x) = -f(x - L/4)$.

We consider the sine terms of $g(x) = f(x - L/4)$, and substitute $s = x - L/4$:

$$\begin{aligned} b_n &= \frac{2}{L} \int_{x_0}^{x_0+L} f(x - L/4) \sin \frac{2\pi nx}{L} dx \\ &= \frac{2}{L} \int_{x_0-L/4}^{x_0-L/4+L} f(s) \sin \left[\frac{2\pi ns}{L} + \frac{\pi n}{2} \right] ds \\ &= \frac{2}{L} \int_{x_0}^{x_0+L} f(s) \sin \left[\frac{2\pi ns}{L} + \frac{\pi n}{2} \right] ds \end{aligned}$$

where the limits of integration can be changed since f is periodic. We observe that

$$\sin \left[\frac{2\pi ns}{L} + \frac{\pi n}{2} \right] = \sin \frac{2\pi ns}{L} \cos \frac{\pi n}{2} + \cos \frac{2\pi ns}{L} \sin \frac{\pi n}{2}$$

so the trigonometric portion of the integrand is odd if n is even and even if n is odd. Then if $f(s)$ is even and n is even the integral is zero, and similarly if $f(s)$ is odd and n is odd the integral is zero. For the cosine coefficients we have

$$\cos \left[\frac{2\pi ns}{L} + \frac{\pi n}{2} \right] = \cos \frac{2\pi ns}{L} \cos \frac{\pi n}{2} - \sin \frac{2\pi ns}{L} \sin \frac{\pi n}{2}$$

which is even if n is even and odd if n is odd. Then if $f(s)$ is even and n is odd, the terms a_n are zero, and if $f(s)$ is odd and n is even, the terms a_n are zero. In summary:

- If $f(x)$ is even about $L/4$, then $a_{2n-1} = 0$ and $b_{2n} = 0$ for all $n \geq 1$,
- If $f(x)$ is odd about $L/4$, then $a_{2n} = 0$ and $b_{2n+1} = 0$ for all $n \geq 0$.

6.2.0 Discontinuous and Non-Periodic Functions

Discontinuities

Definition 6.2.1. The error term for the Fourier series representation of f when expressed as a partial sum with highest term N is

$$E_N(x) = \left| f(x) - \left[\frac{a_0}{2} + \sum_{n=1}^N \left(a_n \cos \frac{2\pi nx}{L} + b_n \sin \frac{2\pi nx}{L} \right) \right] \right|$$

If $f(x)$ is discontinuous at a point a in the domain of interest, then the Fourier series for $f(x)$ does not produce a discontinuity at a but rather converges to the value

$$\frac{1}{2}[f(a+) + f(a-)]$$

where $f(a+) = \lim_{x \rightarrow a^+} f(x)$ is the one-sided limit from above, and $f(a-) = \lim_{x \rightarrow a^-} f(x)$ is the one-sided limit from below. Then, there exists sequences u_n and v_n such that $u_n, v_n \rightarrow a$, with $u_n < a$ for all n and $v_n > a$ for all n and

$$E_N(u_N) \approx 0.9|f(a-) - f(a+)| \quad E_N(v_N) \approx 0.9|f(a-) - f(a+)|$$

so the maximum value of the error $E_N(x)$ near a does not approach zero as $N \rightarrow \infty$, but rather occurs closer and closer to a , and is essentially independent of N . This is known as the **Gibbs' phenomenon**.

Non-Periodic Functions

We often wish to analyze non-periodic functions using Fourier analysis, and this can be done by using appropriate periodic extensions:

Theorem 6.2.1. Suppose h is differentiable on $[0, L]$; that is, $h'(x)$ exists for $0 < x < L$, and the one-sided derivatives

$$h'_+ = \lim_{x \rightarrow 0^+} \frac{h(x) - h(0)}{x} \quad \text{and} \quad h'_-(L) = \lim_{x \rightarrow L^-} \frac{h(x) - h(L)}{x - L}$$

both exist.

- Let O denote the odd periodic extension of h to $(-\infty, \infty)$ defined by

$$O(x) = \begin{cases} h(x), & 0 \leq x \leq L \\ -h(-x), & -L < x < 0, \end{cases} \quad \text{and } O(x + 2L) = O(x), \quad \forall x \in \mathbb{R}$$

Then O is differentiable on $(-\infty, \infty)$ if and only if

$$h(0) = h(L) = 0$$

- Let E denote the even periodic extension of h to $(-\infty, \infty)$, defined by

$$E(x) = \begin{cases} h(x), & 0 \leq x \leq L \\ h(-x), & -L < x < 0, \end{cases} \quad \text{and } E(x + 2L) = E(x), \quad \forall x \in \mathbb{R}$$

Then E is differentiable on $(-\infty, \infty)$ if and only if

$$h'_+(0) = h'_-(L) = 0$$

6.3.0 Integration and Differentiation

Theorem 6.3.1. If $f(x)$ satisfies the Dirichlet conditions, then integrating the Fourier series of $f(x)$ term by term produces a Fourier series which converges to the integral of $f(x)$, modulo an arbitrary constant.

Theorem 6.3.2. *If $f(x)$ satisfies the Dirichlet conditions, is differentiable, and $f'(x)$ satisfies the Dirichlet conditions, then the Fourier series obtained by differentiating f 's Fourier series term by term converges to $f'(x)$.*

For general functional series we have the following important result:

Theorem 6.3.3. *A convergent infinite series*

$$W(z) = \sum_{n=1}^{\infty} w_n(z)$$

can be differentiated term by term on a closed interval $[z_1, z_2]$ to obtain

$$W'(z) = \sum_{n=1}^{\infty} w'_n(z)$$

provided that w'_n is continuous on $[z_1, z_2]$ and there exists a sequence M_n of constants such that $\sum_{n=1}^{\infty} M_n$ converges and

$$|w'_n(z)| \leq M_n, \quad z_1 \leq z \leq z_2, \quad n = 1, 2, 3, \dots$$

6.4.0 Complex Fourier Series

Recall that by DeMoivre's Formula we have the following for the complex exponential:

$$\exp \{ix\} = \cos x + i \sin x$$

Definition 6.4.1. *For a function $f(x)$ of period L , its complex Fourier series expansion is given by*

$$f(x) \sim \sum_{n=-\infty}^{\infty} c_n \exp \left\{ \frac{2\pi i n x}{L} \right\}$$

We remark that the set of functions

$$\left\{ 1, \exp \left\{ \frac{2\pi i x}{L} \right\}, \exp \left\{ \frac{2\pi i 2x}{L} \right\}, \dots, \exp \left\{ \frac{2\pi i n x}{L} \right\}, \dots \right\}$$

forms an orthogonal set under the complex inner product

$$\langle f, g \rangle = \int_{x_0}^{x_0+L} f(x) \overline{g(x)} dx$$

for x_0 fixed with the relation

$$\int_{x_0}^{x_0+L} \exp \left\{ \frac{2\pi i n x}{L} \right\} \exp \left\{ -\frac{2\pi i m x}{L} \right\} dx = \begin{cases} L & \text{if } m = n \\ 0 & \text{if } m \neq n \end{cases}$$

Then the Fourier coefficients are given by

$$c_n = \frac{1}{L} \int_{x_0}^{x_0+L} f(x) \exp \left\{ -\frac{2\pi i n x}{L} \right\} dx \quad (6.4.1)$$

We expand the complex exponential in the Fourier series as follows:

$$\begin{aligned} \sum_{n=-\infty}^{\infty} c_n \exp \left\{ \frac{2\pi i n x}{L} \right\} &= \sum_{n=1}^{\infty} c_{-n} \exp \left\{ \frac{-2\pi i n x}{L} \right\} + c_0 + \sum_{n=1}^{\infty} c_n \exp \left\{ \frac{2\pi i n x}{L} \right\} \\ &= c_0 + \sum_{n=1}^{\infty} \left[c_{-n} \left(\cos \frac{2\pi n x}{L} - i \sin \frac{2\pi n x}{L} \right) + c_n \left(\cos \frac{2\pi n x}{L} - i \sin \frac{2\pi n x}{L} \right) \right] \\ &= c_0 + \sum_{n=1}^{\infty} \left[(c_{-n} + c_n) \cos \frac{2\pi n x}{L} + (i c_n - i c_{-n}) \sin \frac{2\pi n x}{L} \right] \end{aligned}$$

From this expansion we find that

$$\begin{aligned} c_0 &= \frac{a_0}{2} \\ c_{-n} + c_n &= a_n \\ i c_n - i c_{-n} &= b_n \end{aligned}$$

for $n \geq 1$. Then we have that

$$c_n = \frac{1}{2}(a_n - i b_n) \quad \text{and} \quad c_{-n} = \frac{1}{2}(a_n + i b_n)$$

It follows that if $f(x)$ is real, so a_n and b_n are real, $c_{-n} = \overline{c_n}$.

Parseval's Theorem

Theorem 1 (Parseval's Theorem).

Suppose that $A(x)$ and $B(x)$ are two complex valued functions on \mathbb{R} of period $2L$ that are square integrable with respect to the Lebesgue measure over intervals of period length with complex Fourier series

$$A(x) = \sum_{n=-\infty}^{\infty} a_n \exp \left\{ \frac{i\pi n x}{L} \right\}, \quad \text{and} \quad B(x) = \sum_{n=-\infty}^{\infty} b_n \exp \left\{ \frac{i\pi n x}{L} \right\}$$

Then

$$\sum_{n=-\infty}^{\infty} a_n \overline{b_n} = \frac{1}{2L} \int_{-L}^L A(x) \overline{B(x)} dx \quad (6.4.2)$$

Proof. Suppose $A(x)$ and $B(x)$ are as above, with corresponding Fourier series, and observe that

$$\frac{1}{2L} \int_{-L}^L A(x) \overline{B(x)} dx = \sum_{n=-\infty}^{\infty} a_n \frac{1}{2L} \int_{-L}^L \overline{B(x)} \exp \left\{ \frac{2\pi i n x}{2L} \right\} dx$$

$$\begin{aligned}
 &= \sum_{n=-\infty}^{\infty} a_n \left[\frac{1}{2L} \int_{-L}^L B(x) \exp \left\{ \frac{-2\pi i n x}{2L} \right\} dx \right] \\
 &= \sum_{n=-\infty}^{\infty} a_n \overline{b_n}
 \end{aligned}$$

as desired. ■

As a corollary, we have that for any function $f(x)$ of period L ,

$$\frac{1}{L} \int_{x_0}^{x_0+L} |f(x)|^2 dx = \sum_{n=-\infty}^{\infty} |c_n|^2 = \frac{|a_0|^2}{4} + \frac{1}{2} \sum_{n=1}^{\infty} (a_n^2 + b_n^2)$$

This says that the sum of the moduli squared of the complex Fourier coefficients is equal to the average value of $|f(x)|^2$ over one period.

Chapter 7

Integral Transforms

7.1.0 Fourier Transform

Construction and Definitions

Let $f(x)$ be defined on $-\infty < x < \infty$ such that it satisfies the Dirichlet conditions on $-L < x < L$. Then recall it has complex Fourier series expansion

$$f(x) = \sum_{n=-\infty}^{\infty} c_n \exp \left\{ \frac{in\pi x}{L} \right\} dx$$

with coefficients given by

$$c_n = \frac{1}{2L} \int_{-L}^L f(x) \exp \left\{ \frac{-in\pi x}{L} \right\} dx$$

We now define

$$\mu_n = \frac{n\pi}{L}, \quad n = 0, \pm 1, \pm 2, \dots$$

so we have

$$\Delta\mu_n = \mu_{n+1} - \mu_n = \frac{\pi}{L}$$

Note here we are constructing f with period $2L$. Then the coefficients can be rewritten as

$$c_n = \frac{\Delta\mu_n}{2\pi} \int_{-L}^L f(x) \exp \{-i\mu_n x\} dx$$

and we define

$$F_L(\mu) = \frac{1}{2\pi} \int_{-L}^L f(x) \exp \{-i\mu x\} dx$$

so we can write the series as

$$f(x) = \sum_{n=-\infty}^{\infty} F_L(\mu_n) \exp \{i\mu_n x\} \Delta\mu_n, \quad \text{and } c_n = F_L(\mu_n) \Delta\mu_n$$

In this form we take the limit as $L \rightarrow \infty$, which gives us the Riemann-integral

$$f(x) = \int_{-\infty}^{\infty} F(\mu) \exp \{i\mu x\} d\mu \quad (7.1.1)$$

and

$$F(\mu) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(x) \exp \{-i\mu x\} dx \quad (7.1.2)$$

Theorem 7.1.1. *Given a suitably integrable function $f(x)$ which satisfies the Dirichlet conditions, its **Fourier Inversion Formula** is given by:*

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} f(y) \exp \{-i\mu y\} dy \right] \exp \{i\mu x\} d\mu \quad (7.1.3)$$

The Fourier inversion formula and our previous results motivates the following definition:

Definition 7.1.1. *Given a function $f(x)$ suitably integrable, we define the **Fourier transform** of $f(x)$ to be*

$$F(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) \exp \{-iux\} dx \quad (7.1.4)$$

and consequently the function is given by

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} F(u) \exp \{iux\} du \quad (7.1.5)$$

We change the factor from $1/2\pi$ for one and 1 for the other integral to $1/\sqrt{2\pi}$ for both to emphasize symmetry.

The Uncertainty Principle

Definition 7.1.2. *We define the **dispersion about zero** of a complex-valued function $f(x)$, $-\infty < x < \infty$ by the formula*

$$D_0(f) = \frac{\int_{-\infty}^{\infty} x^2 |f(x)|^2 dx}{\int_{-\infty}^{\infty} |f(x)|^2 dx}$$

which is defined when the appropriate integrals are finite.

Proposition 7.1.2 (Uncertainty Principle). *Let $f(x)$, $-\infty < x < \infty$, be a complex-valued function with Fourier transform $F(\mu)$, for which the integrals defining $D_0(f)$ and $D_0(F)$ are finite. Then we have the inequality*

$$D_0(f)D_0(F) \geq \frac{1}{4} \quad (7.1.6)$$

Equality holds if and only if f is a normal density function centered at $x = 0$; in detail, $f(x) = C_1 e^{-x^2/2\sigma^2}$, $F(\mu) = C_2 e^{-\sigma^2\mu^2/2}$ for suitable constants C_1, C_2 , and σ^2 .

Proof. Both the numerator and denominator of the expressions defining $D_0(f)$ and $D_0(F)$ may be transformed by Parseval's Theorem. In this way we are led to examine a corresponding integral involving $F'(\mu)$. Specifically, we write the real part of the integral of $\mu \bar{F} F'$ in two ways:

$$2\mathbf{Re} \int \mu \bar{F} F' d\mu = \int \mu (F' \bar{F} + F \bar{F}') d\mu = \int \mu (F \bar{F})' d\mu = \int \mu (|F|^2)' d\mu = - \int |F|^2 d\mu$$

using integration by parts, and

$$-\mathbf{Re} \int \mu \bar{F} F' d\mu \leq \left| \int \mu \bar{F}(\mu) F'(\mu) d\mu \right| \leq \left(\int |\mu \bar{F}(\mu)|^2 d\mu \right)^{1/2} \left(\int |F'(\mu)|^2 d\mu \right)^{1/2}$$

applying the Schwarz inequality. Now we apply Parseval's theorem twice, and use the fact that the Fourier transform of $xf(x)$ is $iF'(\mu)$:

$$\int |F(\mu)|^2 d\mu = \frac{1}{2\pi} \int |f(x)|^2 dx, \quad \int |F'(\mu)|^2 d\mu = \frac{1}{2\pi} \int x^2 |f(x)|^2 dx$$

Squaring both sides of the Schwarz inequality and making these substitutions gives

$$\frac{1}{4} \int |f|^2 \int |F|^2 \leq \int |xf|^2 dx \int |\mu F|^2$$

which gives the desired inequality.

In case of equality we require that the imaginary part of $\int \mu \bar{F} F' d\mu$ is zero, and F must satisfy the differential equation $F'(\mu) = -A\mu F(\mu)$ for some complex constant A . ■

This implies that we cannot localize both $f(x)$ and $F(\mu)$ in their respective spaces. If $f(x)$ is localized about $x = 0$, then $D_0(f)$ will be small; the uncertainty principle then asserts that $D_0(F)$ must be correspondingly large, indicating a lack of localization about $\mu = 0$.

Now, we consider the normalized Gaussian distribution centered at zero:

$$f(t) = \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{t^2}{2\sigma^2} \right\}, \quad -\infty < t < \infty$$

Then the Fourier transform of $f(t)$ is

$$\begin{aligned} F(u) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{t^2}{2\sigma^2} \right\} \exp \{ -iut \} dt \\ &= \frac{1}{2\pi\sigma} \int_{-\infty}^{\infty} \exp \left\{ -\frac{(t + iu\sigma^2)^2}{2\sigma^2} - \frac{u^2\sigma^2}{2} \right\} dt \\ &= \frac{\exp \left\{ -\frac{u^2\sigma^2}{2} \right\}}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{(t + iu\sigma^2)^2}{2\sigma^2} \right\} dt \\ &= \frac{1}{\sqrt{2\pi}} \exp \left\{ -\frac{u^2\sigma^2}{2} \right\} \end{aligned}$$

Thus, $F(u)$ is another Gaussian distribution centered at zero with standard deviation $1/\sigma$. Hence, the spreads in t and u are inversely related, and independent of the actual value of σ . If t represents time, u represents frequency, and we find that the narrower a time measurement is, the greater the spread of frequency components it must contain.

Note 7.1.1. Note that the fourier transform of a function $f(t)$ of time t is a function $F(u)$ of angular frequency, $u = 2\pi/T$.

Recall the de Broglie and Einstein relationships for momentum and energy:

$$p = \hbar k \quad \text{and} \quad E = \hbar \omega$$

where $\hbar = h/2\pi$, for Planck's constant h . If $f(t)$ is the wave function, the distribution of the wave intensity in time is given by $|f|^2$, which is Gaussian. Similarly, the intensity distribution in frequency is given by $|F|^2$. These distributions have respective standard deviations $\sigma/\sqrt{2}$ and $1/(\sqrt{2}\sigma)$, given

$$\Delta E \Delta t = \frac{\hbar}{2} \quad \text{and} \quad \Delta p \Delta x = \frac{\hbar}{2}$$

In general, the equalities are inequalities with $\geq \frac{\hbar}{2}$ if the distributions are possibly not Gaussian.

Dirac δ -function

Definition 7.1.3. Rigorously the Dirac δ -function is a distribution or measure, not a function. But, simplistically it can be thought of a function such that

$$\delta(t) = \begin{cases} +\infty & t = 0 \\ 0 & t \neq 0 \end{cases}$$

with the property that

$$\int_{-\infty}^{\infty} \delta(t) dt = 1$$

or in general

$$\int_{-\infty}^{\infty} f(t) \delta(t - a) dt = f(a) \quad (7.1.7)$$

which is also valid as long as a is contained in the bounds of integration, and is zero otherwise.

Proposition 7.1.3. The δ function is even, absolutely inverse homogeneous, and trivial upon multiplication by t :

$$\delta(t) = \delta(-t)$$

$$\delta(at) = \frac{1}{|a|} \delta(t)$$

$$t\delta(t) = 0$$

Proof. First, $\delta(t) = \delta(-t)$ is immediate for all t . Then, suppose $a > 0$. It follows that

$$\int_{-\infty}^{\infty} \delta(at) dt = \int_{-\infty}^{\infty} \delta(x) \frac{dx}{a} = \frac{1}{a}$$

If $a < 0$ then we have

$$\int_{-\infty}^{\infty} \delta(at) dt = \int_{\infty}^{-\infty} \delta(x) \frac{dx}{a} = -\frac{1}{a}$$

In either case we find the integral is equal to $1/|a|$, so $\delta(at) = \frac{1}{|a|} \delta(t)$. ■

Proposition 7.1.4. *If g is a continuously differentiable function with g' nowhere zero, then if g is also nowhere zero*

$$\delta(g(x)) = 0$$

Otherwise, if g has zeros x_i , then

$$\delta(g(x)) = \sum_i \frac{\delta(x - x_i)}{|g'(x_i)|}$$

Proposition 7.1.5. *The k th distributional derivative of the delta function, $\delta^{(k)}(t)$, is defined by*

$$\int_{-\infty}^{\infty} f(t) \delta^{(k)}(t) dt = (-1)^k f^{(k)}(0)$$

Definition 7.1.4. *The delta function generalized to n -dimensional euclidean space gives*

$$\int_{\mathbb{R}^n} f(\mathbf{x}) \delta(\mathbf{x}) d\mathbf{x} = f(\mathbf{0})$$

As a probability measure on \mathbb{R} , the delta measure has cumulative distribution function equal to the unit step function, or **Heaviside function**:

$$H(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases}$$

so we can loosely see $H'(t) = \delta(t)$.

Returning to the Fourier transform, we note that we can write the Fourier inversion formula as

$$\begin{aligned} f(t) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} f(x) \exp\{-iux\} dx \right] \exp\{iut\} du \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} f(x) \left[\int_{-\infty}^{\infty} \exp\{iu(t-x)\} du \right] dx \end{aligned}$$

Comparison of this with the δ function's property

$$\int_{-\infty}^{\infty} f(x) \delta(t-x) dx = f(t)$$

This implies that we can write

$$\boxed{\delta(t-x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \exp\{iu(t-x)\} du} \quad (7.1.8)$$

This formulation implies that we can interpret the dirac δ -function as having a very narrow peak at $t = x$ resulting from the superposition of a complete spectrum of harmonic waves, all frequencies having the same amplitude and all waves being in phase at $t = x$.

Now, consider a uniform distribution of unit height extending from $-\Omega$ to Ω , considered as the Fourier transform of some function f_Ω . Then we take the inverse Fourier transform to obtain

$$\begin{aligned} f_\Omega(t) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} F_\Omega(u) \exp\{iut\} du \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\Omega}^{\Omega} \exp\{iut\} du \\ &= \frac{1}{\sqrt{2\pi}} \frac{1}{it} \exp\{iut\} \Big|_{u=-\Omega}^{u=\Omega} \\ &= \frac{2\Omega}{\sqrt{2\pi}} \frac{\sin \Omega t}{\Omega t} \end{aligned}$$

In the limit as $\Omega \rightarrow \infty$, $f_\Omega(t)$ as defined by the inverse Fourier transform tends to $\sqrt{2\pi}\delta(t)$ by nature of our realization of the delta function from the Fourier inversion formula. Hence, we may conclude that the δ function can also be represented by

$$\delta(t) = \lim_{\Omega \rightarrow \infty} \frac{\sin \Omega t}{\pi t}$$

Finally, the Fourier transform of a δ function is simply

$$F_\delta(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \delta(t) \exp\{-iut\} dt = \frac{1}{\sqrt{2\pi}}$$

Properties of Fourier Transforms

The following theorem gives conditions for the existence of the integrals in the Fourier inversion formula and convergence of the Fourier transform:

Theorem 7.1.6. *Let $f(x)$, $-\infty < x < \infty$, be a piecewise smooth function on each finite interval such that $\int_{-\infty}^{\infty} |f(x)| dx < \infty$. Define the Fourier transform $F(\mu)$ as above. Then for each x ,*

$$\lim_{M \rightarrow \infty} \int_{-M}^M F(\mu) \exp\{i\mu x\} d\mu = \frac{1}{2}[f(x+) + f(x-)]$$

Note that this limit is a **Cauchy principal value**.

Proposition 7.1.7. *Let $f(x)$ be a function and let \mathcal{F} denote the Fourier transform, so in our previous notation $\mathcal{F}[f(x)](\mu) = F(\mu)$. Then \mathcal{F} satisfies the following properties:*

- *If f is differentiable, such that $f'(x)$ is absolutely integrable on the real line, then we have that*

$$\mathcal{F}[f'(x)](\mu) = i\mu \mathcal{F}[f(x)](\mu) \quad (7.1.9)$$

In general, if $f(x)$ is n times differentiable such that $f^{(n)}$ is absolutely integrable on the real line, then

$$\mathcal{F}[f^{(n)}(x)](\mu) = i^n \mu^n \mathcal{F}[f(x)](\mu) \quad (7.1.10)$$

- If $f(x)$ is absolutely integrable on the real line, we have

$$\mathcal{F}\left[\int_{-\infty}^t f(x)dx\right](\mu) = \frac{1}{i\mu} \mathcal{F}[f(x)](\mu) + 2\pi c\delta(\mu) \quad (7.1.11)$$

where the term $2\pi c\delta(\mu)$ represents the Fourier transform of the constant of integration associated with the indefinite integral.

- For any complex number $a \in \mathbb{C}^\times$, we have

$$\mathcal{F}[f(at)](\mu) = \frac{1}{a} \mathcal{F}[f(t)](\mu/a) \quad (7.1.12)$$

- For any complex number $a \in \mathbb{C}^\times$ we have

$$\mathcal{F}[f(t+a)](\mu) = \exp\{ia\mu\} \mathcal{F}[f(t)](\mu) \quad (7.1.13)$$

- For any $\alpha, \beta \in \mathbb{C}$, and g also absolutely integrable on the real line, we have

$$\mathcal{F}[\alpha f(t) + \beta g(t)](\mu) = \alpha \mathcal{F}[f(t)](\mu) + \beta \mathcal{F}[g(t)](\mu) \quad (7.1.14)$$

- For any $\alpha \in \mathbb{C}$, we have that

$$\mathcal{F}[\exp\{\alpha t\} f(t)](\mu) = \mathcal{F}[f(t)](\mu + i\alpha) \quad (7.1.15)$$

Odd and Even Functions

First consider an odd function $f(t) = -f(-t)$, and analyze its Fourier transform $F(\mu)$:

$$\begin{aligned} F(\mu) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(t) \exp\{-i\mu t\} dt \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(t) (\cos \mu t - i \sin \mu t) dt \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} -if(t) \sin \mu t dt \\ &= \frac{-2i}{\sqrt{2\pi}} \int_0^{\infty} f(t) \sin \mu t dt \end{aligned}$$

where an integral even about the center of an odd function is 0, and an integral even about the center of an even function is two times half the integral. Further we observe that when $f(t)$ is odd, $F(\mu)$ is also odd, so $F(\mu) = -F(-\mu)$. Hence,

$$f(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} F(\mu) \exp\{i\mu t\} d\mu = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} F(\mu) (\cos \mu t + i \sin \mu t) d\mu$$

$$\begin{aligned}
 &= \frac{2i}{\sqrt{2\pi}} \int_0^\infty F(\mu) \sin \mu t d\mu \\
 &= \frac{2i}{\sqrt{2\pi}} \int_0^\infty \left[\frac{-2i}{\sqrt{2\pi}} \int_0^\infty f(x) \sin \mu x dx \right] \sin \mu t d\mu \\
 &= \frac{2}{\pi} \int_0^\infty \left[\int_0^\infty f(x) \sin \mu x dx \right] \sin \mu t d\mu
 \end{aligned}$$

Definition 7.1.5. If $f(x)$ is odd and absolutely integrable over the real line, we define the Fourier sine transform pair to be:

$$F_s(\mu) = \sqrt{\frac{2}{\pi}} \int_0^\infty f(t) \sin \mu t dt \quad (7.1.16)$$

$$f(t) = \sqrt{\frac{2}{\pi}} \int_0^\infty F_s(\mu) \sin \mu t d\mu \quad (7.1.17)$$

Next, consider an even function $f(t) = f(-t)$, and analyze its Fourier transform $F(\mu)$:

$$\begin{aligned}
 F(\mu) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty f(t) \exp \{-i\mu t\} dt \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty f(t) (\cos \mu t - i \sin \mu t) dt \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty f(t) \cos \mu t dt \\
 &= \frac{2}{\sqrt{2\pi}} \int_0^\infty f(t) \cos \mu t dt
 \end{aligned}$$

using properties of even and odd functions, as in the odd case. Moreover, we observe that when $f(t)$ is even, $F(\mu)$ is also even, so $F(\mu) = F(-\mu)$. Hence,

$$\begin{aligned}
 f(t) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty F(\mu) \exp \{i\mu t\} d\mu = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty F(\mu) (\cos \mu t + i \sin \mu t) d\mu \\
 &= \frac{2}{\sqrt{2\pi}} \int_0^\infty F(\mu) \cos \mu t d\mu \\
 &= \frac{2}{\sqrt{2\pi}} \int_0^\infty \left[\frac{2}{\sqrt{2\pi}} \int_0^\infty f(x) \cos \mu x dx \right] \cos \mu t d\mu \\
 &= \frac{2}{\pi} \int_0^\infty \left[\int_0^\infty f(x) \cos \mu x dx \right] \cos \mu t d\mu
 \end{aligned}$$

Definition 7.1.6. If $f(x)$ is even and absolutely integrable over the real line, we define the Fourier cosine transform pair to be:

$$F_s(\mu) = \sqrt{\frac{2}{\pi}} \int_0^\infty f(t) \cos \mu t dt \quad (7.1.18)$$

$$f(t) = \sqrt{\frac{2}{\pi}} \int_0^\infty F_s(\mu) \cos \mu t d\mu \quad (7.1.19)$$

Convolution and Deconvolution

When performing a physical measurement, we are always limited by the finite resolution of our measuring apparatus. Let $f(x)$ represent the physical quantity we wish to measure which is a function of the independent variable x . Then, as mentioned, we have a resolution function $g(y)$, which is some form of distribution, where the probability that an output value y will be recorded instead as between y and $y + dy$ is $g(y)dy$. Ideally, we want $g(y)$ to be as close to a δ -function as possible, centered at the true value.

Given the true distribution $f(x)$ and the resolution function of our apparatus $g(y)$, we wish to calculate what the observed distribution $h(z)$ will be. The probability that a true reading lying between x and $x + dx$, which is of probability $f(x)dx$ of being selected by the experiment, will be moved by the instrumental resolution to an amount $z - x$ into a small interval of width dz is $g(z - x)dz$. Hence, the combined probability that the interval dx will give rise to an observation appearing in the interval dz is $f(x)dxg(z - x)dz$. We add together the contributions from all values of x that can lead to an observation in the range z to $z + dz$, and we find that the observed distribution is given by

$$h(z) = \int_{-\infty}^{\infty} f(x)g(z - x)dx$$

This integral is a specific case of the generalized notion of a convolution:

Definition 7.1.7. If f and g are functions absolutely integrable over the entire real line, we define their convolution to be

$$(f * g)(u) = \int_{-\infty}^{\infty} f(x)g(u - x)dx \quad (7.1.20)$$

Proposition 7.1.8. For any functions f and g for which the convolution $f * g$ is defined, we have that $*$ is commutative, associative, and distributive over addition.

We now consider the Fourier transform of the convolution $f * g$:

$$\begin{aligned} \mathcal{F}[(f * g)(t)](\mu) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (f * g)(t) \exp\{-i\mu t\} dt \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} f(x)g(t - x)dx \right] \exp\{-i\mu t\} dt \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) \left[\int_{-\infty}^{\infty} g(t - x) \exp\{-i\mu t\} dt \right] dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) \left[\int_{-\infty}^{\infty} g(u) \exp\{-i\mu(u + x)\} du \right] dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) \sqrt{2\pi} \mathcal{F}[g(u)](\mu) \exp\{-i\mu x\} dx \\ &= \sqrt{2\pi} \mathcal{F}[f(u)](\mu) \mathcal{F}[g(u)](\mu) \end{aligned}$$

Similarly, for the Fourier transform of the product we have:

$$\begin{aligned}
 \mathcal{F}[f(t)g(t)](\mu) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(t)g(t) \exp\{-i\mu t\} dt \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \left[\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \mathcal{F}[f(t)](u) \exp\{iut\} du \right] g(t) \exp\{-i\mu t\} dt \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \mathcal{F}[f(t)](u) \left[\int_{-\infty}^{\infty} g(t) \exp\{-i(\mu - u)t\} dt \right] du \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \mathcal{F}[f(t)](u) \mathcal{F}[g(t)](\mu - u) du \\
 &= \frac{1}{\sqrt{2\pi}} (\mathcal{F}[f(t)] * \mathcal{F}[g(t)])(\mu)
 \end{aligned}$$

Thus, we obtain the following result:

Theorem 7.1.9 (Convolution Theorem). *If f and g are absolutely integrable over the real line, we have that*

$$\boxed{\mathcal{F}[(f * g)(u)](\mu) = \sqrt{2\pi} \mathcal{F}[f(u)](\mu) \mathcal{F}[g(u)](\mu)} \quad (7.1.21)$$

and

$$\boxed{\mathcal{F}[f(t)g(t)](\mu) = \frac{1}{\sqrt{2\pi}} (\mathcal{F}[f(t)] * \mathcal{F}[g(t)])(\mu)} \quad (7.1.22)$$

To find the true distribution $f(x)$ given an observed distribution $h(z)$ and a resolution function $g(y)$ we use **deconvolution**: We observe that $\mathcal{F}[(f * g)(t)](\mu) = \sqrt{2\pi} \mathcal{F}[f(t)](\mu) \mathcal{F}[g(t)](\mu)$. Then it follows that

$$\mathcal{F}[f(t)](\mu) = \frac{1}{\sqrt{2\pi}} \frac{\mathcal{F}[(f * g)(t)](\mu)}{\mathcal{F}[g(t)](\mu)}$$

so taking the inverse Fourier transform we find

$$f(t) = \frac{1}{\sqrt{2\pi}} \mathcal{F}^{-1} \left[\frac{\mathcal{F}[(f * g)(t)](\mu)}{\mathcal{F}[g(t)](\mu)} \right] = \frac{1}{\sqrt{2\pi}} \left[\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\mathcal{F}[(f * g)(t)](\mu)}{\mathcal{F}[g(t)](\mu)} \exp\{i\mu t\} d\mu \right] \quad (7.1.23)$$

Correlation Functions and Energy Spectra

Definition 7.1.8. The cross-correlation of two functions f and g is defined by

$$C(z) := \int_{-\infty}^{\infty} \overline{f(x)} g(x + z) dx \quad (7.1.24)$$

The cross-correlation provides a quantitative measure of the similarity of two functions f and g as one is displaced through a distance z relative to the other. We shall use the notation $C = f \oplus g$. \oplus is both associative and distributive, but not commutative, and in fact

$$(f \oplus g)(z) = \overline{(g \oplus f)(-z)} \quad (7.1.25)$$

Theorem 7.1.10 (Wiener-Kinchin Theorem). *For functions f and g we have that*

$$\mathcal{F}[(f \oplus g)(t)](\mu) = \sqrt{2\pi} \overline{\mathcal{F}[f(t)](\mu)} \mathcal{F}[g(t)](\mu)$$

Proof. Observe that

$$\begin{aligned} \mathcal{F}[(f \oplus g)(t)](\mu) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (f \oplus g)(t) \exp\{-i\mu t\} dt \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} \overline{f(x)} g(x+t) dx \right] \exp\{-i\mu t\} dt \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \overline{f(x)} \left[\int_{-\infty}^{\infty} g(x+t) \exp\{-i\mu t\} dt \right] dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \overline{f(x)} \left[\int_{-\infty}^{\infty} g(u) \exp\{-i\mu(u-x)\} du \right] dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \overline{f(x)} \frac{1}{\sqrt{2\pi}} \mathcal{F}[g(t)](\mu) \exp\{i\mu x\} dx \\ &= \frac{1}{\sqrt{2\pi}} \overline{\mathcal{F}[f(t)](\mu)} \mathcal{F}[g(t)](\mu) \end{aligned}$$

as desired. ■

Similarly, we can derive the converge theorem

$$\mathcal{F}[\overline{f(x)g(x)}](\mu) = \frac{1}{\sqrt{2\pi}} (\mathcal{F}[f(t)] \oplus \mathcal{F}[g(t)])(\mu) \quad (7.1.26)$$

Definition 7.1.9. The auto-correlation function of $f(x)$ is defined to be

$$(f \oplus f)(t) = \int_{-\infty}^{\infty} \overline{f(x)} f(x+t) dx \quad (7.1.27)$$

Using the Wiener-Kinchin theorem we see that

$$\begin{aligned} (f \oplus f)(t) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \mathcal{F}[(f \oplus f)(t)](\mu) \exp\{i\mu t\} d\mu \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \sqrt{2\pi} \overline{\mathcal{F}[f(t)](\mu)} \mathcal{F}[f(t)](\mu) \exp\{i\mu t\} d\mu \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \sqrt{2\pi} |\mathcal{F}[f(t)](\mu)|^2 \exp\{i\mu t\} d\mu \end{aligned}$$

so that $(f \oplus f)(t)$ is the inverse Fourier transform of $\sqrt{2\pi} |\mathcal{F}[f(t)](\mu)|^2$, which is called the energy spectrum of f .

Parseval's Theorem

From our above result, we have that for functions f and g

$$(f \oplus g)(t) = \int_{-\infty}^{\infty} \overline{f(x)} g(x+t) dx = \int_{-\infty}^{\infty} \overline{\mathcal{F}[f(t)](\mu)} \mathcal{F}[g(t)](\mu) \exp\{i\mu t\} d\mu$$

This gives the **multiplication theorem** when t is set to 0:

Theorem 7.1.11 (Multiplication Theorem). *Let f and g be absolutely integrable on the real line. Then*

$$\int_{-\infty}^{\infty} \overline{f(x)} g(x) dx = \int_{-\infty}^{\infty} \overline{\mathcal{F}[f(t)](\mu)} \mathcal{F}[g(t)](\mu) d\mu \quad (7.1.28)$$

By letting $g = f$ we obtain Parseval's Theorem:

Theorem 2 (Parseval's Theorem).

Let f be absolutely integrable, then

$$\int_{-\infty}^{\infty} |f(x)|^2 dx = \int_{-\infty}^{\infty} |\mathcal{F}[f(t)](\mu)|^2 d\mu \quad (7.1.29)$$

7.2.0 Multidimensional Fourier Transform

Definition 7.2.1. *In three dimensions we may define the Fourier transform of a function which is absolutely integrable over \mathbb{R}^3 by*

$$\mathcal{F}[f(x, y, z)](\mu_x, \mu_y, \mu_z) = \frac{1}{\sqrt{2\pi^3}} \int_{\mathbb{R}^3} f(x, y, z) \exp\{-i\mu_x x\} \exp\{-i\mu_y y\} \exp\{-i\mu_z z\} dx dy dz \quad (7.2.1)$$

with inverse

$$f(x, y, z) = \frac{1}{\sqrt{2\pi^3}} \int_{\mathbb{R}^3} \mathcal{F}[f(x, y, z)](\mu_x, \mu_y, \mu_z) \exp\{i\mu_x x\} \exp\{i\mu_y y\} \exp\{i\mu_z z\} d\mu_x d\mu_y d\mu_z \quad (7.2.2)$$

Definition 7.2.2. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be an absolutely integrable scalar field. Denote (x_1, x_2, \dots, x_n) by \mathbf{x} and $(\mu_{x_1}, \mu_{x_2}, \dots, \mu_{x_n})$ by μ . Then*

$$\mathcal{F}[f(\mathbf{x})](\mu) = \frac{1}{\sqrt{2\pi^n}} \int_{\mathbb{R}^n} f(\mathbf{x}) \exp\{-i\mu \cdot \mathbf{x}\} d\mathbf{x} \quad (7.2.3)$$

and

$$f(\mathbf{x}) = \frac{1}{\sqrt{2\pi^n}} \int_{\mathbb{R}^n} \mathcal{F}[f(\mathbf{x})](\mu) \exp\{i\mu \cdot \mathbf{x}\} d\mu \quad (7.2.4)$$

From these definitions and the Fourier inverse theorem we obtain the n -dimensional dirac δ function:

$$\delta(\mathbf{x}) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \exp\{i\mu \cdot \mathbf{x}\} d\mu \quad (7.2.5)$$

Suppose in three dimension $f(\mathbf{r})$ is spherically symmetric, so $f(\mathbf{r}) = f(r)$, where $r = |\mathbf{r}|$. Since f is spherically symmetric we can choose the vector μ of the Fourier transform to lie along the polar axis $\theta = 0$ without loss of generality. Then we have the change of volume differentials:

$$d\mathbf{r} = r^2 \sin \theta dr d\theta d\phi, \quad \text{and} \quad \mu \cdot \mathbf{r} = \mu r \cos \theta \quad (7.2.6)$$

where $\mu = |\mu|$. The Fourier transform is then given by

$$\begin{aligned} \mathcal{F}[f(r)](\mu) &= \frac{1}{\sqrt{2\pi^3}} \int_{\mathbb{R}^3} f(r) \exp\{-i\mu \cdot \mathbf{r}\} d\mathbf{r} \\ &= \frac{1}{\sqrt{2\pi^3}} \int_0^\infty dr \int_0^\pi d\theta \int_0^{2\pi} d\phi f(r) r^2 \sin \theta \exp\{-i\mu r \cos \theta\} \\ &= \frac{1}{\sqrt{2\pi^3}} \int_0^\infty dr 2\pi f(r) r^2 \int_0^\pi d\theta \sin \theta \exp\{-i\mu r \cos \theta\} \\ &= \frac{1}{\sqrt{2\pi^3}} \int_0^\infty dr 2\pi f(r) r^2 \frac{1}{\mu r} 2 \sin(\mu r) \\ &= \frac{1}{\sqrt{2\pi^3}} \int_0^\infty dr 4\pi f(r) r^2 \frac{\sin(\mu r)}{\mu r} \end{aligned}$$

7.3.0 Laplace Transform

The Laplace transform is an alternative to the Fourier transform, and applies to functions which aren't necessarily absolutely integrable over the real line. Moreover, we be interested in a function for a positive time, $t > 0$, and hence only place restrictions on the function in such a region:

Definition 7.3.1. Let $f(t)$ be some real function. Then the Laplace transform of f is defined to be

$$\mathcal{L}[f(t)](s) = \int_0^\infty f(t) e^{-st} dt \quad (7.3.1)$$

provided the integral exists.

Definition 7.3.2. A function f is said to be of exponential order s_0 if there are constants M and t_0 such that

$$|f(t)| \leq M e^{s_0 t}, \quad t \geq t_0$$

Theorem 7.3.1. If f is piecewise continuous on $[0, \infty)$ and of exponential order s_0 , then $\mathcal{L}[f(t)](s)$ is defined for $s > s_0$.

This theorem provides sufficient, but not necessary conditions, for f to have a Laplace transform.

Properties 7.3.3. The Laplace transforms of functions f, g_1, \dots, g_n satisfy the following properties, where $a_1, \dots, a_n \in \mathbb{R}$ are constants:

- (Linearity) The Laplace transform is linear

$$\mathcal{L}\left[\sum_{i=1}^n a_i g_i(t)\right](s) = \sum_{i=1}^n a_i \mathcal{L}[g_i(t)](s) \quad (7.3.2)$$

Consequently, the inverse Laplace transform is also linear.

- The Laplace transform of a derivative of a function is

$$\mathcal{L}\left[\frac{df}{dt}\right](s) = s\mathcal{L}[f(t)](s) - f(0), \quad s > 0 \quad (7.3.3)$$

and in general we have

$$\mathcal{L}[f^{(n)}(t)](s) = s^n \mathcal{L}[f(t)](s) - \sum_{i=0}^{n-1} s^{n-1-i} f^{(i)}(0), \quad s > 0 \quad (7.3.4)$$

- The integral of a Fourier transform is given by

$$\mathcal{L}\left[\int_0^t f(u)du\right](s) = \frac{1}{s} \mathcal{L}[f(t)](s) \quad (7.3.5)$$

- (Shift I) for $a \in \mathbb{R}$ we have that

$$\mathcal{L}[e^{at}f(t)](s) = \mathcal{L}[f(t)](s-a) \quad (7.3.6)$$

- (Shift II) for the unit step function $u_b(t) = 1, t > b, 0$ otherwise, for $b \in \mathbb{R}$, we have

$$\mathcal{L}[u_b(t)f(t)](s) = e^{-bt} \mathcal{L}[f(t+b)](s) \quad (7.3.7)$$

and

$$e^{-bt} \mathcal{L}[f(t)](s) = \mathcal{L}[u_b(t)f(t-b)](s) \quad (7.3.8)$$

- For any $a \in \mathbb{R}, a \neq 0$, we have that

$$\mathcal{L}[f(at)](s) = \frac{1}{a} \mathcal{L}[f(t)](s/a) \quad (7.3.9)$$

- For any $n \in \mathbb{N}$, we have that

$$\mathcal{L}[t^n f(t)](s) = (-1)^n \frac{d^n}{ds^n} \mathcal{L}[f(t)](s) \quad (7.3.10)$$

- If $\lim_{t \rightarrow 0} [f(t)/t]$ exists, then we have

$$\mathcal{L}\left[\frac{f(t)}{t}\right](s) = \int_s^\infty \mathcal{L}[f(t)](u)du \quad (7.3.11)$$

- The Laplace transform of the convolution is given by

$$\mathcal{L}[(f * g)(t)](s) = \mathcal{L}[f(t)](s) \mathcal{L}[g(t)](s) \quad (7.3.12)$$

There is no general inverse Laplace transform, unlike with the Fourier transform, so often we rely on our identities for the Laplace transform of common functions to compute inverses. We have the following table of standard Laplace transforms:

Table 7.1: Common Laplace Transforms

$f(t)$	$\mathcal{L}[f(t)](s)$	s_0
c	$\frac{c}{s}$	0
ct^n	$\frac{cn!}{s^{n+1}}$	0
$\sin bt$	$\frac{b}{(s^2+b^2)}$	0
$\cos bt$	$\frac{s}{(s^2+b^2)}$	0
e^{at}	$\frac{1}{(s-a)}$	a
$t^n e^{at}$	$\frac{n!}{(s-a)^{n+1}}$	a
$\sinh at$	$\frac{a}{(s^2-a^2)}$	$ a $
$\cosh at$	$\frac{s}{(s^2-a^2)}$	$ a $
$e^{at} \sin bt$	$\frac{b}{[(s-a)^2+b^2]}$	a
$e^{at} \cos bt$	$\frac{s-a}{[(s-a)^2+b^2]}$	a
$t^{1/2}$	$\frac{\sqrt{\pi}}{2\sqrt{s^3}}$	0
$t^{-1/2}$	$\frac{\sqrt{\pi}}{\sqrt{s}}$	0
$\delta(t-t_0)$	e^{-st_0}	0
$H(t-t_0) = \begin{cases} 1 & \text{if } t \geq t_0 \\ 0 & \text{if } t < t_0 \end{cases}$	$\frac{e^{-st_0}}{s}$	0

General Transform

Definition 7.3.4. A general integral transform of a function $f(t)$ is of the form

$$F(\alpha) = \int_a^b K(\alpha, t) f(t) dt \quad (7.3.13)$$

where $F(\alpha)$ is the transform of $f(t)$ with respect to the kernel $K(\alpha, t)$, and α is the transform variable.

Often the inverse transform is itself an integral transform, forming a transform pair - the Fourier transform is one such example, and here are two others:

- The Hankel transform:

$$F(k) = \int_0^\infty f(x) J_n(kx) x dx$$

$$f(x) = \int_0^\infty F(k) J_n(kx) k dk$$

where the J_n are Bessel functions of order n ,

- The *Mellin Transform*

$$F(z) = \int_0^{\infty} t^{z-1} f(t) dt$$
$$f(t) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} t^{-z} F(z) dz$$

Chapter 8

Separation of Variables and Other Methods

8.1.0 Separation of Variables

8.2.0 Applying Integral Transforms

8.3.0 Inhomogeneous Problems

Appendices