

And so begins the actual work towards my thesis

Kristaps J. Balodis

July 13, 2022

Contents

Contents	1
1 The yoga of multi-segments	3
1.1 The spectral multiplicity matrix	3
2 The geometry of $GL(n)$	7
2.1 The reduction of Vogan varieties of $GL(n)$	7
2.2 Explicit descriptions on V_λ in extremal cases	12
3 The book of $GL(4)$	19
3.1 The determinantal case	20
3.2 An "unresolved" case: Zelevinsky's 11.4	25
3.3 An explicit description of a supercuspidal representation of $GL(2)$	33
4 The groups $SO(n)$	35
5 The Kazhdan-Lusztig hypothesis for affine Hecke algebras	37
5.1 Chapter 8 of C+G	37
5.2 The final pieces	39
5.3 Speculation towards the full proof	39
Bibliography	41

The main idea behind this document, is that in order to compute either the spectral or geometric multiplicity matrices for $GL_n(F)$, it suffices to compute the multiplicity matrices "run-by-run",

which will be explained in the following section. The proof of this on the spectral side is easily dispatched by a theorem of Zelevinsky [11]. On the geometric side, considerably more work is required.

In the respective sections, after establishing the main result, we turn to an analysis of extremal cases, being those of multiplicity free support, and of "complete multiplicity".

Chapter 1

The yoga of multi-segments

1.1 The spectral multiplicity matrix

In this section, we rely on the theory of multi-segments as it appear in [11] to prove the main result, and explore some examples.

Spectral a-ray-ngments

In this section, we will show that given some support S of supercuspidal representations, one can group together the representations based on which *rays* (defined below) they belong to, and then furthermore, into *runs* within each ray. We then demonstrate that in order to determine the spectral multiplicity matrix for all representations of support S , it suffices to determine the multiplicity matrices for the individual runs, treated as the supports of a collection of representations.

Definition 1.1.1. Let C_n be the collection of (isomorphism classes of) supercuspidal representations of $GL_n(F)$, and define $\mathcal{C} := \bigcup_{n \in \mathbb{N}} C_n$.

For any choice of $\rho \in \mathcal{C}$, a *ray* is

$$\{\nu^a \rho \mid a \in \mathbb{Z}\},$$

where we consider the elements only up to isomorphism.

Definition 1.1.2. Let $S = \{\rho_1, \dots, \rho_r\}$ be a multi-set of supercuspidal representations belonging to the same ray. We define a *short run* to be a sub-multi-set R of S , such that the underlying set of R determines a segment (when appropriately ordered). We say the a short run R is simply a *run* if no other element of $S - R$ can be added to R while having in remain a run.

Suppose a_1, \dots, a_s are distinct runs (allowing for the possibility they even belong to distinct rays), and set $a = a_1 + a_2 + \dots + a_s$. By *Theorem 8.6*, [11], we know that

$$\langle a \rangle = \langle a_1 \rangle \times \langle a_2 \rangle \times \dots \times \langle a_s \rangle.$$

Indeed, we can also compute that

$$\pi(a) = \pi(a_1) \times \dots \times \pi(a_s).$$

Writing $a_i = \sum_{j=1}^{k_i} m(b_{ij}; a_i) \langle b_{ij} \rangle$, we find that

$$\pi(a) = \left(\sum_{j=1}^{k_1} m(b_{1j}; a_1) \langle b_{1j} \rangle \right) \times \dots \times \left(\sum_{j=1}^{k_s} m(b_{sj}; a_s) \langle b_{sj} \rangle \right) \quad (1.1)$$

$$= \sum_{(j_1, \dots, j_s)} m(b_{1j_1}; a_1) \cdot \dots \cdot m(b_{sj_s}; a_s) \langle b_{1j_1} \rangle \times \dots \times \langle b_{sj_s} \rangle. \quad (1.2)$$

The factors of each term

$$\langle b_{1j_1} \rangle \times \dots \times \langle b_{sj_s} \rangle,$$

belong to different runs, none of them can be linked. Hence by *Theorem 8.5*, [11] we know that

$$\langle b_{1j_1} \rangle \times \dots \times \langle b_{sj_s} \rangle = \langle b_{1j_1} + \dots + b_{sj_s} \rangle.$$

Namely, we know that each term is irreducible. Therefore, it suffices to separately determine the multiplicities $m(b_{ij}; a_i)$ for each $1 \leq i \leq s$ and $1 \leq j \leq k_i$. In other words, one need only consider the multi-segments belonging to a single run.

Multi-segments with multiplicity free support

Having demonstrated that it suffices to determine the multiplicity matrix run-by-run, we now turn to extreme of examples of the multiplicities of a given run. First, in this section, we turn to the case of *multiplicity free support*.

Let a be a multi-segment with multiplicity free support. In other words, for each $i \neq j$ we have $\Delta_i \cap \Delta_j = \emptyset$. Let

$$a_{\min} = \{\Delta_1, \dots, \Delta_r\}, \quad a_{\max} = \{[\rho], \dots, [\rho']\},$$

be the respectively minimal and maximal elements below and above a in the simple operations ordering on multi-segments.

Since the multi-segments have multiplicity free support, the segments (which are singletons) of a_{\max} are all disjoint.

Suppose $d = \{\Delta'_1, \dots, \Delta'_s\}$ is a multi-segment where each pair of segments have empty intersection. If Δ'_i and Δ'_j are linked, then consider the multi-segment b obtained by applying the elementary operation to Δ'_i and Δ'_j . By assumption $\Delta'_i \cap \Delta'_j = \emptyset$, and thus

$$\begin{aligned} b &= \{\Delta'_1, \dots, \Delta'_{i-1}, \Delta'_i \cup \Delta'_j, \Delta'_{i+1}, \dots, \Delta'_{j-1}, \Delta'_i \cap \Delta'_j, \Delta'_{j+1}, \dots, \Delta'_s\} \\ &= \{\Delta'_1, \dots, \Delta'_{i-1}, \Delta'_i \cup \Delta'_j, \Delta'_{i+1}, \dots, \Delta'_{j-1}, \emptyset, \Delta'_{j+1}, \dots, \Delta'_s\} \\ &= \{\Delta'_1, \dots, \Delta'_{i-1}, \Delta'_i \cup \Delta'_j, \Delta'_{i+1}, \dots, \Delta'_{j-1}, \Delta'_{j+1}, \dots, \Delta'_s\}. \end{aligned}$$

Suppose Δ''_k and Δ''_l are multi-segments of b with non-empty intersection. If Δ''_k and Δ''_l are both segments of a as well, this is a contradiction. Otherwise, Wlog $\Delta''_k = \Delta'_i \cup \Delta'_j$, and Δ''_l is

not. Hence, Δ_i'' is a segment of d , and thus $\Delta_i'' = \Delta_m'$ for some $m \neq i, j$. If it were the case that $\Delta_i'' = \Delta_m'$ had non-empty intersection with $\Delta_k'' = \Delta_i' \cup \Delta_j'$, then it must be the case that Δ_m' has non-empty intersection with either Δ_i' or Δ_j' , which would be contrary to our assumption on d . Thus, given a multi-segment with disjoint segments, and multi-segment obtained from it by elementary operations will also contain disjoint segments.

Returning to our segment a , by successively applying the above argument to a_{\max} , it must be that the segments of b are pair-wise disjoint for all multi-segment of the support of a . Therefore, by *proposition 9.13* of [11], we conclude that in $\mathfrak{R}(G)$

$$\pi(b) = \sum_{d \leq b} \langle d \rangle.$$

Thus, the geometric multiplicity matrix, indexed by segments obtained from a_{\max} , has the form

$$m(b; a) = \begin{cases} 1, & b \leq a \\ 0, & \text{else.} \end{cases} \quad (1.3)$$

Chapter 2

The geometry of $GL(n)$

2.1 The reduction of Vogan varieties of $GL(n)$

In this section, we describe several ways in which the study of Vogan varieties V_λ for infinitesimal parameters λ of $GL_n(F)$ over a p -adic field F may be reduced to simpler cases.

In the first section, we demonstrate that one need only determine V_λ and its corresponding multiplicity matrix m_{geo}^λ in the case where the cuspidal support of λ belongs to a single *run* (defined below). While this is not articulated explicitly, we believe that this is a manifestation of Bernstein decomposition on the "geometric side".

In the second section, we prove that given an infinitesimal parameter λ of $GL_{dN}(F)$ whose cuspidal support belongs to a single run, determined by a supercuspidal representation whose infinitesimal parameter has dimension d , there exist an (unramified) infinitesimal parameter λ^b of $GL_N(F)$ whose cuspidal support consists entirely of (unramified) characters, such that $V_\lambda \cong V_{\lambda^b}$ as varieties, and $\text{Per}_{H_\lambda}(V_\lambda) \cong \text{Per}_{H_{\lambda^b}}(V_{\lambda^b})$.

Recently, Clifton has dubbed the varieties V_{λ^b} "Athena" or "Athenian" varieties, as they simply "popped" out of Kristaps' head in much the same way Athena sprung forth from Zeus' head. It is expected that λ^b is simply λ_{ur} as it appears in [5], and that this procedure will allow one to easily generalize previous results of Connor Riddlesden on orbits and their duals, and hence generalize results of Mishty Ray on ABV-packets.

Geometric a-ray-ngments

Given a supercuspidal representation ρ of $GL_m(F)$, the *ray* or *line* determined by ρ is the set of (isomorphism classes of elements of)

$$\{v^k \rho : k \in \mathbb{Z}\}.$$

A *run* is a finite subset R of a ray such that if $v^l \rho, v^n \rho \in R$ with $l \leq n$, then (the isomorphism class of) $v^k \rho$ is in R for all k such that $l \leq k \leq n$.

Consider a finite collection of supercuspidals

$$\{\rho_{11}, \rho_{12}, \dots, \rho_{1,m_1}, \rho_{2,1}, \rho_{2,2}, \dots, \rho_{s,m_s}\},$$

in such a way that for any $(i, j), (k, l)$ we know that $i = k$ if and only if $\rho_{i,j}$ and $\rho_{k,l}$ belong to the same run. Writing the infinitesimal parameters as $\sigma_{ij} := \lambda_{\rho_{ij}}$, we have

$$\lambda = \bigoplus_{i,j} \sigma_{i,j}.$$

Note that if ρ and ρ' belong to the same run, then $\rho = v^k \rho'$ for some $k \in \mathbb{Z}$, hence for the associated infinitesimal parameters $\lambda_\rho = v^k \lambda_{\rho'}$. In particular, the dimension d_i of any infinitesimal parameter σ_{ij} will be the same for fixed i and valid choices of j . We will write each $X \in V_\lambda$ and $\lambda(w)$ in block-form where the $((i, j), (k, l))^{\text{th}}$ block is

$$(\dim \sigma_{ij}) \times \dim \sigma_{kl}.$$

The defining equation for $X \in V_\lambda$ is

$$\text{Ad}(\lambda(w))X = v(w)X, \quad \forall w \in W_F.$$

Since $\lambda(w)$ is diagonal, the condition on each block becomes that for all $w \in W_F$

$$\sigma_{ij}(w)X_{(ij),(kl)}\sigma_{kl}^{-1}(w) = v(w)X_{(ij),(kl)} \Rightarrow \sigma_{(ij)}(w)X_{(ij),(kl)} = X_{(ij),(kl)}v\sigma_{(kl)}(w).$$

Therefore, we may realize each block $X_{(ij),(kl)}$ as an intertwining operator for $\sigma_{(ij)}$ and $v\sigma_{(kl)}$. As it's not possible, by definition, for infinitesimal parameters arising from distinct runs to be isomorphic to a single twist of v of each other, this is only possible when $i = k$. Thus, we may re-write $X \in V_\lambda$ in terms of different blocks X_{ij} of size $(m_i \dim \lambda_{\rho_{i,-}}) \times (m_j \dim \lambda_{\rho_{j,-}})$. Then, by the above argument $X_{ij} = 0$ for $i \neq j$.

Applying an entirely similar argument, we can partition $g \in H_\lambda$ in the same manner as above to find that $g_{ij} = 0$ unless $i = j$. Therefore, the action of H_λ on V_λ is given by

$$\begin{bmatrix} g_{11} & 0 & \dots & 0 \\ 0 & g_{22} & \dots & 0 \\ \dots & & & \\ 0 & 0 & \dots & g_{ss} \end{bmatrix} \cdot \begin{bmatrix} X_{11} & 0 & \dots & 0 \\ 0 & X_{22} & \dots & 0 \\ \dots & & & \\ 0 & 0 & \dots & X_{ss} \end{bmatrix} = \begin{bmatrix} g_{11}X_{11}g_{11}^{-1} & 0 & \dots & 0 \\ 0 & g_{22}X_{22}g_{22}^{-1} & \dots & 0 \\ \dots & & & \\ 0 & 0 & \dots & g_{ss}X_{ss}g_{ss}^{-1} \end{bmatrix}.$$

Thus, writing

$$\lambda_i = \bigoplus_{j=1}^{m_i} \lambda_{\rho_{ij}},$$

for each $i \in \{1, \dots, s\}$, we can identify

$$V_\lambda \cong V_{\lambda_1} \times V_{\lambda_2} \times \dots \times V_{\lambda_s},$$

and

$$H_\lambda \cong H_{\lambda_1} \times H_{\lambda_2} \times \dots \times H_{\lambda_s},$$

where the action is given by

$$(g_1, \dots, g_s) \cdot (x_1, \dots, x_s) = (g_1 x_1 g_1^{-1}, \dots, g_s x_s g_s^{-1}).$$

Thus every orbit C of V_λ decomposes as

$$C = C_1 \times C_2 \times \dots \times C_s.$$

We also note that for all $X = (x_1, \dots, x_s) \in V_\lambda$,

$$\frac{\text{Stab}_{H_\lambda}(X)}{\text{Stab}_{H_\lambda}(X)^0} \cong \left(\frac{\text{Stab}_{H_{\lambda_1}}(x_1)}{\text{Stab}_{H_{\lambda_1}}(x_1)^0} \right) \times \dots \times \left(\frac{\text{Stab}_{H_{\lambda_s}}(x_s)}{\text{Stab}_{H_{\lambda_s}}(x_s)^0} \right).$$

Thus, for any orbit $C = C_1 \times \dots \times C_s$ and $X = (x_1, \dots, x_s) \in C$ we have

$$\pi_{H_\lambda}^{\text{ét}}(C, x) \cong \bigoplus_{i=1}^s \pi_{H_{\lambda_i}}^{\text{ét}}(C_i, x_i).$$

In a later section I will demonstrate that H_{λ_i} for a single run is itself isomorphic to a product of some copies of GL 's of varying size, and the stabilizer of any point will be isomorphic to a product of stabilizers within each GL . By the argument in *Lemma 3.6.3* [4], the stabilizer of any point under GL conjugation is connected. Therefore, the only irreducible H_{λ_i} -equivariant sheaves on a given orbit is the trivial one.

From now on, for each constituent V_{λ_i} of V_λ , we will write the s_i orbits as C_{ij} . Therefore, every orbit C of V_λ may be written as

$$C_{j_1, \dots, j_s} = C_{1j_1} \times \dots \times C_{sj_s}.$$

As a short-hand, we define $\mathbb{1}_{ij} := \mathbb{1}_{C_{ij}}$, and $\mathbb{1}_{j_1, \dots, j_s} := \mathbb{1}_{C_{j_1, \dots, j_s}}$. From [2], we know that there's a functor

$$\boxtimes : D_G^b(X, \mathbb{C}) \times D_H^b(Y, \mathbb{C}) \rightarrow D_{G \times H}^b(X \times Y, \mathbb{C}).$$

I conjecture that for every C_{j_1, \dots, j_s} of V_λ ,

$$\mathbb{1}_{j_1, \dots, j_s} \cong \mathbb{1}_{1j_1} \boxtimes \dots \boxtimes \mathbb{1}_{sj_s},$$

which should follow from *Lemma 3.3.14*, [2], or the results leading up to it. Speaking of *Lemma 3.3.14*, it does tell us that for any C_{j_1, \dots, j_s} of V_λ , and any finite-type local systems \mathcal{L}_{i,j_i} on each C_{i,j_i} ,

$$IC(C_{1j_1}, \mathcal{L}_{1j_1}) \boxtimes \dots \boxtimes IC(C_{sj_s}, \mathcal{L}_{sj_s}) \cong IC(C_{1j_1} \times \dots \times C_{sj_s}, \mathcal{L}_{1j_1} \boxtimes \dots \boxtimes \mathcal{L}_{sj_s}).$$

Suppose on faith, just for the time being, that

1. The only irreducible equivariant local systems on Vogan varieties of $GL(n)$ are the trivial ones (which I believe I know that proof, I only have to write it down above),
2. $\mathbb{1}_C \cong \mathbb{1}_{1j_1} \boxtimes \dots \boxtimes \mathbb{1}_{sj_s}$,
for every orbit $C = C_{1j_1} \times \dots \times C_{sj_s}$, (with appropriate shifts),
3. The distributivity of \boxtimes over \oplus for (perverse) sheaves (and/or local systems).

Then, writing the multiplicity of $\mathbb{1}_{i,j_i}$ in $IC(C_{i,k_i})$ as m_{j_i,k_i} , we find that

$$\begin{aligned}
 IC(\mathbb{1}_{j_1,\dots,j_s}) &= IC(C_{1,j_1} \times \dots \times C_{s,j_s}, \mathbb{1}_{1,j_1} \boxtimes \dots \boxtimes \mathbb{1}_{s,j_s}) \\
 &= IC(C_{1,j_1}, \mathbb{1}_{1,j_1}) \boxtimes \dots \boxtimes IC(C_{s,j_s}, \mathbb{1}_{s,j_s}) \\
 &= \left[\sum_{k_1} m_{k_1,j_1} \mathbb{1}_{1,k_1} \right] \boxtimes \dots \boxtimes \left[\sum_{k_s} m_{k_s,j_s} \mathbb{1}_{s,k_s} \right] \\
 &= \sum_{(k_1,\dots,k_s)} m_{k_1,j_1} \cdot \dots \cdot m_{k_s,j_s} \left(\mathbb{1}_{1,k_1} \boxtimes \dots \boxtimes \mathbb{1}_{s,k_s} \right) \\
 &= \sum_{(k_1,\dots,k_s)} m_{k_1,j_1} \cdot \dots \cdot m_{k_s,j_s} \cdot \mathbb{1}_{k_1,\dots,k_s},
 \end{aligned}$$

where the sum is in the Grothendieck group, and taken over all valid choices of (k_1, \dots, k_s) .

Therefore, we see that in order to compute the geometric multiplicity matrix of V_λ , it suffices to compute the multiplicity matrices of each of the constituent V_{λ_i} .

Note that the implication is that as categories?

$$\text{Per}_{H_\lambda}(V_\lambda) \cong \text{Per}_{H_{\lambda_1}}(V_{\lambda_1}) \otimes \dots \otimes \text{Per}_{H_{\lambda_s}}(V_{\lambda_s}),$$

which, on the spectral side predicts that there exists an equivalence of Grothendieck groups

$$\mathfrak{R}^\lambda(G) \cong \mathfrak{R}^{\lambda_i}(G) \otimes \dots \otimes \mathfrak{R}^{\lambda_s}(G).$$

The description V_λ in the case of a single run

The goal of this section is to prove that on the geometric side, we may reduce to the case of Vogan varieties whose infinitesimal parameters have cuspidal supports only consisting of characters (and by the previous section, those characters need only belong to a single run).

To be more specific, consider an infinitesimal parameter λ of $GL_N(F)$ corresponding to a single run, determined by a cuspidal representation ρ whose infinitesimal parameter λ_ρ is of dimension d , and for which the respective multiplicities of successive twists $v^i \rho$ are given by $m_0, \dots, m_i, \dots, m_s$. We will show that there is an isomorphism between $\text{Per}_{H_\lambda}(V_\lambda)$ and $\text{Per}_{H_{\lambda^b}}(V_{\lambda^b})$ where V_{λ^b} is a Vogan variety attached to $GL_N(F)$, attached to an infinitesimal parameter λ^b whose cuspidal support strictly contains the characters $1, v, \dots, v^s$ (potentially with multiplicity).

As we've seen, we can reduce to the case where for some infinitesimal parameter $\lambda : W_F \rightarrow GL_n(\mathbb{C})$ has support S whose elements belong to a single run. Thus, we can organize the support such that it has the form

$$S = \{\rho_{00}, \dots, \rho_{0,m_0}, \dots, \rho_{s,m_s}\},$$

where $\rho_{ij} = v^i \rho$ for some fixed supercuspidal representation ρ .

Writing σ_{ij} for the infinitesimal parameter of ρ_{ij} as in the previous question, and each $X \in V_\lambda$ and $\lambda(w)$ in $(\dim \sigma) \times (\dim \sigma)$ blocks, the defining equation for V_λ becomes that for all $w \in W_F$,

the $(i, j)^{\text{th}}$ block satisfies

$$\begin{aligned}\sigma_{ij}(w)X_{(ij),(kl)} &= X_{(ij),(kl)}\nu\sigma_{kl}(w) \\ \Rightarrow \nu^i\sigma(w)X_{(ij),(kl)} &= X_{(ij),(kl)}\nu\nu^k\sigma(w) \\ \Rightarrow \nu^i\sigma(w)X_{(ij),(kl)} &= X_{(ij),(kl)}\nu^{k+1}\sigma(w).\end{aligned}$$

Therefore we may realize $X_{(ij),(kl)}$ as an intertwining operator for $\nu^i\sigma$ and $\nu^{k+1}\sigma$. Hence, $X_{(ij),(kl)}$ will be zero unless $\nu^i\sigma \cong \nu^{k+1}\sigma$, in which case $i = k + 1$. Note that in this case, $X_{(ij),(kl)}$ be only be a scalar matrix of size $d := \dim \sigma$, and may be any such scalar matrix. Hence, each element $X \in V_\lambda$ may be written in block-form

$$X = \begin{bmatrix} 0 & 0 & 0 & \dots & 0 & 0 & 0 \\ X_{21} & 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & X_{32} & 0 & \dots & 0 & 0 & 0 \\ \dots & & & \dots & & & \dots \\ 0 & 0 & 0 & \dots & 0 & X_{s+1,s} & 0 \end{bmatrix},$$

where $X_{i+1,i}$ is a $(dm_i \times dm_{i-1})$ block, which itself may be written in terms of $(d \times d)$ blocks

$$X_{m_i, m_{i-1}} = \begin{bmatrix} Y_{11}^i & Y_{12}^i & \dots & Y_{1, m_{i-1}}^i \\ Y_{21}^i & Y_{22}^i & \dots & Y_{2, m_{i-1}}^i \\ \dots & \dots & \dots & \dots \\ Y_{m_i, 1}^i & \dots & \dots & Y_{m_i, m_{i-1}}^i \end{bmatrix},$$

where each Y_{jk}^i is a scalar matrix. Like-wise, each $g \in H_\lambda$ may be written as

$$G = \begin{bmatrix} G_1 & 0 & \dots & 0 \\ 0 & G_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & G_{s+1} \end{bmatrix} \in GL_{(m_0 + \dots + m_s)d}(\mathbb{C}),$$

where $G_i \in GL_{m_{i-1}}(\mathbb{C})$, which can be further written in $(d \times d)$ blocks

$$G_i = \begin{bmatrix} H_{11}^i & \dots & H_{1, m_{i-1}}^i \\ \dots & \dots & \dots \\ H_{m_{i-1}, 1}^i & \dots & H_{m_{i-1}, m_{i-1}}^i \end{bmatrix},$$

where H_{jk}^i is any scalar matrix such that $G_i \in G_{m_{i-1}}(\mathbb{C})$.

Athenian varieties

Given the support S of the previous sub-section, we form a new multi-set S^b which consists precisely of the characters ν^i occurring with multiplicity m_i . Let λ^b be the infinitesimal parameter

with support S^b , then by the results of the previous section, each $x \in V_{\lambda^b}$ is of the form

$$x = \begin{bmatrix} 0 & 0 & 0 & \dots & 0 & 0 & 0 \\ x_{21} & 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & x_{32} & 0 & \dots & 0 & 0 & 0 \\ \dots & & & & & & \\ 0 & 0 & 0 & \dots & 0 & x_{s+1,s} & 0 \end{bmatrix} \in V_{\lambda^b},$$

where $x_{i,i-1}$ is a block

$$X_{i,i-1} = \begin{bmatrix} y_{11}^i & y_{12}^i & \dots & y_{1,m_{i-1}}^i \\ y_{21}^i & y_{22}^i & \dots & y_{2,m_{i-1}}^i \\ \dots & & \dots & \dots \\ y_{m_{i-1},1}^i & \dots & \dots & y_{m_{i-1},m_{i-1}}^i \end{bmatrix}$$

size $(m_i \times m_{i-1})$ for $i \in \{1, 2, \dots, s\}$, and each $g \in H_{\lambda^b}$ is of the form

$$g = \begin{bmatrix} g_1 & 0 & \dots & 0 \\ 0 & g_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & g_{s-1} \end{bmatrix} \in GL_{(m_0+\dots+m_s)}(\mathbb{C}),$$

where $g_i \in GL_{n_0}(\mathbb{C})$, can be written as

$$g_i = \begin{bmatrix} h_{11}^i & \dots & h_{1,m_{i-1}}^i \\ \dots & \dots & \dots \\ h_{m_{i-1},1}^i & \dots & h_{m_{i-1},m_{i-1}}^i \end{bmatrix}$$

Define the isomorphism of varieties $\varphi : V_{\lambda^b} \rightarrow V_{\lambda}$ such that for any $x \in V_{\lambda^b}$ and its corresponding y_{jk}^i , the block Y_{jk}^i of the image $\varphi(x)$ is $y_{jk}^i I_{d,d}$. Like-wise, we define an isomorphism of algebraic groups $\psi : H_{\lambda^b} \rightarrow H_{\lambda}$ such that $g \in H_{\lambda} \subseteq GL_N(\mathbb{C})$ gets sent to the element $G \in H_{\lambda} \subseteq GL_{dN}(\mathbb{C})$ with sub-blocks $H_{jk}^i = h_{jk}^i I_{d,d}$.

The fact that these are indeed isomorphisms, and that φ is equivariant in the sense that for all $x \in V_{\lambda^b}$ and $g' \in H_{\lambda^b}$, we have

$$\varphi(\psi(g') \cdot x) = \psi(g') \cdot \varphi(x),$$

is intuitively "obvious", though I do plan to write out the relatively trivial but tedious details eventually.

Therefore $\text{Per}_{H_{\lambda^b}}(V_{\lambda^b}) \cong \text{Per}_{H_{\lambda}}(V_{\lambda})$, and the rest of the claims of the introduction to this section follow by the local Langlands correspondence.

2.2 Explicit descriptions on V_{λ} in extremal cases

The variety V_{λ} in the case of multiplicity free support

Now that we seen that the computation of the geometric multiplicity matrix may always be carried out run-by-run, we will turn to an examination of some of the extremal cases of the multiplicities

of a run. In this section, we will analyze the case of multiplicity-free support, and see that it agrees with the expectations of our computations on the spectral side, and the p -KLH.

Suppose $S = \{\rho_1, \dots, \rho_r\}$ is a finite collection of supercuspidal representations such that $\rho_i \not\cong \rho_j$ for $i \neq j$. Writing σ_i for the infinitesimal parameter of ρ_i , by the previous section we know that any representation with support S has the infinitesimal parameter

$$\lambda = \bigoplus_{i=1}^r \sigma_i.$$

The only block of $X \in V_\lambda$ which can be non-zero are precisely those X_{ij} for which $\sigma_i \cong \hat{\sigma}_j$. In this case X_{ij} can be any non-zero scalar matrix.

Like-wise, we see that for any $g \in H_\lambda$, the only blocks that can be non-zero are g_{ii} , each such block must be a scalar matrix (and such is valid), and since $g \in GL_n(\mathbb{C})$, each of the diagonal blocks must actually be non-zero.

Adopting the simplified notation $g_i := g_{ii}$, we find that the action of H_λ on V_λ is given by

$$\begin{aligned} g \cdot X &= \begin{bmatrix} g_1 & 0 & 0 & \dots & 0 & 0 \\ 0 & g_2 & 0 & \dots & 0 & 0 \\ \dots & & & & & \\ 0 & 0 & 0 & \dots & 0 & g_r \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 0 & \dots & 0 \\ X_{21} & 0 & 0 & \dots & 0 & 0 \\ \dots & & & & & \\ 0 & 0 & 0 & \dots & X_{r,r-1} & 0 \end{bmatrix} \begin{bmatrix} g_1^{-1} & 0 & 0 & \dots & 0 & 0 \\ 0 & g_2^{-1} & 0 & \dots & 0 & 0 \\ \dots & & & & & \\ 0 & 0 & 0 & \dots & 0 & g_r^{-1} \end{bmatrix} \\ &= \begin{pmatrix} 0 & 0 & 0 & 0 & \dots & 0 \\ g_2 X_{21} g_1^{-1} & 0 & 0 & \dots & 0 & 0 \\ \dots & & & & & \\ 0 & 0 & 0 & \dots & g_r X_{r,r-1} g_{r-1}^{-1} & 0 \end{pmatrix} \end{aligned}$$

Thus, we see that for any $g \in H_\lambda$ applied to some $X \in V_\lambda$, we see that the $(i, j)^{\text{th}}$ block of X_{ij} is non-zero iff the $(i, j)^{\text{th}}$ block of $g \cdot X$ is non-zero.

Given $X \in V_\lambda$, if $X_{21} = 0$ set $g_1 = I$, otherwise set $g_1 = X_{21}$. In either case set $g_2 = I$. Now supposing we've defined g_j for each $j \leq i$, for some $i \geq 2$. If $X_{i+1,i} = 0$ set $g_{i+1} = I$, otherwise set $g_{i+1} = g_i X_{21}^{-1}$.

Therefore, for the action of $g \cdot X$, either the block X_{ij} was zero and the block $[g \cdot X]_{ij}$ will remain zero, or X_{ij} is non-zero, where $j = i - 1$, and the block $[g \cdot X]_{i,i-1} = I$. Thus we see that H_λ is transitive among any $X \in V_\lambda$ with fixed choices of non-zero blocks. Therefore, for those i , say i_1, \dots, i_m such that $X_{i+1,i}$ can be nonzero, an orbit is simply a choice of which block will be non-zero. That is, there are 2^m orbits in total.

The above argument also tells us that for each orbit, we can choose the representative to be given by setting $X_{i+1,i} = I$ for those blocks we're choosing to be non-zero. Thus, the stabilizer of a any point will simply have $g_{i+1} = g_i$ for those i for which $X_{i+1,i}$ is non-zero. Therefore, suppose there are l blocks which are non-zero, then the stabilizer of a point in that orbit is isomorphic to $(\mathbb{C}^\times)^l$, which is connected. Therefore

$$\pi_{H_\lambda}^{\text{ét}}(C, X) = \frac{\text{Stab}_{H_\lambda}(X)}{\text{Stab}_{H_\lambda}(X)^0} \cong 0,$$

hence, on each orbit there is a single simple H_λ -equivariant local system.

Moreover, any orbit with l non-zero blocks will be isomorphic to $(\mathbb{C}^\times)^l$, whose closure in $V_\lambda \cong \mathbb{A}_{\mathbb{C}}^m$ is simply \mathbb{C}^l , which is smooth. Thus, recalling our basic results on perverse sheaves

1. $IC(\mathcal{L}_C)|_C = \mathcal{L}(C)[\dim C]$,
2. If \bar{C} is smooth, then $IC(\mathbb{1}_C)|_{C'} = \begin{cases} \mathbb{1}_{C'}[\dim C], & C' \subseteq \bar{C} \\ 0, & \text{otherwise} \end{cases}$
3. $IC(\mathcal{L}_C)|_{C'} = 0$ if $C' \not\subseteq \bar{C}$.

Taking the total dimension of the stalks of sheaves to obtain the entries of m_{geo}^λ , we can also label the matrix by non- \leq -decreasing orbits, in which case the p KLH is verified in comparison with our result 1.3.

The case of determinantal varieties

In this subsection, we consider the Vogan varieties which correspond to *Example 11.3* of Zelevinsky's [11]. In this case, the strata of V_λ are all *determinantal varieties*, which has come to be known as the "Tom Braden" case by the Voganish group.

Consider a support of the form

$$S = \left\{ \underbrace{\rho, \dots, \rho}_{k \text{ times}}, \underbrace{\nu\rho, \dots, \nu\rho}_{l \text{ times}} \right\}.$$

Let σ_i be the infinitesimal parameter λ_ρ of ρ for $1 \leq i \leq k$, and $\sigma_j := \nu\lambda_\rho$ for $k+1 \leq j \leq k+l$, and set $r = \dim \lambda_\rho$. Any representation with support S has infinitesimal parameter

$$\lambda = \bigoplus_{i=1}^{k+l} \sigma_i.$$

Writing $X \in V_\lambda$ in r by r blocks, from ?? we know that X_{ij} is non-zero iff $\sigma_i \cong \nu\sigma_j$. This is possible if, and only if $1 \leq i \leq k$ and $k+1 \leq j \leq k+l$. For each such choice of i and j , it must be that that X_{ij} is a scalar matrix.

An entirely similar argument shows us that $(i, j)^{\text{th}}$ block of $g \in H_\lambda$ can be non-zero iff $\sigma_i \cong \sigma_j$. That is, the only (potential) non-zero blocks are g_{ij} for $1 \leq i, j \leq k$ and $k+1 \leq i, j \leq k+l$. In any such case, g_{ij} is an intertwining operator, and thus a scalar matrix. Therefore, the element of H_λ can be written in blocks

$$g = \begin{bmatrix} A_{kr,kr} & 0_{kr,lr} \\ 0_{lr,kr} & B_{lr,lr} \end{bmatrix} = \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix},$$

where A and B are composed of r -by- r blocks, subject to the condition that $g \in GL_{(k+l)r}(\mathbb{C})$. Writing $X \in V_\lambda$ as

$$X = \begin{bmatrix} 0_{k,k} & 0_{k,l} \\ Z_{l,k} & 0_{l,l} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ Z & 0 \end{bmatrix}$$

the action of H_λ on V_λ is given by

$$\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \begin{bmatrix} 0 & 0 \\ Z & 0 \end{bmatrix} \begin{bmatrix} A^{-1} & 0 \\ 0 & B^{-1} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ BZA^{-1} & 0 \end{bmatrix}.$$

claim: The proof of the Tom Braden case which gives us the rank-stratification goes through entirely similarly to tell us that, V_λ is stratified into orbits

$$C_0 < C_1 < C_2 < \dots < C_k,$$

where C_i is the orbit of those block-scalar matrices of rank ir , and by adapting this argument, $\dim C_i = i(k + l - i)$. The stabilizer of any point $Z_i \in C_i$ will be isomorphic to

The closure \bar{C}_i will be smooth for $i = 0, k$, and for each $i \neq 0, k$ it will be the union of all smaller orbits, which will be singular.

We may identify each

$$X = \begin{bmatrix} 0 & 0 \\ Z & 0 \end{bmatrix} \in V_\lambda,$$

with the matrix $\tilde{Z} \in M_{l,k}(\mathbb{C})$ whose $(i, j)^{\text{th}}$ entry is the scalar of the $(i, j)^{\text{th}}$ r -by- r block of Z . Like-wise, we may identify H_λ with $GL_l \times GL_k$.

Take as a representative of the orbit C_i , the matrix Z_i which consists of the i -by- i identity matrix in the upper left block, and 0's everywhere else. While I have not yet done so here in full generality and formality, it appears that the stabilizer will always be connected. However, an explicit description of this is somewhat complicated and tedious. I am hoping that I might find a more elegant explanation than direct abstract computation.

Once the connectedness of stabilizers is established, we can conclude that the only irreducible equivariant local systems are the trivial ones. Thus, as a short-hand we define $\mathbb{1}_i := \mathbb{1}_{C_i}$.

Claim: According to this Wikipedia entry for *Determinantal varieties*,

$$\tilde{C}_i = \{(Z, V) \in M_{l,k}(\mathbb{C}) \times G(i, l) \mid Z(\mathbb{C}^k) \subseteq V\},$$

is a resolution of singularities for \bar{C}_i , via the projection map π_i on the first factor, where $G(i, l)$ is the complex Grassmannian of i -planes in l -space. However, by the symmetry of the situation, in order to ensure that our maps are semi-small, we will instead take

$$\tilde{C}_i = \{(Z^t, V) \in M_{k,l}(\mathbb{C}) \times G(i, k) \mid Z(\mathbb{C}^l) \subseteq V\},$$

where π_i is now the transpose of projection onto the first factor.

For $j \leq i$, let $Z_j \in C_j$, the fiber $\pi_i^{-1}\{Z_j\}$ must consist of all pairs (Z_j, V) where $V \in G(i, k)$, and $V \supseteq Z_j(\mathbb{C}^l)$. To re-phrase this, we could view it as making the choice of quotient $V/Z_j(\mathbb{C}^l)$ within $\mathbb{C}^k/Z_j(\mathbb{C}^l)$. Thus, the fiber $\pi_i^{-1}\{Z_j\}$ can be realized any any choice of a subspace of dimension

$$\dim(V/Z_j(\mathbb{C}^l)) = \dim V - \dim Z_j(\mathbb{C}^l) = i - j,$$

(recalling that Z has rank j), from a total space of dimension

$$\dim(\mathbb{C}^k/Z_j(\mathbb{C}^l)) = \dim \mathbb{C}^k - \dim Z_j(\mathbb{C}^l) = k - j.$$

In other words, we can make the identification $\pi_i^{-1}\{Z_j\} \cong G(i-j, k-j)$.

In order to prove that these covers are in fact semi-small we need to confirm that for all $C_j \subseteq \bar{C}_i$, that is, all C_j for $j \leq i$, and $Z_j \in C_j$,

$$\begin{aligned} \dim C_j + 2 \dim \pi_i^{-1}\{Z_j\} &\leq \dim \bar{C}_i \\ j(k+l-j) + 2 \dim G(i-j, k-j) &\leq i(k+l-i) \\ j(k+l-j) + 2(i-j)(k-i) &\leq i(k+l-i). \end{aligned}$$

For $j = 0$ this reduces to $k-i \leq l$ which is certainly true. Proceeding by induction... For some reason I'm having the damndest time trying to prove this, but having tested this inequality for many values of $0 \leq j \leq i \leq k \leq l$ in Python, I'm confident that this map is indeed semi-small, and more-over that the only relevant orbit of π_i is C_i itself. Thus, pending a more formal verification, for any choice of (i, j) such that $0 \leq j \leq i \leq k \leq l$, and $Z \in C_i$,

$$\pi_{i*} \mathbb{1}_i[i(k+l-i)] = mIC(\mathbb{1}_i).$$

Given $Z \in C_j$, it's a general fact about IC 's that

$$IC(\mathbb{1}_i)_Z = IC(\mathbb{1}_i)|_{C_i} = \mathbb{1}_i[i(k+l-i)].$$

Meanwhile, the fiber $\pi_i^{-1}\{Z\} = G(k-i, k-i) \cong \{pt\}$, and as

$$H^i(\{pt\}) = \begin{cases} \mathbb{1}_{\bar{C}_i}, & i = 0 \\ 0, & \text{else} \end{cases},$$

we conclude that

$$\begin{aligned} mIC(\mathbb{1}_i)_Z &= \left(\pi_{i*} \mathbb{1}_{\bar{C}_i}[i(k+l-i)] \right)_{Z_j} \\ IC(\mathbb{1}_i)|_{C_i} &= H^\bullet(\{pt\})[i(k+l-i)] \\ \mathbb{1}_i[i(k+l-i)] &= \mathbb{1}_i[i(k+l-i)], \end{aligned}$$

meaning $m = 1$.

More generally, according to *Lie Groups, Bump, pages 522*, the rank of the cohomology ring of $G(i-j, k-j) = G(k-i, k-j)$ is $\binom{k-j}{k-i}$, which is the "total" dimension of $IC(C_i, \mathbb{1}_i)|_{C_j}$. For the purposes of the p -KLH, we're interested in the total dimension $\dim IC(\mathbb{1}_i)|_{C_j}$, and thus, for any (i, j) such that $0 \leq j \leq i \leq k \leq l$ and $Z_j \in C_i$, we see that

$$\begin{aligned} \dim IC(\mathbb{1}_i)_{Z_j} &= \dim \left(\pi_{i*} \mathbb{1}_{\bar{C}_i}[i(k+l-i)] \right)_{Z_j} \\ \dim IC(\mathbb{1}_i)|_{C_j} &= \dim (H^\bullet(\pi_i^{-1}\{Z_j\})) \\ &= \dim H^\bullet(G(i-j, k-j)) \\ &= \binom{k-j}{k-i}. \end{aligned}$$

This completes the computation of the geometric multiplicity matrix in this case, that is,

$$[m_{\text{geo}}^\lambda]_{ij} = \dim IC(\mathbb{1}_i)|_{C_j} = \binom{k-j}{k-i},$$

where we define $\binom{k-j}{k-i} = 0$ for $k-i > k-j$. According to *Example 11.3* of [11], the spectral multiplicity matrix is given by

$$[m_{\text{rep}}^\lambda]_{ij} = \binom{k-i}{k-j} = [m_{\text{geo}}^\lambda]_{ji},$$

and thus ${}^t m_{\text{rep}}^\lambda = m_{\text{geo}}^\lambda$, verifying the p -KLH in this case.

Chapter 3

The book of $GL(4)$

In this chapter, we give explicit descriptions of the multiplicity tables for $GL(4)$ on both the spectral and geometric sides.

By Bernstein decomposition on the spectral side, or correspondingly the Vogan decomposition on the geometric side, we know that we need only consider the cases of cuspidal support belonging to a single ray. More-over, barring some proper references, we need only consider the multiplicities of "twists" by the determinant character ν that arise, and, with respect to the aims of the p -adic Kazhdan-Lusztig hypothesis, we also do not care which representations have particular multiplicities, only the multiplicities that appear.

We begin with the principal series representations, which is to say, twists of the identity representation 1 , by ν . We will describe the cuspidal support by the powers of ν that appear. For example, we will describe $\{1, \nu, \nu, \nu^2\}$ by the integer sequence $\{0, 1, 1, 2\}$.

First, consider the case where we only have a single representation, for which of course, there is a single possibility $\{0, 0, 0, 0\}$.

If we have two representations, then either one appears with multiplicity one, and the other with multiplicity three, or both occur with multiplicity two. That is $\{0, 0, 1\}$ or $\{0, 0, 1, 1\}$.

For the case of three representations, the only possibility is that one representation occurs with multiplicity two, and the others occur with multiplicity one, hence $\{0, 0, 1, 2\}$.

Finally, there is a unique support where each representation occurs with multiplicity one: $\{0, 1, 2, 3\}$.

In summary, the possible supports corresponding to principal series representations, up to multiplicities of representations, and arrangements of their positions, are

$$\{0, 0, 0, 0\}, \{0, 0, 0, 1\}, \{0, 0, 1, 1\}, \{0, 0, 1, 2\}, \{0, 1, 2, 3\}.$$

Aside from considering a supercuspidal representation of $GL(4)$ itself

3.1 The determinantal case

The spectral side

Consider the following family of multi-segments with support $\{\varphi, \varphi, \nu\varphi, \nu\varphi\}$,

$$\begin{aligned} a_0 &= \{[\varphi], [\varphi], [\nu\varphi], [\nu\varphi]\} \\ a_1 &= \{[\varphi, \nu\varphi], [\varphi], [\nu\varphi]\} \\ a_2 &= \{[\varphi, \nu\varphi], [\varphi, \nu\varphi]\}. \end{aligned}$$

By *Example 11.3* of [11], we know that $a_0 > a_1 > a_2$, and in the Grothendieck group $\mathfrak{R}(G)$,

$$\pi(a_i) = \sum_{i \leq j \leq k} C_{k-j}^{k-i} \langle a_j \rangle.$$

Therefore

$$\begin{aligned} \pi(a_0) &= \sum_{0 \leq j \leq 2} C_{2-j}^2 \langle a_j \rangle \\ &= C_2^2 \langle a_0 \rangle + C_1^2 \langle a_1 \rangle + C_0^2 \langle a_2 \rangle \\ &= \langle a_0 \rangle + 2 \langle a_1 \rangle + \langle a_2 \rangle. \\ \pi(a_1) &= \sum_{1 \leq j \leq 2} C_{2-j}^1 \langle a_j \rangle \\ &= C_1^1 \langle a_1 \rangle + C_0^1 \langle a_2 \rangle \\ &= \langle a_1 \rangle + \langle a_2 \rangle. \\ \pi(a_2) &= \sum_{2 \leq j \leq 2} C_{2-j}^0 \langle a_j \rangle \\ &= C_0^0 \langle a_2 \rangle \\ &= \langle a_2 \rangle. \end{aligned}$$

Therefore, the spectral multiplicity matrix can be seen to be

m_{rep}	$\langle a_0 \rangle$	$\langle a_1 \rangle$	$\langle a_2 \rangle$
$\pi(a_0)$	1	2	1
$\pi(a_1)$	0	1	1
$\pi(a_2)$	0	0	1

Determination of V_λ , H_λ , and the orbits

Note that this theory was completely worked out in greater generality in previous sections, but we take an overly explicit and naive approach here just to liven things up a little, and gain some broader perspective.

Let $\psi : W_F \rightarrow GL_4(\mathbb{C})$ be the infinitesimal parameter of φ . Therefore, the infinitesimal parameter of the irreducible and standard representations appearing in the previous section is

$$\lambda(w) = \begin{bmatrix} \psi(w) & 0 & 0 & 0 \\ 0 & \psi(w) & 0 & 0 \\ 0 & 0 & v\psi(w) & 0 \\ 0 & 0 & 0 & v\psi(w) \end{bmatrix}.$$

Writing $U \in V_\lambda$ in 2 by 2 blocks,

$$U = \begin{bmatrix} X & Y \\ Z & W \end{bmatrix},$$

we can expand the defining condition on V_λ as being, for all $w \in W_F$,

$$\begin{aligned} \lambda(w)X\lambda(w)^{-1} &= v(w)X \\ \begin{bmatrix} \psi(w)I_2 & 0 \\ 0 & v\psi(w)I_2 \end{bmatrix} \begin{bmatrix} X & Y \\ Z & W \end{bmatrix} \begin{bmatrix} \psi(w)^{-1}I_2 & 0 \\ 0 & v\psi(w)^{-1}I_2 \end{bmatrix} &= \begin{bmatrix} v(w)X & v(w)Y \\ v(w)Z & v(w)W \end{bmatrix} \\ \begin{bmatrix} X & v^{-1}(w)Y \\ v(w)Z & W \end{bmatrix} &= \begin{bmatrix} v(w)X & v(w)Y \\ v(w)Z & v(w)W \end{bmatrix}. \end{aligned}$$

As this must hold for all $w \in W_F$, this is only possible if $X = Y = W = 0$. Meanwhile, any choice for $Z \in M_{22}(\mathbb{C})$ is valid, hence

$$V_\lambda = \left\{ \begin{bmatrix} 0 & 0 \\ Z & 0 \end{bmatrix} : Z \in M_{22}(\mathbb{C}) \right\}.$$

By an entirely similar argument

$$H_\lambda = \left\{ \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \in GL_4(\mathbb{C}) \right\}.$$

Now

$$\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \cdot \begin{bmatrix} 0 & 0 \\ X & 0 \end{bmatrix} = \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \begin{bmatrix} 0 & 0 \\ Z & 0 \end{bmatrix} \begin{bmatrix} A^{-1} & 0 \\ 0 & B^{-1} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ BX A^{-1} & 0 \end{bmatrix}.$$

Thus, it follows that the orbits of V_λ are determined entirely by the rank of Z . We index them by their corresponding rank C_0, C_1, C_2 , where

$$\dim C_0 = 0, \quad \dim C_1 = 3, \quad \dim C_2 = 4.$$

Determination of the stabilizers/equivariant fundamental groups

First we note that the stabilizer of the zero matrix, being the only element of C_0 is all of H_λ , and since

$$H_\lambda \cong GL_2(\mathbb{C}) \times GL_2(\mathbb{C}),$$

is connected, hence $\pi_{H_\lambda}^{\text{ét}}(C_0, 0) = 0$.

For

$$\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \in H_\lambda,$$

write

$$B = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \quad A^{-1} = \begin{bmatrix} \alpha & \beta \\ \gamma & \delta \end{bmatrix},$$

we attempt to compute the stabilizer of a point in the rank one orbit C_1 , by demanding

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \alpha & \beta \\ \gamma & \delta \end{bmatrix} = \begin{bmatrix} a\alpha & a\beta \\ c\alpha & c\beta \end{bmatrix}.$$

Equating both sides, we determine that $\alpha = a^{-1}$, hence neither are zero, and this forces $\beta = c = 0$. Therefore, by taking the inverse and making appropriate substitutions, we find that

$$A = \frac{1}{\alpha\delta - \beta\gamma} \begin{bmatrix} \delta & -\beta \\ -\gamma & \delta \end{bmatrix} = \frac{a}{\delta} \begin{bmatrix} \delta & 0 \\ -\gamma & a^{-1} \end{bmatrix}.$$

Therefore, the stabilizer is given by

$$\begin{bmatrix} a & 0 & 0 & 0 \\ -\gamma/\delta & 1/\delta & 0 & 0 \\ 0 & 0 & a & b \\ 0 & 0 & 0 & d \end{bmatrix},$$

with the only restrictions being $a, \delta, d \neq 0$. Hence the variety is isomorphic to

$$\mathbb{C}^\times \times \mathbb{C}^\times \times \mathbb{C}^\times \times \mathbb{C} \times \mathbb{C},$$

which is connected. That is, for any $Z \in C_1$ we find $\pi_{H_\lambda}^{\text{ét}}(C_1, X) = 0$.

Choosing the identity matrix I for the representative of C_2 , we find that the stabilizer must satisfy $BA^{-1} = I \Rightarrow A = B$. Therefore the stabilizer is just $GL_2(\mathbb{C})$ as a variety, which is connected, and thus $\pi_{H_\lambda}^{\text{ét}}(C_2, I) = 0$.

Thus we have seen that each of the equivariant fundamental groups will be trivial, and therefore the only sheaves to account for are those that arise from trivial the local system on each orbit.

Note that $\bar{C}_0 = 0$ and $\bar{C}_2 \cong V_\lambda \cong \mathbb{A}_{\mathbb{C}}^4$ are both smooth. Thus, by the result that

1. $IC(\mathcal{L}_C)|_C = \mathcal{L}(C)[\dim C]$,
2. If \bar{C} is smooth, then $IC(\mathbb{1}_C)|_{C'} = \begin{cases} \mathbb{1}_{C'}[\dim C], & C' \subseteq \bar{C} \\ 0, & \text{otherwise} \end{cases}$
3. $IC(\mathcal{L}_C)|_{C'} = 0$ if $C' \not\subseteq \bar{C}$,

we can begin to fill out the geometric multiplicity matrix as

m_{geo}	$P _{C_0}$	$P _{C_1}$	$P _{C_2}$
$IC(\mathbb{1}_{C_0})$	$\mathbb{1}_{C_0}[0]$	0	0
$IC(\mathbb{1}_{C_1})$?	$\mathbb{1}_{C_1}[3]$	0
$IC(\mathbb{1}_{C_2})$	$\mathbb{1}_{C_0}[4]$	$\mathbb{1}_{C_1}[4]$	$\mathbb{1}_{C_2}[4]$

As we'll see below, \bar{C}_1 is singular, and thus we'll need to find a resolution. We know from the KS paper, for example, that the closure of the orbit of rank 1 matrices is the collection of matrices with rank *at most* 1. Thus, since we're identifying the total space with $M_{22}(\mathbb{C})$, we can make the realization

$$\bar{C}_1 \cong \{(a, b, c, d) \in \mathbb{A}_{\mathbb{C}}^4 \mid ad - bc = 0\}.$$

Differentiating the equation at the four variables, we find that we have precisely one singular point $(a, b, c, d) = (0, 0, 0, 0)$. Taking another perspective $\bar{C}_1 = \{X \in GL(2) \mid \det X = 0\}$, and

$$\tilde{C}_1 := \{(Z, v) \in \bar{C}_1 \times \mathbb{P}^1 \mid Zv = 0\},$$

gives us a smooth resolution by projection on the first factor $f : \tilde{C}_1 \rightarrow \bar{C}_1$. I will not prove this here, but am taking it on faith that this is true given other calculations that I have seen. So, it is possible this a place where I am going wrong.

Recall that in order for f to be semi-small, for every orbit $C \subseteq \bar{C}_1$, and every $Z \in C$ we have

$$\dim C + 2 \dim f^{-1}\{Z\} \leq \dim \tilde{C}_1.$$

Those orbits for which we have equality are said to be *relevant*. The orbits contained in \bar{C}_1 are exactly C_0 and C_1 .

Since, for the zero-matrix $0v = 0$ for all $v \in \mathbb{P}^1$, we see that the fiber

$$f^{-1}\{0\} = \{(0, v) \in \bar{C}_1 \times \mathbb{P}^1 \mid Xv = 0\} \cong \mathbb{P}^1.$$

Therefore,

$$\dim C_0 + 2 \dim f^{-1}\{0\} = 0 + 2(1) = 2 < 3 \dim \tilde{C}_1.$$

Hence, C_0 does fulfill the requirement upon it for f to be semi-small, but it is *not* relevant.

As a representative of $Z \in C_1$, consider

$$Z_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}.$$

The fiber is given by

$$f^{-1}\{Z_0\} = \{(Z_0, v) \in \bar{C}_1 \times \mathbb{P}^1 \mid Z_0 v = 0\}.$$

Writing $v \in \mathbb{P}^1$ as

$$v = \begin{bmatrix} a \\ b \end{bmatrix},$$

we can compute

$$\begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} b \\ 0 \end{bmatrix} \sim \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

Assuming this behavior is generic, the fiber above any point of C_1 is a single point. Thus, for any $Z \in C_1$ we compute

$$\dim C_1 + 2 \dim f^{-1}\{Z\} = 3 + 2(0) = 3 \leq 3 = \dim \tilde{C}_1.$$

Therefore, any point $Z \in C_1$ will satisfy the semi-small condition, hence we conclude that f itself is semi-small, and C_1 is the only relevant orbit.

By the *Decomposition Theorem*

$$\begin{aligned} f_* \mathbb{L}_{\tilde{C}_1}[\dim \tilde{C}_1] &= mIC(\mathbb{L}_{C_1}) \\ f_* \mathbb{L}_{\tilde{C}_1}[3] &= mIC(\mathbb{L}_{C_1}). \end{aligned}$$

Let $Z \in C_1$. We know in general that for $Z_1 \in C_1$

$$IC(\mathbb{L}_{C_1})_{Z_1} \cong IC(\mathbb{L}_{C_1})|_{C_1} = \mathbb{L}_{C_1}[3].$$

More-over,

$$(f_* \mathbb{L}_{\tilde{C}_1}[3])_Z = H^\bullet(f^{-1}\{Z\})[3] = H^\bullet(pt)[3].$$

Since

$$H^i(pt) = \begin{cases} \mathbb{L}_{\tilde{C}_1}, & i = 0 \\ 0, & \text{else} \end{cases}$$

we see that

$$\begin{aligned} (f_* \mathbb{L}_{\tilde{C}_1}[3])_Z &= mIC(\mathbb{L}_{C_1})_Z \\ \mathbb{L}_{C_1}[3] &= m\mathbb{L}_{C_1}[3]. \end{aligned}$$

Therefore $m = 1$.

Taking the stalk at the zero matrix,

$$IC(\mathbb{L}_{C_1})|_{C_0} \cong IC(\mathbb{L}_{C_1})_0 = (f_* \mathbb{L}_{\tilde{C}_1}[3])_0 = H^\bullet(f^{-1}\{0\})[3],$$

where

$$H^i(f^{-1}\{0\}) = H^i(\mathbb{P}^1) = \begin{cases} \mathbb{L}_{\tilde{C}_1}, & i = 0, 2 \\ 0, & \text{otherwise} \end{cases}$$

For some reason I don't understand, when we take cohomology, we shift to the negative degree, therefore the above reduces to

$$\begin{aligned} IC(\mathbb{L}_{C_1})_0 &= (f_* \mathbb{L}_{\tilde{C}_1}[3])_0 \\ &= (\mathbb{L}_{\tilde{C}_1}[0] \oplus \mathbb{L}_{\tilde{C}_1}[-2])_0[3] \\ &= \mathbb{L}_{C_0}[3] \oplus \mathbb{L}_{C_0}[1]. \end{aligned}$$

Hence, we may complete the table as

m_{geo}^λ	$P _{C_0}$	$P _{C_1}$	$P _{C_2}$
$IC(\mathbb{1}_{C_0})$	$\mathbb{1}_{C_0}[0]$	0	0
$IC(\mathbb{1}_{C_1})$	$\mathbb{1}_{C_0}[1] \oplus \mathbb{1}_{C_0}[3]$	$\mathbb{1}_{C_1}[3]$	0
$IC(\mathbb{1}_{C_2})$	$\mathbb{1}_{C_0}[4]$	$\mathbb{1}_{C_1}[4]$	$\mathbb{1}_{C_2}[4]$

Thus, counting the (total) dimensions of the stalks, we compare the multiplicity matrices side-by-side

m_{geo}^λ	$P _{C_0}$	$P _{C_1}$	$P _{C_2}$	m_{rep}	$\langle a_0 \rangle$	$\langle a_1 \rangle$	$\langle a_2 \rangle$
$IC(\mathbb{1}_{C_0})$	1	0	0	$\pi(a_0)$	1	2	1
$IC(\mathbb{1}_{C_1})$	2	1	0	$\pi(a_1)$	0	1	1
$IC(\mathbb{1}_{C_2})$	1	1	1	$\pi(a_2)$	0	0	1

which is consistent with what we expect by our computations on the spectral side and the Kazhdan-Lusztig hypothesis.

3.2 An "unresolved" case: Zelevinsky's 11.4

The spectral side


From [11], we consider *Example 11.4*. Specializing to the case of $\rho = \mathbb{1}$, the trivial representation, we may write the support as

$$S = \{0, 1, 1, 2\},$$

with multi-segments,

$$a_{\max} = \{0, 0, 1, 2\}, a_l = \{[0, 1], 1, 2\}, a_r = \{0, 1, [1, 2]\}, a_0 = \{[0, 1], [1, 2]\}, a_{\min} = \{[0, 2], [1]\}.$$

The corresponding lattice structure is given by



Then, as demonstrated in [11], the spectral multiplicity matrix is given by

m_{geo}	$\langle a_{\max} \rangle$	$\langle a_l \rangle$	$\langle a_r \rangle$	$\langle a_0 \rangle$	$\pi(a_{\min})$
$\pi(a_{\max})$	1	1	1	2	1
$\pi(a_l)$	0	1	0	1	1
$\pi(a_r)$	0	0	1	1	1
$\pi(a_0)$	0	0	0	1	1
$\pi(a_{\min})$	0	0	0	0	1

The Geometry

The Vogan variety corresponding to the support $S = \{1, v, v, v^2\}$ can be seen to be of the form

$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ * & 0 & 0 & 0 \\ * & 0 & 0 & 0 \\ 0 & * & * & 0 \end{bmatrix}$$


Letting X be the column matrix on the left, and Y the row matrix on the bottom, we write the orbits as

$$C_{ijk} = \{(X, Y) \in M_{2,1}(\mathbb{C}) \times M_{1,2}(\mathbb{C}) \mid \text{rk}(X) = i, \text{rk}(Y) = j, \text{rk}(YX) = k\}.$$

The full list of orbits is

$$C_{000}, C_{010}, C_{100}, C_{110}, C_{111}.$$

It would appear from the KS paper that the closure of any of these orbits, simply has the effect of replacing the "=" sign in their definitions with " \leq ". Thus the corresponding lattice structure is given by



which is exactly what we expect from the spectral side. From our expectations of the Kazhdan-Lusztig hypothesis, I then suppose it will be the case that

$$\dim IC(C_{110}, \mathbb{1}_{110})|_{C_{000}} = 2.$$

Given the expectations that all irreducible local systems are trivial, we might write this more simply as

$$\dim IC_{110}|_{000} = 2.$$

The dimensions of the orbits are given by

$$\dim C_{111} = 4??? \quad \dim C_{110} = 3, \quad \dim C_{100} = 2, \quad \dim C_{010} = 2, \quad \dim C_{000} = 0.$$

Side note: I still wonder, based off the work of [4], if we can base change the moment map $\mu : \tilde{\mathfrak{g}} \rightarrow \mathfrak{g}$ to \tilde{C}_{110} to get

$$\tilde{C}_{110} = \{(x, p) \in \tilde{C}_{110} \times G/P_\lambda \mid x \in p\}.$$

Since arbitrary flag varieties can be singular, we may then have to apply the Bott-Smaleson resolution to G/P_λ , and compose with the moment map μ to get a resolution of singularities that we can use. Note that by the claims of *Elijah*, the parabolic P_λ we see, is "the same" parabolic arising in Zelevinsky's short paper in which he embeds the closures of orbits into Schubert varieties.

Again, without even resolving singularities, we may begin to fill out the multiplicity table. Recall that we have our 3 basic tools,

1. $IC(\mathcal{L}_C)|_C = \mathcal{L}(C)[\dim C]$,
2. If \bar{C} is smooth, then $IC(\mathbb{1}_C)|_{C'} = \begin{cases} \mathbb{1}_{C'}[\dim C], & C' \subseteq \bar{C} \\ 0, & \text{else} \end{cases}$
3. $IC(\mathcal{L}_C)|_{C'}$ if $C' \subsetneq \bar{C}$.

Therefore, properties 1. and 3. immediately give us the diagonal and upper diagonal respectively. That is, writing $\mathbb{1}_{ijk} := \mathbb{1}_{C_{ijk}}$, and $IC_{ijk} := IC(C_{ijk}, \mathbb{1}_{ijk})$.

m_{geo}^λ	$ _{C_{000}}$	$ _{C_{010}}$	$ _{C_{100}}$	$ _{C_{110}}$	$ _{C_{111}}$
IC_{000}	$\mathbb{1}_{000}[0]$	0	0	0	0
IC_{010}		$\mathbb{1}_{010}[2]$	0	0	0
IC_{100}			$\mathbb{1}_{100}[2]$	0	0
IC_{110}				$\mathbb{1}_{110}[3]$	0
IC_{111}					$\mathbb{1}_{111}[4]$

Moreover, since $\bar{C}_{111} \cong \mathbb{A}_{\mathbb{C}}^4$ is smooth, property two of the above allows us to fill in the bottom row as

m_{geo}^λ	$ _{C_{000}}$	$ _{C_{010}}$	$ _{C_{100}}$	$ _{C_{110}}$	$ _{C_{111}}$
IC_{000}	$\mathbb{1}_{000}[0]$	0	0	0	0
IC_{010}		$\mathbb{1}_{010}[2]$	0	0	0
IC_{100}			$\mathbb{1}_{100}[2]$	0	0
IC_{110}				$\mathbb{1}_{110}[3]$	0
IC_{111}	$\mathbb{1}_{111}[0]$	$\mathbb{1}_{111}[2]$	$\mathbb{1}_{111}[2]$	$\mathbb{1}_{111}[3]$	$\mathbb{1}_{111}[4]$

Likewise, $\bar{C}_{010} \cong \mathbb{A}_{\mathbb{C}}^2 \cong \bar{C}_{100}$, and thus again by property 2. we can complete the entries of the table as

m_{geo}^λ	$ _{C_{000}}$	$ _{C_{010}}$	$ _{C_{100}}$	$ _{C_{110}}$	$ _{C_{111}}$
IC_{000}	$\mathbb{1}_{000}[0]$	0	0	0	0
IC_{010}	$\mathbb{1}_{000}[2]$	$\mathbb{1}_{010}[2]$	0	0	0
IC_{100}	$\mathbb{1}_{000}[2]$	0	$\mathbb{1}_{100}[2]$	0	0
IC_{110}				$\mathbb{1}_{110}[3]$	0
IC_{111}	$\mathbb{1}_{111}[0]$	$\mathbb{1}_{111}[2]$	$\mathbb{1}_{111}[2]$	$\mathbb{1}_{111}[3]$	$\mathbb{1}_{111}[4]$

where $IC(\mathbb{1}_{100})|_{C_{010}}$ as $C_{010} \subsetneq \bar{C}_{100}$.

Now, we turn to analyzing the resolution $\pi : \tilde{C}_{110} \rightarrow \tilde{C}_{110}$ outlined in *Joel's* notes. In order to do this, I hastily jump over to his notation, where now we conceive of the respective 1-by-2 and 2-by-1 sub-matrices x and y as maps $x : E_1 \rightarrow E_q$ and $y : E_{q-1} \rightarrow E_1$. The notation here reflects the conception of these as linear operators between eigenspaces with eigenvalues consisting of the diagonal entries of $\lambda(\mathbf{frob})$. Note too that above I was working with lower triangular Vogans, while Joel is working in the upper triangular case. I will now switch to the upper triangular version of this story to stay consistent with *Joel's* notes, and later edit things so that everything is upper-triangular. That is, from here on out we will write

$$V_\lambda = \begin{bmatrix} 0 & x & 0 \\ 0 & 0 & y \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & x_1 & x_2 & 0 \\ 0 & 0 & 0 & y_1 \\ 0 & 0 & 0 & y_2 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

We describe \tilde{C}_{110} as consisting of all $(Z, E_1^{(1)}) \in V_\lambda \times \text{Gr}(1, E_1)$ such that, writing $Z = (x, y)$

$$x(E_1) \subseteq E_q, \quad x(E_1^{(1)}) \subseteq 0, \quad y(E_{q-1}) \subseteq E_1^{(1)},$$

for some 1-dimensional subspace $E_1^{(1)}$ of E_1 .

We wish verify that this map is indeed semi-small, for this, as well as later computations, we must determine its fibers (at least up to homology and dimension).

First, letting e_i denote the standard basis vectors in \mathbb{C}^4 , we write

$$E_q = \text{span}_{\mathbb{C}}\{e_1\}, \quad E_1 = \text{span}_{\mathbb{C}}\{e_2, e_3\}, \quad E_{q-1} = \text{span}_{\mathbb{C}}\{e_4\}.$$

First, we make the following computation

$$\begin{bmatrix} 0 & * & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 0 \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ d \\ 0 \end{bmatrix}.$$

If we take as a representative Z_{110} of C_{110} which is the above matrix with $*$ = 1, then by the conditions on \tilde{C}_{110} , this forces $E_1^{(1)}$ to be the space on the right, running over all d (and that x_{110} satisfies its condition as well). This exhausts all possible choices, and thus we conclude that

$$\pi^{-1}\{Z_{110}\} \cong \{\text{pt}\}.$$

Consider the representative Z_{010} of C_{010} given now by the above matrix with $*$ = 0, and note that

$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 0 \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ d \\ 0 \end{bmatrix}.$$

By the same reasoning as before, $E_1^{(1)}$ is fixed, and we conclude that

$$\pi^{-1}\{Z_{010}\} \cong \{\text{pt}\}.$$

Now, we compute the action of a representative Z_{100} of C_{100} on E_1 to see that

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ b \\ c \\ 0 \end{bmatrix} = \begin{bmatrix} b \\ 0 \\ 0 \\ 0 \end{bmatrix}.$$

Since we have the condition that $x(E_1^{(1)})$, this forces

$$E_1^{(1)} = \text{span}_{\mathbb{C}}\{e_3\}.$$

Therefore, we again may conclude that

$$\pi^{-1}\{Z_{100}\} \cong \{\text{pt}\}.$$

Finally, we note that for the zero matrix, all the conditions are trivially satisfied, and thus

$$\pi^{-1}\{0\} \cong \text{Gr}(1, E_1) \cong \mathbb{P}_{\mathbb{C}}^1.$$

Now we are in place to verify the map is semi-small. Recall that the condition is that for each orbit $C \leq C_{110}$, we must find that

$$\dim C + 2 \dim \pi^{-1}\{Z\} \leq \tilde{C}_{110},$$

where Z is any representative of C . For now, one will have to intuitively see why $\dim \tilde{C}_{110} = 3$, but I plan to make this rigorous before too long. Observe

$$\begin{aligned} \dim C_{110} + 2 \dim \pi^{-1}\{Z_{110}\} &= 3 + 2 \dim\{\text{pt}\} \\ &= 3 + 2(0) \\ &= 3 \\ &= \dim \tilde{C}_{110}. \\ \dim C_{010} + 2 \dim \pi^{-1}\{Z_{010}\} &= 2 + 2 \dim\{\text{pt}\} \\ &= 2 + 2(0) \\ &< 3 \\ &= \dim \tilde{C}_{110}. \\ \dim C_{100} + 2 \dim \pi^{-1}\{Z_{100}\} &= 2 + 2 \dim\{\text{pt}\} \\ &= 2 + 2(0) \\ &< 3 \\ &= \dim \tilde{C}_{110}. \\ \dim C_{000} + 2 \dim \pi^{-1}\{0\} &= 0 + 2 \dim \mathbb{P}^1 \\ &= 0 + 2(1) \\ &< 3 \\ &= \dim \tilde{C}_{110}. \end{aligned}$$

Therefore π is semi-small, and the only relevant orbit is C_{110} itself. Hence, writing $\tilde{\mathbb{I}} := \mathbb{I}_{\tilde{C}_{110}}$, by the decomposition theorem, we know that there is some non-negative integer m such that

$$\pi_* \tilde{\mathbb{I}}[\dim \tilde{C}_{110}] = mIC_{110}.$$

Taking stalks at $Z_{010} \in C_{010}$, we find that

$$\begin{aligned} m(IC_{110})_{Z_{010}} &= (\pi_* \tilde{\mathbb{I}})_{Z_{010}}[3] \\ mIC_{110}|_{C_{010}} &= H^*(\pi^{-1}\{Z_{010}\})[3] \\ &= H^*(\{\text{pt}\})[3] \\ &= (\mathbb{I}_{010}[0])[3] \\ &= \mathbb{I}_{010}[3]. \end{aligned}$$

Therefore $\mathbb{I}_{010}[3]$ must appear at least once on the left hand side, so $m > 0$. However, if $m > 1$, then it would have to appear on the right hand side more than once, which is a contradiction. Therefore $m = 1$ and

$$IC_{110}|_{010} = \mathbb{I}_{010}[3].$$

An entirely similar argument demonstrates to us that

$$IC_{110}|_{100} = \mathbb{I}_{100}[3].$$

Finally, keeping in mind that we already know $m = 1$,

$$\begin{aligned} (IC_{110})_0 &= (\pi_* \tilde{\mathbb{I}})_0[3] \\ IC_{110}|_{000} &= H^*(\pi^{-1}\{0\})[3] \\ &= H^*(\mathbb{P}^1)[3] \\ &= (\mathbb{I}_{000}[0] \oplus \mathbb{I}_{000}[2])[3] \\ &= \mathbb{I}_{000}[3] \oplus \mathbb{I}_{000}[5]. \end{aligned}$$

At last we are able to complete our stalk table as

m_{geo}^λ	$ _{C_{000}}$	$ _{C_{010}}$	$ _{C_{100}}$	$ _{C_{110}}$	$ _{C_{111}}$
IC_{000}	$\mathbb{I}_{000}[0]$	0	0	0	0
IC_{010}	$\mathbb{I}_{000}[2]$	$\mathbb{I}_{010}[2]$	0	0	0
IC_{100}	$\mathbb{I}_{000}[2]$	0	$\mathbb{I}_{100}[2]$	0	0
IC_{110}	$\mathbb{I}_{000}[3] \oplus \mathbb{I}_{000}[5]$	$\mathbb{I}_{010}[3]$	$\mathbb{I}_{100}[3]$	$\mathbb{I}_{110}[3]$	0
IC_{111}	$\mathbb{I}_{111}[0]$	$\mathbb{I}_{111}[2]$	$\mathbb{I}_{111}[2]$	$\mathbb{I}_{111}[3]$	$\mathbb{I}_{111}[4]$

Reducing to the total dimension in each entry, we obtain

m_{geo}^λ	$ _{C_{000}}$	$ _{C_{010}}$	$ _{C_{100}}$	$ _{C_{110}}$	$ _{C_{111}}$
IC_{000}	1	0	0	0	0
IC_{010}	1	1	0	0	0
IC_{100}	1	0	1	0	0
IC_{110}	2	1	1	1	0
IC_{111}	1	1	1	1	1

Indeed, this is consistent with our previous computations on the spectral side, and the expectations of the p -adic Kazhdan-Lusztig hypothesis.

An immediate generalization

Note that the results of the previous section immediately generalize. That is, consider the cuspidal support

$$S = \{0, \underbrace{1, 1, \dots, 1}_{k\text{-times}}, 2\}.$$

We have the analogous multi-segments

$$\begin{aligned} a_{\max} &= \{0, \underbrace{1, 1, \dots, 1}_{k\text{-times}}, 2\}, a_l = \{[0, 1], \underbrace{1, 1, \dots, 1}_{(k-1)\text{-times}}, 2\}, a_r = \{0, [1, 2], \underbrace{1, 1, \dots, 1}_{(k-1)\text{-times}}\}, \\ a_0 &= \{[0, 1], [1, 2], \underbrace{1, 1, \dots, 1}_{(k-2)\text{-times}}\}, a_{\min} = \{[0, 2], \underbrace{1, 1, \dots, 1}_{(k-1)\text{-times}}\}. \end{aligned}$$

The multi-segment lattice takes on the exact same shape as before. Presumably, it would not be difficult to copy Zelevinsky's computations on the spectral side to reproduce the same multiplicity matrix. However, at the moment the geometry is more clear to me, and thus I will start there and fill in the spectral side later.

Conjecture: In some (all?) cases, it's not even the multiplicities that appear which determine the ranks, it's simply the orbit structure. Or, to put it another way, it's only the *overlaps* that appear.

Again, taking the perspective of the upper-triangular Vogans,

$$V_\lambda = \left\{ \begin{bmatrix} 0 & x & 0 \\ 0 & 0 & y \\ 0 & 0 & 0 \end{bmatrix} : x \in M_{1,k}(\mathbb{C}), y \in M_{k,1}(\mathbb{C}) \right\},$$

and

$$H_\lambda = \left\{ \begin{bmatrix} a & 0 & 0 \\ 0 & B & 0 \\ 0 & 0 & c \end{bmatrix} : a, c \in \mathbb{C}, B \in GL_k(\mathbb{C}) \right\}.$$

The orbits are again given by

$$C_{ijk} = \{(x, y) \in M_{k,1}(\mathbb{C}) \times M_{1,k}(\mathbb{C}) \mid \text{rk}(x) = i, \text{rk}(y) = j, \text{rk}(yx) = k\},$$

with all possible orbits being

$$C_{000}, C_{010}, C_{100}, C_{110}, C_{111},$$

which again, take on the same shape. However in this case, the dimensions can differ. Indeed

$$\dim C_{000} = 0, \quad \dim C_{010} = k = \dim C_{100}, \quad \dim C_{110} = ???, \quad \dim C_{111} = k^2.$$

I suspect that since C_{110} is just C_{111} carved out by a single linear equation, that it's dimension is $k^2 - 1$. I will assume this for now.

As always, we immediately know the upper triangle of the matrix, and since $\bar{C}_{010} \cong \mathbb{A}_{\mathbb{C}}^k \cong \bar{C}_{100}$ and $\bar{C}_{111} \cong \mathbb{A}_{\mathbb{C}}^{k^2}$, we find

m_{geo}^λ	$ C_{000}$	$ C_{010}$	$ C_{100}$	$ C_{110}$	$ C_{111}$
IC_{000}	$\mathbb{1}_{000}[0]$	0	0	0	0
IC_{010}	$\mathbb{1}_{000}[k]$	$\mathbb{1}_{010}[k]$	0	0	0
IC_{100}	$\mathbb{1}_{000}[k]$	0	$\mathbb{1}_{100}[k]$	0	0
IC_{110}				$\mathbb{1}_{110}[k^2 - 1]$	0
IC_{111}	$\mathbb{1}_{111}[0]$	$\mathbb{1}_{111}[k]$	$\mathbb{1}_{111}[k]$	$\mathbb{1}_{111}[k^2 - 1]$	$\mathbb{1}_{111}[k^2]$

Again, we have

$$E_q = \text{span}_{\mathbb{C}}\{e_1\}, \quad E_1 = \text{span}_{\mathbb{C}}\{e_2, \dots, e_{k+1}\}, \quad \text{span}_{\mathbb{C}}\{e_{k+2}\},$$

where $x : E_1 \rightarrow E_q$ and $y : E_{q-1} \rightarrow E_1$, and

$$\tilde{C}_{110} = \{(x, y, E_1^{(1)}) \in V_\lambda \times \text{Gr}(1, E_1) : x(E_1) \subseteq E_q, x(E_1^{(1)}) \subseteq 0, y(E_{q-1}) \subseteq E_1^{(1)}\}.$$

As a representative of C_{110} , consider the matrix Z_{110} such that $x = (1, 0, 0, \dots, 0)$ and $y = (0, 0, \dots, 0, 1)^t$. Then, as before, this forces $E_1^{(1)}$ to be the space with all entries but the penultimate to be zero, hence

$$\pi^{-1}\{Z_{110}\} \cong \{\text{pt}\}.$$

The same arguments carry over to show that

$$\pi^{-1}\{Z_{010}\} \cong \{\text{pt}\} \cong \pi^{-1}\{Z_{100}\},$$

and finally

$$\pi^{-1}\{0\} \cong \text{Gr}(1, E_1) \cong \mathbb{P}^{k-1}.$$

It is certainly much harder to intuit the dimension of \tilde{C}_{110} here, though I suspect it might also be either $\dim \tilde{C}_{110}$ or $(k-1) \times \dim \tilde{C}_{110}$. If the former is true, then I'm wrong about the multiplicity table being the same as before. However in the latter case, there's no relevant orbits, and then I'm stuck on what to do.

If, it is the former case of dimensions, then we will have

$$IC_{110}|_{010} = \mathbb{1}_{010}[k^2 - 1], \quad IC_{110}|_{100} = \mathbb{1}_{100}[k^2 - 1],$$

and

$$IC_{110}|_{000} = \bigoplus_{i=0}^{k-1} \mathbb{1}_{000}[2i].$$

3.3 An explicit description of a supercuspidal representation of $GL(2)$

In this section we will construct an explicit supercuspidal representation ρ of $GL(4)$, and determine its Langlands parameter $\phi := \phi_\rho$, its infinitesimal parameter $\lambda := \lambda_\phi$, the Vogan variety V_λ , and the other (equivalence classes of) Langlands parameters associated to λ .

We fix now a p -adic field, and an extension E/F such that $\text{Gal}(E/F) \cong S_3$. The "standard" irreducible representation $S_3 \rightarrow GL_2(\mathbb{C})$ is given by

$$\begin{aligned} e &\mapsto \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \\ (12) &\mapsto \begin{bmatrix} -1 & -1 \\ 0 & 1 \end{bmatrix} & (123) &\mapsto \begin{bmatrix} -1 & -1 \\ 1 & 0 \end{bmatrix} \\ (23) &\mapsto \begin{bmatrix} 1 & 0 \\ -1 & -1 \end{bmatrix} & (132) &\mapsto \begin{bmatrix} 0 & 1 \\ -1 & -1 \end{bmatrix} \end{aligned}$$

$$\begin{array}{ccc} W_F & \xrightarrow{\quad\quad\quad} & GL_2(\mathbb{C}) \\ & \searrow \quad \swarrow & \\ & W_E/W_F \cong \text{Gal}(E/F) & \end{array}$$

***** for complex numbers $z = re^{i\theta}$ the r is the hyperbolic and the $e^{i\theta}$ is elliptic. In fact, Clifton actually thinks about writing things in terms of powers of q . ***

Anyways, send

$$\sigma \mapsto \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad \tau \mapsto \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Then

$$\begin{aligned} \phi_\rho(frob) &= \phi(\tau^{-1}) \\ &= \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}_{\text{elliptic}}. \end{aligned}$$

For our supercuspidal ρ , we set $\mathfrak{m}_0\{[\rho], [\nu\rho]\}$ and $\mathfrak{m}_1 = \{[\rho, \nu\rho]\}$, then we get

$$0 \rightarrow \pi_1 \rightarrow M_0 \text{Ind}(\rho \otimes \nu\rho) \rightarrow \pi_0 \rightarrow 0.$$

Meanwhile $\pi_1 \cong M_1 = \text{Ind}(\langle[\rho, \nu\rho]\rangle)$.

Now we want to figure out λ , but first lets do the we have not yet wroked it out buuuuuut

Truths to be worked out

For \mathfrak{m}_1 ,

$$\lambda(w) = \phi_{\pi_0}(w, x) = \phi_\rho(w, x) \oplus |w|\phi_\rho(w, x)$$

and

$$\phi_\rho = \text{Ind}_{W_E}^{W_F}(\phi_\psi),$$

and for $\mathfrak{m}_1 = \{[\rho, \nu\rho]\}$

Chapter 4

The groups $SO(n)$

In this section, we carry out analysis of the group $SO_n(\mathbb{C})$, as for a p -adic field F , we have ${}^\vee\mathrm{Sp}_n \cong SO_{2n+1}(\mathbb{C})$ and ${}^\vee\mathrm{SO}_{2n}(F) \cong SO_{2n}(\mathbb{C})$.

To simplify the exposition, we will define the group $SO_n(\mathbb{C})$ with respect to the form $J = {}^tI_n$. We consider an infinitesimal parameter $\lambda : W_F \rightarrow {}^\vee G$ in any case such that ${}^\vee G \cong SO_n(\mathbb{C})$, where

$$\lambda = 1 \oplus \nu \oplus \dots \oplus \nu^{n-1},$$

where as usual, we write ν both for the determinant character of the group, as well as its image under the (local) Langlands correspondence.

Thus, realizing λ as a diagonal matrix, the same analysis may be carried out as in the case of multiplicity free support for $GL(n)$. However, we must now impose the further condition that for all such matrices X , we have

$${}^tXJ + JX = 0,$$

as $V_\lambda \subseteq \mathfrak{so}_n(\mathbb{C})$. Thus, we impose the condition that

$$0 = {}^tXJ + JX$$

$$\begin{aligned} &= \begin{bmatrix} 0 & 0 & \dots & 0 & 0 \\ x_{21} & 0 & \dots & 0 & 0 \\ 0 & x_{32} & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & x_{n,n-1} & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 1 & 0 & \dots & 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 1 & 0 & \dots & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & \dots & 0 & 0 \\ x_{21} & 0 & \dots & 0 & 0 \\ 0 & x_{32} & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & x_{n,n-1} & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & x_{21} \\ 0 & 0 & \dots & x_{32} & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & x_{n,n-1} & \dots & 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & x_{n,n-1} \\ 0 & 0 & \dots & x_{n-1,n-2} & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & x_{2,1} & \dots & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & x_{21} + x_{n,n-1} \\ 0 & 0 & \dots & x_{32} + x_{n-1,n-2} & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & x_{n,n-1} + x_{21} & \dots & 0 & 0 \end{bmatrix}. \end{aligned}$$

Therefore, we find that for all integers i such that $1 \leq i \leq n/2$, we find that $x_{i+1,i} = x_{n-(i-1),n-(i-1)-1}$. In particular, if n is even, this tells us that this forces $x_{n/2+1,n/2} = 0$. An entirely similar computation tells us that for $h = \text{diag}(t_1, \dots, t_n) \in H_\lambda$, when $n = 2m$ we have $h = \text{diag}(t_1, t_2, \dots, t_m, t_m^{-1}, \dots, t_1^{-1})$ and when $n = 2m + 1$ we have $h = \text{diag}(t_1, \dots, t_m, 1, t_m^{-1}, \dots, t_1^{-1})$.

... The moral of the story is that the theory is basically the same here, the stabilizers will be trivial, and the multiplicity table will be immediate.

Chapter 5

The Kazhdan-Lusztig hypothesis for affine Hecke algebras

The goal of this chapter is to understand the statement of *Theorem 8.6.23* of [4], and explicitly describe the notation and the way in which it provides us with an analogue of the Kazhdan-Lusztig hypothesis for affine Hecke algebras.

To begin, we first state the theorem as it appears in [4]. Fix parameters $\psi = (\mathbb{O}_\psi, \chi_\psi)$, $\phi = (\mathbb{O}_\phi, \chi_\phi)$, take $x \in \mathbb{O}_\psi$, and let $i_x : \{x\} \hookrightarrow N$.

Theorem 5.0.1 (Theorem 8.6.23, Chriss & Ginzburg). *The multiplicity of the simple $H_\bullet(Z)$ -module L_ϕ in the composition series of the $H_\bullet(Z)$ -module $H_\bullet(M_x)_\psi$ (resp. $H^\bullet(M_x)_\psi$) is given by*

$$\begin{aligned} [H_\bullet(M_x)_\psi L : LL_\phi] &= \sum_k \dim H^k(i_x^! IC_\phi)_\psi \\ (\text{resp.}) \quad [H^\bullet(M_x)_\psi L : L_\phi] &= \sum_k \dim H^k(i_x^! IC_\phi)_\psi \end{aligned}$$

5.1 Chapter 8 of C+G

Intros

It's claimed that the center of the Hecke algebra \mathbb{H} is

$$Z(\mathbb{H}) \cong R(G)[q, q^{-1}] \cong R(G \times \mathbb{C}^\times),$$

which is certainly believable, but it's not clear that this is proved for p -adic groups G . $\mathbb{C} \otimes_{\mathbb{Z}} Z(\mathbb{H})$ is then identified with the algebra of regular class functions on $G \times \mathbb{C}^\times$, by mapping a character to its representation. Therefore, any algebra homomorphism $Z(\mathbb{H}) \rightarrow \mathbb{C}$ can be identified with the evaluation homomorphism sending $z \in R(G \times \mathbb{C}^\times)$ to $z(a)$ at the semi-simple element $a = (s, t) \in G \times \mathbb{C}^\times$.

Given $a = (s, t) \in G \times \mathbb{C}^*$ semi-simple, we denote \mathbb{C}_a the 1-dimensional subspace as an $Z(\mathbb{H})$ -module, or equivalently a $R(G \times \mathbb{C}^*)$ -module by means of

$$\begin{aligned} R(G \times \mathbb{C}^*) \times \mathbb{C}_a &\rightarrow \mathbb{C}_a \\ (z, x) &\mapsto z(a) \cdot x. \end{aligned}$$

Definition 5.1.1 (8.1.1, [4]). We call the tensor product $\mathbb{H}_a := \mathbb{C}_a \otimes_{Z(\mathbb{H})} \mathbb{H}$ the *Hecke algebra specialized at a* .

Observe that

$$\mathbb{C} \otimes_{\mathbb{Z}} Z(\mathbb{H}) \cong Z(\mathbb{H}_{\mathbb{C}}) \cong Z(\mathbb{C} \otimes_{\mathbb{Z}} \mathbb{H}).$$

The contravariant form

Consider a variety X , which i don't believe need necessarily be complex for the next few statements. Let $Z \subseteq X$ be a locally closed subset, and $i : Z \hookrightarrow X$. For any complex $A \in D^b(X)$, there exists a natural morphism

$$\begin{aligned} \{\text{sections supported on } Z\} &\rightarrow \{\text{sections supported on a nbd of } Z\} \\ i^! A &\rightarrow i^* A, \end{aligned}$$

obtained by applying i^* to the canonical morphism $i_! i^! A \rightarrow A$ by the fact that $i^* i_! (i^! A) \cong i^! A$.

We will generally be taking $A = IC(Y, \mathcal{L})$, with Y a locally closed subvariety of N , and $Z = \{x\}$, with $i_x : \{x\} \hookrightarrow X$.

Let M be a smooth complex algebraic variety, $\mu : M \rightarrow N$ a proper map, and C_M the constant perverse sheaf on M , with direct image $\mu_* C_M$ on N .

Lemma 5.1.2 (8.5.4, [4]). *If M has pure complex dimension m (yet to be defined), then there are natural isomorphisms*

$$H^\bullet(M_x) \cong H^{\bullet-m}(i_x^* C_M), \quad h_\bullet(M_x) \cong H^{m-\bullet}(i_x^! C_M).$$

Suppose now that $\mu : M \rightarrow N$ is G -equivariant, in addition to previous assumptions, and then $N = \coprod \mathbb{O}$ is partitioned into finitely many G -orbits. Write \mathbb{O}_x for the orbit containing the point $x \in N$, let $G(x)$ be the stabilizer of x , and $K(x)$ a maximal compact subgroup.

Let S be a *local transverse slice* (Definition 3.2.19, [4]) and $\tilde{S} := \mu^{-1}(S)$.

Proposition 5.1.3 (8.5.16, [4]). *Given $x \in N$, and irreducible representation χ of $G(x)/G(x)^\circ$, and writing $\phi = (\mathbb{O}, \chi)$ where \mathbb{O} is the orbit of x , then there is a natural isomorphism of vector spaces*

$$L_\phi \cong \text{Im}[H_\bullet(M_x)_\chi \rightarrow H_\bullet(\tilde{S})_\chi].$$

L_x is the "associated multiplicity vector space in $\mu_* C_M$ "???

5.2 The final pieces

Given $x \in N$, and χ an irreducible representation of $G(x)/G(x)^\circ$,

Theorem 5.2.1 (8.6.22, [4]). *Every $H_\bullet(Z)$ -module*

$$L_{x,\chi} := \text{Im}[H_\bullet(M_x)_\chi \rightarrow H_\bullet(\tilde{S})_\chi],$$

is either trivial or simple, and every simple $H_\bullet(Z)$ -module arises in this way.

5.3 Speculation towards the full proof

In this section I speculate on large swaths of the p -KLH which might be proved by the use of the result by Chriss & Ginzburg [4]. We begin by carrying out some exposition from [8].

Let G be a reductive p -adic group, and M a Levi subgroup. Given a supercuspidal representation (σ, E) of M , let \mathfrak{s} be the inertial class it determines.

Let M^1 be the subgroup of M which is generated by all the compact open subgroups. Define $B = \mathbb{C}[M^1/M]$ and $E_B := E \otimes_{\mathbb{C}} B$, and $\Pi := \text{Ind}_P^G(E_B)$. Then, we have an equivalence of categories

$$\begin{aligned} \text{Rep}(G)^{\mathfrak{s}} &\xrightarrow{\mathcal{E}} \text{End}_G(\Pi) - \text{Mod} \\ V &\mapsto \text{Hom}_G(\Pi, V). \end{aligned}$$

For classical groups and inner forms of GL_n , the algebras $\text{End}_G(\Pi)$ are affine Hecke algebras.

It seems promising that this functor is in fact additive and exact, and thus will induce the coefficient preserving map on Grothendieck groups that I think it ought to.

Bibliography

- [1] J. Adams, D. Barbasch and D. A. Vogan, Jr. – The Langlands classification and irreducible characters for real reductive groups, Progress in Mathematics, vol. 104, Birkhauser Boston, Inc., Boston, MA, 1992.
- [2] P. Achar, Perverse Sheaves
- [3] A. Borel – Automorphic L-functions, in Automorphic forms, representations and L-functions, Part 2, p. 27-61 Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979.
- [4] N. Chriss, V. Ginzburg - Representation theory and complex geometry, Modern Birkhauser classics, Birkhauser Boston, 2010.
- [5] C. Cunningham, A. Fiori, A. Moussaoui, J. Mracek, B. Xu - Arthur packets for p -adic groups by way of microlocal vanishing cycles of perverse sheaves, with examples. Memoirs of the American Mathematical Society, Volume 276, Number 1353, 2022.
- [6] J. E. Humphreys, Linear algebraic groups, Springer Verlag New York 1975.
- [7] T. Konno, - A note on Langlands' classification and irreducibility of induced representations of p -adic groups. Kyushu Journal of Mathematics 57(2), January 2003.
- [8] <https://www.math.ru.nl/~solleveld/EndomorphismAlgebras.pdf>
- [9] D. Vogan - The local Langlands conjecture, Representation theory of groups and algebras (J. Adams et al., eds. Contemporary Mathematics 145. American Mathematical Society, 1993.
- [10] A. Zelevinsky - p -adic analogue of the Kazhdan-Lusztig conjecture, Funktsional Anal, i Prilozhen. 15, 1981.
- [11] A. Zelevinsky - Induced representations of p -adic groups II.