

Regulatory coupling across development and technological change

A graded causal framework connecting the RCCX–Hunter–Dąbrowski cascade, perinatal $\alpha_2\delta$ /CaV2 biology, and the polymathic singularity

E. T. Ellis

Research synthesis • 25 September 2026 • Version 0.1

Source boundary. This version integrates two preserved author source manuscripts and a newly verified primary-literature review of the $\alpha_2\delta$ /CaV2 extension. The named *Perinatal Regulatory Initialization / $\alpha_2\delta$ –CaV2 Extension* draft was not available in the local workspace at preparation. Its exact text, diagrams, equations, and author-specific priorities have therefore not been represented as recovered. This is a substantive source gap requiring reconciliation before this version can be called a complete integration of the canonical three-document corpus.

Abstract

This article formalizes a multiscale research program in which inherited variation, conditions around birth, later exposure, physiological regulation, attention, social feedback, and technological tools jointly shape the conversion of human capacity into visible work. Its distinctive claim is not a one-way genotype-to-destiny cascade. It is a set of time-indexed, recursively coupled paths with heterogeneous effects and multiple possible developmental trajectories. We separate four levels of assertion: established mechanisms in bounded systems, supported syntheses across studies, proposed mechanisms requiring direct tests, and horizon hypotheses about long-run cultural selection. The RCCX region contains well-characterized genes and clinically demonstrated CYP21A2–TNXB contiguous-gene disease, but the proposed RCCX-to-attention-to-creativity bridge remains untested. $\alpha_2\delta$ -1 regulates CaV2 trafficking in specific systems and also participates in thrombospondin-associated synaptogenesis; neither result demonstrates that ordinary perinatal exposures initialize a later regulatory phenotype. Human stress-epigenetic observations are informative but mixed and do not establish direct intergenerational methylation transfer. Event-rate experiments and online foraging tasks support environment-sensitive attention in bounded settings, while AI-assisted short-story experiments show that individual gains can coexist with reduced collective diversity. We propose three discriminating study designs and a source-linked edge ledger that make the theory falsifiable without mistaking background biology for proof of an unsupported bridge.

Keywords: regulatory coupling; development; RCCX; CACNA2D1; CaV2; $\alpha_2\delta$; attention; heterogeneity; human–AI interaction; causal inference

1. Scope and claim grammar

The RCCX–Hunter–Dąbrowski Cascade v3.2 places constitution, ecological fit, stress, autonomic state, attention, adaptation, identity, reorganization, and historical output in one developmental map. *The Polymathic Singularity* v0.1 extends the map to institutions and AI-mediated production. The third named draft would add a perinatal $\alpha 2\delta$ /CaV2 layer; its exact author text is pending recovery. The present paper tests whether these layers can be connected without laundering uncertainty between them.

Four claim classes govern every interpretation. **Established mechanism** means the stated causal relation was directly manipulated or otherwise strongly identified in the cited system and outcome. **Supported synthesis** combines findings with an explicit transfer boundary. **Proposed mechanism** names a causal path that requires a direct test. **Horizon hypothesis** predicts a broader developmental or civilizational pattern beyond present identification. Grades A through D operationalize this distinction in the accompanying edge ledger: A is direct evidence in the stated system; B is convergent but bounded; C is plausible synthesis; D is an unmeasured bridge or horizon claim. A grade attaches to an *arrow*, never to an evocative node name.

The practical translation of the theory is simple: a person's behavior depends partly on the kind of feedback their body and attention can use, the environments available to them, and what happens after they act. The model does not make “Hunter,” “Farmer,” “gifted,” “traumatized,” “ADHD,” or “polymath” into biological kinds. Those terms refer to different constructs and scales. They must be measured separately.

2. The bounded molecular and developmental starting points

2.1 RCCX: real locus, proposed cascade

The RCCX module includes C4, CYP21A2, TNXB, and neighboring repeat structures. Recombination involving CYP21A2 and TNXB can cause the clinically defined CAH-X contiguous-gene syndrome [S01]. This is a strong genetic and clinical anchor for a *specific* endocrine–connective-tissue link. It is not evidence that ordinary RCCX variants create an integrated attention, trauma, and creativity phenotype. Complement C4 structural variation has been associated with schizophrenia risk, and mouse work implicated postnatal synapse elimination [S02]; that result concerns another phenotype and cannot be treated as proof of the broad cascade. ADHD genome-wide analysis identified 27 risk loci [S03], a direct warning against a single-locus etiological story.

The strongest defensible formulation is a subgroup hypothesis: measured RCCX structural variants may, in some people, alter endocrine, immune, or connective-tissue constraints that interact with autonomic and developmental conditions. Selection into hEDS/HSD clinics matters. In one cohort of 102 adults, 48% met a POTS definition [S16]. That figure describes the selected cohort, not all people with hypermobility or the general population. Genotype, diagnosis, symptoms, age, medication, and recruitment must be recorded before attributing attention differences to this pathway.

2.2 Perinatal $\alpha 2\delta$ /CaV2: two functions and a missing bridge

$\alpha 2\delta$ proteins are auxiliary calcium-channel subunits, with isoforms encoded by different CACNA2D genes. CaV2 channels are pore-forming voltage-gated calcium channels important for synaptic transmission. A mouse nerve-injury experiment showed $\alpha 2\delta$ -1 was required for increased CaV2.2 localization in particular sensory neurons and terminals [S11]. In a different experimental system, $\alpha 2\delta$ -1 was identified as a participant in thrombospondin-induced excitatory synapse formation; the new synapses and drug effects were studied in rodent cells and in vivo models [S10]. These are distinct molecular roles. “ $\alpha 2\delta$ affects CaV2 trafficking” and “ $\alpha 2\delta$ participates in synaptogenesis” must not be collapsed into a single pathway.

Stage and tissue are decisive. Mouse brain expression of channel subunits changes by region and developmental stage; in one profiling study, blocking electrical activity did not change the measured expression pattern [S12]. Two reported humans with severe biallelic CACNA2D1 loss-of-function variants had early epileptic encephalopathy, and functional assays showed impaired CaV2 trafficking [S18]. Rare severe variants cannot establish the effects of common variation, stress exposure, or an adult attention style. The proposed arrow from perinatal conditions to $\alpha 2\delta$ /CaV2 timing, and thence to later regulatory coupling, is **grade D**. It earns a place in a research program because its components are measurable, not because the bridge has been shown.

An internal contrary result sharpens the mechanism. A biochemical study reported that gabapentin did not directly inhibit measured binding between soluble $\alpha 2\delta$ -1 and thrombospondin-4 [S13]. This does not negate the earlier observation that gabapentin inhibited synapse formation in the model [S10]. It prevents the simple explanation that the drug necessarily blocks the ligand–receptor binding event itself. Ligand specificity, cell context, channel effects, and downstream steps must be distinguished.

2.3 Stress programming and intergenerational limits


In selected postmortem human hippocampal samples, childhood abuse history was associated with differences in NR3C1 promoter methylation and glucocorticoid receptor expression [So4]. This supports a biologically plausible route between exposure and stress-response biology, but the design does not identify a universal causal sequence. A study of 113 mother–infant pairs found maternal associations while finding no direct transmission of the assayed methylation patterns in newborns [So5]. A separate Holocaust-offspring study replicated an FKBP5 site association in an independent sample of 125 offspring and 31 controls, with maternal developmental programming proposed as one explanation [So6]. A population sample of 3,965 adults did not replicate previously reported childhood-maltreatment interactions at assayed FKBP5 sites [S17]. Tissue, cell mixture, site selection, genotype, shared environment, and timing may all change results. Therefore, “stress is inherited as methylation” is too strong. The defensible claim is that early environments can become biologically associated with later regulation, while the exact causal and intergenerational routes remain unsettled.

3. A time-indexed causal model

A single directed acyclic graph cannot contain a literal feedback cycle. We represent recursion by indexing the same variables over time. At time t , constitutional and perinatal variables precede measured cellular, stress, and attentional mediators. Attention and action alter selected contexts at $t +$

1, whose feedback can then alter physiology and later attention. This preserves the central recursion without hiding simultaneity behind an arrow diagram.

Within-window DAG:



Feedback across windows: $A_t \rightarrow E_{(t+1)} \rightarrow S_{(t+1)} \rightarrow A_{(t+1)}$. Social adaptation can mediate $A_t \rightarrow E_{(t+1)}$. AI access can modify $E_t \rightarrow C_t$; outputs and institutional response may change later opportunity, $Y_t \rightarrow E_{(t+1)}$. These are distinct empirical questions. The illustrated graph is an organizing hypothesis, not a claim that all paths are active in every person.

Potential common causes include family resources, parental genotype, prenatal health, medication, age, sex, educational opportunity, and the selection process that places a participant in a clinic or study. Some mediators are colliders under plausible alternative graphs. Cross-sectional conditioning on diagnosis, for example, can create spurious genotype–environment associations. The complete analysis plan must declare exposure windows, sampling, measurement sites, and adjustment sets before fitting path models.

One possible parameterization of the author's regulatory closure construct is $R_c = \log(G_{\text{exo}} / G_{\text{endo}})$, where G_{exo} is within-person performance or regulatory improvement produced by usable environmental feedback, and G_{endo} is improvement under matched internally maintained control. These are *experimental contrasts*, not intrinsic quantities that can be read from a diagnosis. Gains must share a scale and be greater than zero; when either is near zero, use a predeclared bounded or difference-score alternative. Repeated tasks with matched difficulty, arousal, and reward are needed to separate feedback dependence from task preference.

The author's conversion model distinguishes potential, organization, execution resources, and dissemination. It is best treated as a vector of necessary and substitutable constraints, not as a literal multiplication of unmeasured “magnitudes.” Historical visibility is an outcome with strong institutional selection. A person may have high ability and no opportunity, or substantial output that is not publicly counted. Such cases are predicted by the model and are crucial controls, not exceptions to discard.

4. Behavioral and social bridges

The 2012 event-rate meta-analysis found disproportionate slowing in ADHD groups at slow event rates and larger commission-error effects at fast rates, while general ADHD–control differences persisted and reaction-time variability did not show the predicted event-rate effect [S07]. This supports a bounded environment-sensitive performance mechanism. It does not establish two fixed regulatory human types. In a preregistered online foraging task involving 457 participants, people crossing a positive ADHD screening threshold left depleting resource patches sooner than those below the threshold [S08]. That result concerns a task strategy and a screening measure. It does not demonstrate a hunter genotype, an ancestral adaptation, or superior performance in every environment.

The Hadza actigraphy study found one or more people scored awake or in light sleep during 99.8% of sampled nighttime epochs across 20 days [S09]. The measured phenomenon was group-level chronotype and waking heterogeneity. It cannot be transferred to ADHD, a perinatal channel mechanism, or modern work arrangements without further study. It does, however, make a precise ecological point: variation between individuals can create a group-level function that would be invisible if everyone were forced into one schedule.

Autistic social camouflaging has a validated adult self-report instrument [S15]. That establishes a measurable phenomenon in an identified population. The broader cascade's “homeostatic masking” remains a proposed generalization and should be studied with longitudinal behavior, physiology, and context rather than inferred from personality descriptions. Dąbrowski's overexcitabilities and positive disintegration provide a developmental vocabulary for intensity and reorganization. They are theoretical constructs here. No source in this ledger validates a binary fork from distress to either fragmentation or genius.

5. Human–AI coupling and the horizon claim

AI can supply an external feedback surface, lower some drafting and implementation costs, and alter which work gets evaluated. In a randomized short-story experiment, access to AI-generated ideas improved average ratings of individual stories while stories became more similar to one another [S14]. The two results matter together. More individual output need not imply a more diverse collective frontier. Neither the study nor any source here shows that an identified “polymathic” phenotype uniquely benefits, that AI causes a wholesale economic inversion, or that deep specialization will lose value.

The *polymathic singularity* is therefore a grade-D horizon hypothesis: technological and institutional changes may increase the payoff to people who can integrate domains, use tools, maintain coherence, and convert work into public artifacts. It predicts heterogeneous, context-dependent effects. It also predicts failure modes: dependence on brittle tools, homogenized ideas, unequal access, amplified distraction, and institutional selection that continues to favor credentials or narrow expertise. The relevant unit may be a team of integrators, specialists, and machines rather than a heroic individual. A proper test measures composition, output quality, novelty, and durable adoption separately.

6. Contrary evidence and boundaries that change the conclusion

Claim under pressure	Counterweight	Consequence
RCCX explains broad attention phenotypes	ADHD has a distributed genetic architecture with 27 loci [S03].	Test a predefined RCCX subgroup; do not use the locus as a general cause.
Perinatal exposure controls $\alpha_2\delta$ /CaV2	Mouse channel expression showed stage and region specificity and no change under one activity-blockade condition [S12].	Name exposure, tissue, isoform, stage, and mediator before claiming programming.
Gabapentin blocks TSP- $\alpha_2\delta$ binding	Direct soluble TSP4 binding was gabapentin-insensitive [S13].	Separate binding, trafficking, and downstream synaptogenic effects.
Maternal trauma transfers methylation	A 113-dyad study found no direct transmission at assayed sites [S05]; a population study did not replicate a maltreatment interaction [S17].	Measure common causes and tissue-specific routes; avoid inherited-mark shorthand.
ADHD-related exploration is universally adaptive	The foraging result came from one online task with a screener [S08].	Estimate phenotype-by-environment effects across multiple tasks.
AI expands creative diversity	AI-assisted short stories were more similar across writers [S14].	Measure collective novelty and individual quality separately.

The source literature spans mice, cell cultures, rare human variants, selected clinics, postmortem brain, adult surveys, online tasks, and short-story experiments. There is no license to move an effect between species, tissue, developmental stage, sampling frame, or outcome merely because the vocabulary sounds continuous. The arrow ledger records each such boundary.

7. Minimum discriminating studies

Study 1 — perinatal mediator test. In a longitudinal birth cohort with repeated exposure measures, genotype, medication, gestational age, and maternal health, estimate whether a specified perinatal exposure predicts a prespecified $\alpha_2\delta$ /CaV2 measure in a relevant human biospecimen or validated model, then later regulatory outcomes. A mechanistic arm could use human-derived cellular models with controlled exposure and isoform-specific perturbation. **Discriminator:** mediation should be isoform-, stage-, and tissue-specific and survive family and cell-composition controls. Failure to find the mediator where and when predicted weakens the extension even if exposure and later behavior correlate.

Study 2 — regulatory closure cross-over. Randomize task environments within participants: feedback density, event rate, movement permission, delay, and external representation, with matched task difficulty and reward. Measure performance, state, and recovery repeatedly in people sampled across attention traits and autonomic profiles. Preregister the **R_c** estimation and interaction contrasts. **Discriminator:** if exoregulatory gain is a distinct phenotype, its rank should show within-person stability across several task families and predict cross-over effects beyond diagnosis, arousal, and preference. If the index collapses after these controls, the proposed orientation has not earned construct status.

Study 3 — conversion and diversity. Randomize access to AI assistance and scaffold type in multidisciplinary teams; stratify by measured integration skills rather than self-described polymathy. Score completed artifacts blind for technical quality, originality, cross-domain integration, and independent adoption over time. Track within-team specialization and collective similarity. **Discriminator:** the horizon model predicts an interaction between integrative capacity, scaffold quality,

and conversion resources, not just a general main effect of AI. A universal AI benefit or increasing output with no phenotype interaction would favor simpler explanations; homogenization would count against a strong collective-diversity claim.

These studies should publish null results and adverse effects. They must not assign an individual's worth or clinical treatment from an unvalidated index. Claims about perinatal intervention would require independent safety and efficacy evidence beyond this theory.

8. Discussion and conclusion

The model's useful novelty is its architecture: constitution constrains, development tunes, action selects environments, environments feed back, and tools change conversion opportunities. The strongest empirical pieces live at different scales. The cross-scale bridges remain mostly C or D. Naming that gap makes the project testable. It also preserves the author's central insight: a diagnosis is one possible readout of a coupled history, while publicly visible achievement depends on conversion and dissemination as well as ability.

The decisive scientific task is to identify which arrows operate for which people, at which stages, in which environments, and with what alternative explanations ruled out. Until then, the RCCX-centered pathway, perinatal $\alpha_2\delta$ /CaV2 initialization, regulatory closure phenotype, and polymathic singularity are connected hypotheses with graded evidence, not a single established mechanism.

References

The identifiers and direct links below match the repository's machine-readable source ledger. Source manuscripts are preserved byte-for-byte in `sources/original/` with SHA-256 hashes.

- [S01] Kim et al. (2023). Molecular basis and genetic testing strategies for diagnosing 21-hydroxylase deficiency, including CAH-X syndrome. PubMed.
- [S02] Sekar et al. (2016). Schizophrenia risk from complex variation of complement component 4. *Nature*. PubMed.
- [S03] Demontis et al. (2023). Genome-wide analyses of ADHD identify 27 risk loci. *Nature Genetics*. PubMed.
- [S04] McGowan et al. (2009). Epigenetic regulation of the glucocorticoid receptor in human brain associates with childhood abuse. *Nature Neuroscience*. PubMed.
- [S05] Ramo-Fernández et al. (2019). Effects of childhood maltreatment on epigenetic regulation: an intergenerational approach. *Scientific Reports*. PubMed.
- [S06] Bierer et al. (2020). Intergenerational Effects of Maternal Holocaust Exposure on FKBP5 Methylation. *American Journal of Psychiatry*. PubMed.
- [S07] Metin et al. (2012). A meta-analytic study of event-rate effects on Go/No-Go performance in ADHD. *Biological Psychiatry*. PubMed.
- [S08] Barack et al. (2024). Attention deficits linked with proclivity to explore while foraging. *Proceedings of the Royal Society B*. PubMed.
- [S09] Samson et al. (2017). Chronotype variation drives night-time sentinel-like behaviour in hunter-gatherers. *Proceedings of the Royal Society B*. PubMed.
- [S10] Eroglu et al. (2009). The gabapentin receptor $\alpha_2\delta$ -1 is a neuronal thrombospondin receptor responsible for excitatory CNS synaptogenesis. *Cell*. PubMed.
- [S11] Nieto-Rostro et al. (2023). Nerve injury increases native CaV2.2 trafficking in dorsal root ganglion mechanoreceptors. *Pain*. PubMed.
- [S12] Schlick et al. (2010). Voltage-activated calcium channel expression profiles in mouse brain and cultured hippocampal neurons. *Neuroscience*. PubMed.
- [S13] El-Awaad et al. (2019). Direct, gabapentin-insensitive interaction of soluble $\alpha_2\delta$ -1 with thrombospondin-4. *Scientific Reports*. PubMed.
- [S14] Doshi and Hauser (2024). Generative AI enhances individual creativity but reduces the collective diversity of novel content. *Science Advances*. DOI.
- [S15] Hull et al. (2019). Development and Validation of the Camouflaging Autistic Traits Questionnaire. *Journal of Autism and Developmental Disorders*. PubMed.
- [S16] Celletti et al. (2020). Postural tachycardia syndrome in 102 adults with hEDS/HSD. *Monaldi Archives for Chest Disease*. PubMed.
- [S17] Klinger-König et al. (2019). Methylation of the FKBP5 gene in association with FKBP5 genotypes, childhood maltreatment and depression. *Neuropsychopharmacology*. PubMed.
- [S18] Dahimene et al. (2022). Biallelic CACNA2D1 loss-of-function variants cause early-onset developmental epileptic encephalopathy. *Brain*. PubMed.