

Pedometer-Measured Physical Activity Patterns of Youth

A 13-Country Review

Michael W. Beets, MPH, PhD, Daniel Bornstein, BS, Aaron Beighle, PhD,
Bradley J. Cardinal, PhD, Charles F. Morgan, PhD

Context: Insufficient physical activity among young people aged 5–18 years is a global public health issue, with considerable disparities among countries. A systematic review was conducted to identify studies reporting pedometer daily steps ($\text{steps}\cdot\text{day}^{-1}$) in order to compile comparative, global cross-sectional data on youth physical activity patterns.

Evidence acquisition: Articles were included if they were in English, published by April 2009, and reported $\text{steps}\cdot\text{day}^{-1}$ for boys and girls, separately, and reported $\text{steps}\cdot\text{day}^{-1}$ for age groupings of no more than 4 years (e.g., 5–8 years) or combined no more than three grade levels (e.g., third- to fifth-graders). Studies could have been intervention-based but had to have reported baseline $\text{steps}\cdot\text{day}^{-1}$, which would reflect unadulterated physical activity $\text{steps}\cdot\text{day}^{-1}$ estimates. Inverse variance weighted estimates ($\text{steps}\cdot\text{day}^{-1w}$) were calculated for each country, and random effects models were estimated. Analyses were conducted in May and June 2009.

Evidence synthesis: Forty-three studies, representing young people in 13 countries ($N=14,200$), were included. The majority of studies were from the U.S. (17/43). Overall, there was considerable variation within and among countries in $\text{steps}\cdot\text{day}^{-1w}$. Boys and girls from European and Western Pacific regions had significantly more $\text{steps}\cdot\text{day}^{-1w}$ than young people from the U.S. and Canada. Significantly lower $\text{steps}\cdot\text{day}^{-1w}$ estimates for girls were observed for studies that combined measured $\text{steps}\cdot\text{day}^{-1}$ for weekdays and weekend days, in comparison to weekdays only.

Conclusions: Limited sample sizes and non-population-based data preclude definitive statements regarding projected $\text{steps}\cdot\text{day}^{-1}$ within countries. Nevertheless, these findings provide preliminary information for policymakers and researchers on the extent of the disparities among countries in the physical activity patterns of young people.

(Am J Prev Med 2010;38(2):208–216) © 2010 American Journal of Preventive Medicine

Introduction

The global health burden of physical inactivity is increasingly recognized as a major public health problem for both developing and developed countries.^{1,2} Declines in physical activity are intricately linked to increases in childhood and adolescent

obesity^{3–7} and metabolic syndrome,⁸ and they parallel the rapid secular decline in youth cardiovascular fitness.⁹ Worldwide, 1.9 million deaths annually are attributable to noninfectious diseases related to physical inactivity.² Although there is little doubt that physical activity plays an important role in reversing these trends, public health practitioners, despite best efforts, have made little headway.^{10,11}

Globally, the prevalence of young people (aged 5–18 years) who are physically inactive, in conjunction with the soaring prevalence of obesity, is a major threat to health in the 21st century.² However, considerable disparity exists among countries in the physical activity levels of young people. In some countries, physical activity levels are notably high, and hence, the young people are on average healthier.^{12,13} Conversely, in some countries, young people consistently demonstrate low levels of

From the Department of Exercise Science (Beets, Bornstein), Arnold School of Public Health, University of South Carolina, Columbia, South Carolina; Department of Physical Education (Beighle), University of Kentucky, Lexington, Kentucky; Department of Nutrition and Exercise Sciences (Cardinal), Oregon State University, Corvallis, Oregon; and Kinesiology and Leisure Science Department (Morgan), University of Hawaii, Honolulu, Honolulu, Hawaii

Address correspondence and reprint requests to: Michael W. Beets, MPH, PhD, Department of Exercise Science, Arnold School of Public Health, University of South Carolina, 921 Assembly Street, Room 131, Columbia SC 29208. E-mail: beets@mailbox.sc.edu.

0749-3797/00/\$17.00

doi: 10.1016/j.amepre.2009.09.045

physical activity, high levels of obesity, and hence are on average less healthy.^{3,12–14} The focus here is not on whether the former countries should be emulated, but rather on important lessons gleaned from their experience that may benefit “less healthy” countries.

Public health advocates and policymakers gain from cross-national comparisons in several important ways. Comparisons help identify global rankings of physical activity and inactivity patterns. These rankings assist with establishing priorities for improving physical activity levels and serve as a benchmark for physical activity within individual countries. Population-level data can also serve to inform policymakers regarding trends in physical activity and inactivity patterns over time. As mentioned previously, comparisons afford the opportunity to identify model countries that can inform public health planning in less-active countries. Yet, aside from a few studies,^{12,13,15–17} not many global comparisons of physical activity patterns are available.

Global problems require global solutions, or at a minimum, global studies.¹⁸ Often the spark that ignites global awareness of the magnitude of an issue is the compilation of existing studies.^{6,13,19} Comprehensive reviews, therefore, provide useful information regarding the patterning of physical activity among countries.¹³ Efforts to compile such reviews, however, are complicated by the many forms of measurement used in physical activity data. In the absence of standardized instrumentation (e.g., the International Physical Activity Questionnaire [IPAQ]),¹⁶ aggregating activity levels from either self-report or objective measures (e.g., accelerometers) is methodologically difficult because of differences in the units reported (e.g., estimated energy expenditure from varying formulas, differences among accelerometer cutpoints).^{13,20,21} Thus, there is little information on global patterns of youth physical activity.

The use of pedometers to quantify physical activity has grown over the past decade. Pedometers also provide one of the few forms of physical activity assessment with a standard metric (most commonly, steps-day⁻¹). Pedometers, therefore, are one of the few valid and reliable objective measures of youth physical activity²² that allow direct comparisons across studies. This aspect of pedometer use has led to a growing body of literature reporting daily physical activity levels of young people among numerous countries, in addition to the development of guidelines for recommended steps-day⁻¹ for young people. Recommendations range from 11,000 to 16,500 steps-day⁻¹ for boys and girls.^{23–26} Data on the extent to which young people from various countries achieve recommended pedometer activity levels are limited, however, and comparisons among countries are relatively few.^{12,25} The purpose of the current study was to compile

and explore physical activity patterns of young people (≤ 18 years) from studies that measured physical activity via pedometry from around the world.

Evidence Acquisition

A systematic review of the literature was conducted between December 2008 and April 2009 to identify all peer-reviewed papers published by April 2009 that reported daily pedometer-derived steps (i.e., steps-day⁻¹) of children and adolescents (≤ 18 years). A Boolean search for relevant articles was conducted in PubMed, EBSCOhost, and Google Scholar with pedometer* and the following combination of keywords: child*, adolescent*, and youth*. Additional searches were carried out on citations of included papers and published reviews on youth physical activity and pedometers. Articles were included if they were published prior to May 2009 and reported steps-day⁻¹ for boys and girls, separately, for age groupings of no more than 4 years (e.g., 5–8 years) or combined no more than three grade levels (e.g., third- to fifth-graders). Studies could have been intervention based but had to have reported baseline steps-day⁻¹, which would reflect physical activity steps-day⁻¹ estimates that were unaffected by an intervention. Analyses were conducted in May and June 2009.

Studies were excluded if they reported only steps-day⁻¹ for youth divided into BMI categories (i.e., overweight versus healthy weight)^{25,27}; data were previously published and analyzed for a secondary purpose²⁸; data represented overweight youth only; data for age ranges were combined for more than 4 years or three grade levels; data for boys and girls were combined²⁹; data were collected for less than 1 complete day (e.g., physical education class or recess only); or data were collected on youth from a special population (e.g., indigenous peoples, Amish).^{30,31}

Information was extracted from identified articles, and data were entered into specially designed templates for analysis. The following information was extracted from each article: country; days on which data were monitored (i.e., weekdays, weekend days, or weekdays plus weekend days); sample size; age; gender; type of pedometer used; and mean steps-day⁻¹ and SD. Articles in which seasonal variations were of interest, the spring/fall season was selected as representative and was included in the review, because most data collection occurs during these times.

Aggregation of Data and Statistical Analysis

For ease of interpretation, when studies reported the mean age of the sample, age was rounded to the nearest whole year (e.g., 9.5 to 10 years; 11.4 to 11 years). Where age of the sample was not reported, grade level was used to infer the average expected age for a given grade-level range. For studies within each country, the mean steps-day⁻¹; sample size (n); and SD were used to create an inverse variance sample-weighted estimated steps-day^{-1w} by weighting the contribu-

tion of each study’s mean steps-day^{−1} by the sample size divided by the variance (i.e., square of the SD).³² For studies where the SDs of steps-day^{−1} were not reported (6/43 studies), the SD was estimated by predicting the sample-weighted coefficient of variance (CV) for all studies with the mean steps-day^{−1}, gender, and age. The predicted CV was then used to determine the SD of those studies by multiplying the mean steps-day^{−1} by the predicted CV. For countries where multiple studies reported mean steps-day^{−1} for a single age and gender (e.g., girls aged 8 years), the reported steps-day^{−1} value was multiplied by its respective inverse variance weight; then the values were summed and divided by the sum of the weights to arrive at the steps-day^{−1w} estimate. Studies were divided based on the inclusion of weekend days in the total steps-day^{−1} versus weekdays only. Across studies within a country for a given age (whole years), steps-day^{−1w} were calculated and plotted versus age.

Random effects models, accounting for the clustering of reported steps-day^{−1} within studies (i.e., a single study could have reported multiple age–gender steps-day^{−1} values), were used to estimate the average sample-weighted steps-day^{−1w} across the age range (5–18 years) for all studies for boys and girls, and for weekdays versus weekdays plus weekend days, separately. Age, age squared, and the square root of age were included in the models to account for the nonlinear change in steps-day^{−1} from 5 to 18 years. Random effects models were used to test the difference in weekday only versus weekday plus weekend-day data collection. Tests of significance were also conducted to determine if there was a difference in steps-day^{−1w} for length of pedometer assessment (i.e., number of days). In addition,

steps-day^{−1w} for each age–gender strata within each country were compared to established pedometer steps-day^{−1} guidelines for young people aged 5–12 years^{24–26} and those aged 5–19 years.²³

Comparisons among individual countries were not possible because of the limited sample sizes and ages represented in the majority of the studies. Therefore, countries were placed into regional groupings based on WHO³³ regional distributions in order to compare steps-day^{−1w}. The groupings were as follows: Western Pacific (Australia and New Zealand); Europe (Sweden, Switzerland, Belgium, France, Greece, Czech Republic, and the United Kingdom); and the Americas (Canada and the U.S.). Two countries (China and Saudi Arabia) were the only ones from their WHO regions (Southeast Asia and Eastern Mediterranean, respectively) and therefore were grouped together in an “Other” category in the models. Regional groupings were dummy-coded and entered into the random effects models. The region containing the U.S. and Canada was used as the reference group for these analyses because, in general, it produced some of the lowest steps-day^{−1w} values across age–gender comparisons. All analyses were conducted using Mplus, version 5.2 (www.statmodel.com).

Evidence Synthesis


The database searches yielded a total of 356 hits across all combinations of keywords. After excluding articles that did not meet inclusion criteria (see above), a total of 43 studies, representing 13 countries and approximately

Table 1. Included country characteristics and representation

Region	Country	No. of studies	References	Ages represented (years)														Sample size
				5	6	7	8	9	10	11	12	13	14	15	16	17	18	
North America	Canada	1	34						•	•								133
	U.S.	17	26, 35–50		•	•	•	•	•	•	•	•	•	•	•	•	•	4,228
Europe	Belgium	1	51		•													122
	Czech Republic	1	52			•	•											176
	France	1	53						•		•	•	•	•	•			478
	Greece	1	54								•	•						221
	Sweden	4	55–58			•	•	•	•	•	•	•	•		•		•	1,845
	Switzerland	1	59									•						233
	United Kingdom	6	60–65					•	•									1,255
Other	China	1	66									•	•	•				144
	Saudi Arabia	2	67, 68		•			•	•		•							520
Western Pacific	Australia	4	69–72				•		•		•		•	•	•			2,222
	New Zealand	3	73–75			•	•	•		•		•		•				2,623
Total		43			•	•	•	•	•	•	•	•	•	•	•	•	•	14,200

Results from the random effects models comparing steps·day^{-1w} from studies reporting weekday only versus weekday plus weekend steps·day^{-1w} indicated that for

The final weighted steps-day^{-1w} estimates for boys and girls by age (whole years) are presented in [Figure 1](#) (week-days only) and [Figure 2](#) (weekdays plus weekend days). Considerable variation among countries in steps-day^{-1w} can be seen for both genders and monitoring time frames (see [Figures 1](#) and [2](#)).



Czech, Czech Republic; NZ, New Zealand; SA, Saudi Arabia; Swtz, Switzerland; Swed, Sweden; UK, United Kingdom




Figure 2. Weighted steps·day^{-1w} for studies reporting weekday plus weekend-day steps across countries by age for boys (●, black circles) and girls (○, white circles), separately

Note: Curved lines represent the estimated average steps·day^{-1w} from all studies for boys (—, solid line) and girls (---, dashed line), separately.

^aSteps·day⁻¹ guidelines (from Canada²³) for boys and girls aged 5–19 years, based on an equivalence to 90 minutes of moderate to vigorous physical activity

^bSteps·day⁻¹ guidelines (from New Zealand²⁴) for boys and girls aged 5–12 years, based on bioelectric impedance body fat percentage

^cSteps·day⁻¹ guidelines (from Sweden, Australia, and the U.S.²⁸) for boys and girls aged 5–12 years, based on BMI

^dSteps·day⁻¹ guidelines (from the U.S.²⁶) for boys and girls aged 5–12 years, based on mean values

Czech, Czech Republic; NZ, New Zealand; SA, Saudi Arabia; Swtz, Switzerland; Swed, Sweden; UK, United Kingdom

For weekday steps·day^{-1w} (Figure 1), a notable difference was observed for boys from Sweden, where steps·day^{-1w} are greater than they are in almost all other countries, apart from one (New Zealand, for those aged 9 years). For girls, a similar pattern is observed. In addition, boys and girls from the U.S. have, across the age continuum, the lowest or next-to-lowest steps·day^{-1w} values in comparison to all other countries. Random effects models comparing differences among regions supported these observations, with boys and girls from European countries, on average, taking, respectively, on average 2389 and 2375 steps·day^{-1w} more than young people from the

U.S. and Canada. In addition, Western Pacific girls took 3340 steps·day^{-1w} more than girls from the U.S. and Canada. No comparisons among boys from the Western Pacific region could be made for weekday steps·day^{-1w} because of insufficient representation of boys from this region. No difference was observed between young people from the Other region and those from the U.S. and Canada.

For weekday plus weekend-day steps·day^{-1w} estimates, the ordering of countries was less clear (Figure 2). Boys from Australia and New Zealand consistently demonstrated some of the highest steps·day^{-1w} values in compari-

Table 2. Weighted steps-day^{-1w} comparisons for studies reporting weekdays only and weekdays plus weekend days among geographic regions for boy and girls, separately

Gender	Measurement	Region	Steps-day ⁻¹ (95% CI) ^w
Boys	Weekdays	Other	606 (–1154, 2366)
		Europe	2389 (1060, 3717)
		Western Pacific ^a	NA
	Weekdays + weekend days	Other	–1209 (–3228, 811)
		Europe	–1210 (–2684, 265)
		Western Pacific	1839 (288, 3390)
Girls	Weekdays	Other	–310 (–3687, 3067)
		Europe	2375 (1351, 3399)
		Western Pacific	3340 (2440, 4240)
	Weekdays + weekend days	Other	38 (–2216, 2291)
		Europe	198 (–1053, 1448)
		Western Pacific	1604 (318, 2889)

Note: WHO regional groupings: Other = China and Saudi Arabia; Europe = Sweden, United Kingdom, Belgium, Czech Republic, France, Greece, Switzerland; Americas = Canada and U.S. (comparison group). Random effects models controlled for the linear and nonlinear effects of age on steps-day^{-1w}.

^aUnable to estimate because only one study for this region was included in the review reporting steps-day⁻¹ for this time frame

son to all other countries, apart from two (U.S. boys aged 8 years, United Kingdom boys aged 10 years, and boys from Greece aged 12 years). A similar pattern was observed for girls. The Western Pacific region had, on average, 1839 and 1604 steps-day^{-1w} more, respectively, than boys and girls from the U.S. and Canada (see Table 2).

Average sample-weighted steps-day^{-1w} for those aged 5–18 years across both monitoring time frames suggested similar patterns in activity, with the most steps-day^{-1w} observed for those aged 9–13 years, with a marked decline occurring thereafter. The difference between the two estimates occurs at the youngest ages (<8 years), where for weekdays, steps-day^{-1w} values are lower and gradually increase to a peak around age 9 years (Figure 1). Conversely, for weekdays plus weekend days, steps-day^{-1w} values are greater for those aged 5 years, with lower estimates for those aged 6–12 years, followed by a steeper decline among those aged 12–18 years (Figure 2).

Discussion

Although only one study represented several of the countries, we found clear differences in levels of physical activity as recorded by pedometer among young people from individual countries and regions. These findings are consistent with studies examining the worldwide decline in levels of cardiovascular fitness^{9,14,76} and increases in the prevalence of overweight youth.^{6–8} Across age,

gender, and physical activity–monitoring time frame (i.e., weekdays versus weekday plus weekend days), the Western Pacific region (Australia and New Zealand) demonstrated some of the highest, if not the highest, physical activity levels. Swedish youth are also some of the most physically active. Furthermore, the differences among countries indicate that a single global steps/day recommendation for a given age or gender may be unrealistic.

The reasons that young people in the Western Pacific re-

gion and several European countries were most physically active are unclear and cannot be directly inferred from the original studies. These findings do, however, focus attention on these countries and what might be learned from them. From a socioecologic perspective,^{77,78} these differences imply individual, environmental (both social and physical), policy, and geopolitical variations that have created overall environments and societies that beneficially promote physical activity. Higher levels of physical activity may be a result of less auto-centered lifestyles, and environments where walking and biking are more common. The least-active countries, such as the U.S., rely more heavily on cars for short trips.⁷⁹ The differences may also be a result of varying opportunities to be physically active, either within school (e.g., in daily physical education classes) or outside of school, such as participation in community sports leagues.

Many countries consistently exhibited low levels of physical activity, notably the U.S.⁸⁰ Countries seeking to adopt a lifestyle more similar to that in the U.S. should, therefore, be cautious in making environmental and/or policy-level changes toward this end, such as more motorized transportation. Doing so may potentially remove opportunities for young people to be physically active.⁸¹

A major barrier in evaluating global patterns of physical activity is a lack of standardized measures. This has led to the development of self-report instruments (e.g., IPAQ, www.ipaq.ki.se/ipaq.htm) and multi-

national projects⁸² designed to obtain comparable estimates of physical activity across nations. Self-reported physical activity, although informative, generally provides an overestimate of activity levels when compared to objectively measured physical activity,^{83,84} and it therefore serves as a coarse proxy only. Although there is no one “right” measure of physical activity,⁸⁵ it is accepted that objective measures (e.g., accelerometers, pedometers) provide a more precise estimate of physical activity.

One of the arguments against objective monitoring of physical activity at the population level is the cost-prohibitive nature of collecting such data. However, as demonstrated by the more recent NHANES^{85,86} and Canadian Fitness and Lifestyle Research Institute²³ surveillance study, objective monitoring can be done at a population level, and pedometers may be an inexpensive method. The main critique against pedometers has been their inability to provide estimates of moderate to vigorous physical activity (MVPA). However, with the advancement of technology, new features have been introduced⁸⁷ to pedometers that allow for an estimate of MVPA based on steps·minute⁻¹. In addition, unlike accelerometers,^{20,21} pedometers have a uniform unit of measure (i.e., steps·day⁻¹) that allows for direct comparisons across studies. Thus, pedometers serve as one of the few objective measures that can be readily used to identify global patterns of physical activity.

This study has several limitations. For a majority of the countries, only a few studies, representing a limited age range, were available for review, and many of the studies were not designed to collect data on a nationally representative sample. In addition, there were too few countries represented to further divide regions into smaller parcels (e.g., Mediterranean, Eastern European),⁸ limiting inferences about patterns of activity by region. Variations in data collection procedures may also account for the differences in observed steps·day⁻¹, and because younger youth are more physically active than their older peers,^{28,88} the aggregation of ages in 3-year age spans may downwardly bias steps·day⁻¹ estimates for younger youth. Finally, a number of studies were excluded because they reported data on youth across a large age span (e.g., those aged 5–12 years), on divisions of healthy- and unhealthy-weight young people, or were from select populations (e.g., indigenous people). These factors limit the inferences that can be drawn from the analyses.

Conclusion

Global monitoring of population levels of youth physical activity can inform policymakers and public health advocates regarding the health and well-being of children and adolescents. The information presented in this study

serves as preliminary data for such efforts. This study demonstrates that the use of a widely available, inexpensive tool such as a pedometer might be a feasible way to develop cross-national comparisons of physical activity among young people.

No financial disclosures were reported by the authors of this paper.

References

1. Guthold R, Ono T, Strong KL, Chatterji S, Morabia A. World-wide variability in physical inactivity: a 51-country survey. *Am J Prev Med* 2008;34(6):486–94.
2. WHO. Global strategy on diet, physical activity and health. WHO, 2004.
3. Popkin BM, Conde W, Hou N, Monteiro C. Is there a lag globally in overweight trends for children compared with adults? *Obesity* (Silver Spring) 2006;14(10):1846–53.
4. Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the U.S., 1999–2004. *JAMA* 2006;295(13):1549–55.
5. Ogden CL, Carroll MD, Flegal KM. Epidemiologic trends in overweight and obesity. *Endocrinol Metab Clin North Am* 2003;32(4):741–60, vii.
6. de Onis M, Blossner M. Prevalence and trends of overweight among preschool children in developing countries. *Am J Clin Nutr* 2000;72(4):1032–9.
7. Janssen I, Katzmarzyk PT, Boyce WF, et al. Comparison of overweight and obesity prevalence in school-aged youth from 34 countries and their relationships with physical activity and dietary patterns. *Obes Rev* 2005;6(2):123–32.
8. Kelishadi R. Childhood overweight, obesity, and the metabolic syndrome in developing countries. *Epidemiol Rev* 2007; 29:62–76.
9. Tomkinson GR, Leger LA, Olds TS, Cazorla G. Secular trends in the performance of children and adolescents (1980–2000): an analysis of 55 studies of the 20m shuttle run test in 11 countries. *Sports Med* 2003;33(4):285–300.
10. van Sluijs EM, McMinn AM, Griffin SJ. Effectiveness of interventions to promote physical activity in children and adolescents: systematic review of controlled trials. *Br Med J* 2007; 335(7622):703.
11. Harris KC, Kuramoto LK, Schulzer M, Retallack JE. Effect of school-based physical activity interventions on body mass index in children: a meta-analysis. *CMAJ* 2009;180(7):719–26.
12. Vincent SD, Pangrazi RP, Raustorp A, Tomson M, Cuddihy TF. Activity levels and body mass index of children in the U.S., Sweden, and Australia. *Med Sci Sports Exerc* 2003;35(8): 1367–73.
13. Sisson SB, Katzmarzyk PT. International prevalence of physical activity in youth and adults. *Obes Rev* 2008;9(6):606–14.
14. Beets MW, Pitetti KH. A comparison of 20 meter shuttle run performance of Midwestern youth to national and international counterparts. *Pediatr Exerc Sci* 2004;16(2):94–112.
15. Samdal O, Tynjala J, Roberts C, Sallis JF, Villberg J, Wold B. Trends in vigorous physical activity and TV watching of adolescents from 1986 to 2002 in seven European Countries. *Eur J Public Health* 2007;17(3):242–8.

16. Bauman A, Bull F, Chey T, et al. The International Prevalence Study on Physical Activity: results from 20 countries. *Int J Behav Nutr Phys Act* 2009;6(1):21.
17. Borraccino A, Lemma P, Iannotti RJ, et al. Socioeconomic effects on meeting physical activity guidelines: comparisons among 32 countries. *Med Sci Sports Exerc* 2009;41(4):749–56.
18. Bauman AE, Sallis JF. Global problems require global studies. *Am J Prev Med* 2008;34(6):544–5.
19. Wang Y, Lobstein T. Worldwide trends in childhood overweight and obesity. *Int J Pediatr Obes* 2006;1(1):11–25.
20. Guinhouya CB, Hubert H, Soubrier S, Vilhelm C, Lemdani M, Durocher A. Moderate-to-vigorous physical activity among children: discrepancies in accelerometry-based cut-off points. *Obesity (Silver Spring)* 2006;14(5):774–7.
21. Nilsson A, Brage S, Riddoch C, et al. Comparison of equations for predicting energy expenditure from accelerometer counts in children. *Scand J Med Sci Sports* 2008;18(5):643–50.
22. Beets MW, Patton MM, Edwards S. The accuracy of pedometer steps and time during walking in children. *Med Sci Sports Exerc* 2005;37(3):513–20.
23. Cameron C, Wolfe R, Craig CL. Physical activity and sport: encouraging children to be active. Ottawa ON: Canadian Fitness and Lifestyle Research Institute, 2007.
24. Duncan JS, Schofield G, Duncan EK. Step count recommendations for children based on body fat. *Prev Med* 2007;44(1):42–4.
25. Tudor-Locke C, Pangrazi RP, Corbin CB, et al. BMI-referenced standards for recommended pedometer-determined steps/day in children. *Prev Med* 2004;38(6):857–64.
26. Vincent SD, Pangrazi R. An examination of the activity patterns of elementary school children. *Pediatr Exerc Sci* 2002;14:432–41.
27. Beets MW, Le Masurier GC, Beighle A, et al. Are current body mass index referenced pedometer step count recommendations applicable to U.S. youth? *J Phys ActHealth* 2008;5(5):665–74.
28. Le Masurier GC, Beighle A, Corbin CB. Pedometer-determined physical activity levels of youth. *J Phys Act Health* 2005;2:159–68.
29. Duncan JS, Schofield G, Duncan EK. Pedometer-determined physical activity and body composition in New Zealand children. *Med Sci Sports Exerc* 2006;38(8):1402–9.
30. Bassett DR Jr, Tremblay MS, Esliger DW, Copeland JL, Barnes JD, Huntington GE. Physical activity and body mass index of children in an old order Amish community. *Med Sci Sports Exerc* 2007;39(3):410–5.
31. Fagen M, Beets MW, Brechling V, Flay BR, DuBois D, Allred C. Evaluating the implementation of school-based prevention programs: opportunities and challenges. Portland OR: American Evaluation Association, 2006.
32. Lipsey MW, Milson DB. Practical meta-analysis. Thousand Oaks CA: Sage, 2001.
33. WHO. WHO Data and Statistics. 2009. www.who.int/research/en/.
34. Naylor PJ, Macdonald HM, Warburton DE, Reed KE, McKay HA. An active school model to promote physical activity in elementary schools: action schools! BC. *Br J Sports Med* 2008;42(5):338–43.
35. Wilde BE, Corbin CB, Le Masurier GC. Free-living pedometer step counts of high school students. *Pediatr Exerc Sci* 2004;16:44–53.
36. Morgan CF, Graser SV, Pangrazi RP. A prospective study of pedometer-determined physical activity and physical self-perceptions in children. *Res Q Exerc Sport* 2008;79(2):133–40.
37. Beets MW, Vogel R, Chapman S, Pitetti KH, Cardinal BJ. Parent's social support for children's outdoor physical activity: do weekdays and weekends matter? *Sex Roles* 2007;56(1–2):125–31.
38. Beighle A, Alderman B, Morgan CF, Le MG. Seasonality in children's pedometer-measured physical activity levels. *Res Q Exerc Sport* 2008;79(2):256–60.
39. Beighle A, Pangrazi R. Measuring children's activity levels: the association between step-counts and activity time. *J Phys Act Health* 2006;3:221–9.
40. Eisenmann JC, Laurson KR, Wickel EE, Gentile D, Walsh D. Utility of pedometer step recommendations for predicting overweight in children. *Int J Obes (Lond)* 2007;31(7):1179–82.
41. Flohr JA, Todd MK, Tudor-Locke C. Pedometer-assessed physical activity in young adolescents. *Res Q Exerc Sport* 2006;77(3):309–15.
42. Le Masurier GC, Corbin CB. Steps counts among middle school students vary with aerobic fitness level. *Res Q Exerc Sport* 2006;77(1):14–22.
43. Pangrazi RP, Beighle A, Vehige T, Vack C. Impact of Promoting Lifestyle Activity for Youth (PLAY) on children's physical activity. *J Sch Health* 2003;73(8):317–21.
44. Reed J, Britternham SW, Phillips DA, Carlisle CS. A preliminary examination of the fitness levels of children who meet the President's Council physical activity recommendation. *The Physical Educator* 2007;64(3):159–68.
45. Reed JA, Metzker A, Phillips DA. Relationships between physical activity and motor skills in middle school children. *Percept Mot Skills* 2004;99(2):483–94.
46. Tudor-Locke C, Lee SM, Morgan CF, Beighle A, Pangrazi RP. Children's pedometer-determined physical activity during the segmented school day. *Med Sci Sports Exerc* 2006;38(10):1732–8.
47. Ondrak KS, Morgan DW. Factors influencing bone mineral density and content in young children. *Pediatr Exerc Sci* 2006;17:101–12.
48. Ramirez-Marrero FA, Smith BA, Sherman WM, Kirby TE. Comparison of methods to estimate physical activity and energy expenditure in African American children. *Int J Sports Med* 2004;26(5):363–71.
49. Vadiveloo M, Zhu L, Quatromoni PA. Diet and physical activity patterns of school-aged children. *J Am Diet Assoc* 2009;109(1):145–51.
50. Zizzi S, Vitullo E, Rye J, et al. Impact of a three-week pedometer intervention on high school students' daily step counts and perceptions of physical activity. *Am J Health Educ* 2006;37:35–40.
51. Cardon G, De Bourdeaudhuij I. Comparison of pedometer and accelerometer measures of physical activity in preschool children. *Pediatr Exerc Sci* 2007;19(2):205–14.
52. Sigmund E, Sigmundova D, Ansari WE. Changes in physical activity in pre-schoolers and first-grade children: longitudinal study in the Czech Republic. *Child: Care, Health Devel* 2009;35(3):376–82.
53. Romon M, Lafay L, Bresson JL, et al. Relationships between physical activity and plasma leptin levels in healthy children:

- the Fleurbaix-Laventie Ville Sante II Study. *Int J Obes Relat Metab Disord* 2004;28(10):1227–32.
54. Loucaides CA, Chedzoy SM, Bennett N. Differences in physical activity levels between urban and rural school children in Cyprus. *Health Educ Res* 2004;19(2):138–47.
 55. Raustorp A, Ludvigsson J. Secular trends of pedometer-determined physical activity in Swedish school children. *Acta Paediatr* 2007;96(12):1824–8.
 56. Raustorp A, Stahle A, Gudasic H, Kinnunen A, Mattsson E. Physical activity and self-perception in school children assessed with the Children and Youth Physical Self-Perception Profile. *Scand J Med Sci Sports* 2005;15(2):126–34.
 57. Raustorp A, Svensson K, Perlinger T. Tracking of pedometer-determined physical activity: a 5-year follow-up study of adolescents in Sweden. *Pediatr Exerc Sci* 2007;19(2):228–38.
 58. Raustorp A, Pangrazi RP, Stahle A. Physical activity level and body mass index among schoolchildren in south-eastern Sweden. *Acta Paediatr* 2004;93(3):400–4.
 59. Michaud PA, Cauderay M, Narring F, Schutz Y. Assessment of physical activity with a pedometer and its relationship with VO_{2max} among adolescents in Switzerland. *Soz Präventivmed* 2002;47(2):107–15.
 60. Duncan MJ, Al-Nakeeb Y, Woodfield L, Lyons M. Pedometer determined physical activity levels in primary school children from central England. *Prev Med* 2007;44(5):416–20.
 61. Horne PJ, Hardman CA, Lowe CF, Rowlands AV. Increasing children's physical activity: a peer modelling, rewards and pedometer-based intervention. *Eur J Clin Nutr* 2009;63(2):191–8.
 62. Parfitt G, Eston RG. The relationship between children's habitual activity level and psychological well-being. *Acta Paediatr* 2005;94(12):1791–7.
 63. Rowlands AV, Eston RG. Comparison of accelerometer and pedometer measures of physical activity in boys and girls, ages 8–10 years. *Res Q Exerc Sport* 2005;76(3):251–7.
 64. Rowlands AV, Powell SM, Eston RG, Ingledew DK. Relationship between bone mass and habitual physical activity and calcium intake in 8–11 year-old boys and girls. *Pediatr Exerc Sci* 2002;14:358–68.
 65. Gorely T, Nevill ME, Morris JG, Stensel DJ, Nevill A. Effect of a school-based intervention to promote healthy lifestyles in 7–11 year old children. *Int J Behav Nutr Phys Act* 2009;6:5.
 66. Ng J. The use of pedometer in assessing physical activity of school children. *Hong Kong J Sports Med Sports Sci* 1999:37–43.
 67. Al-Hazzaa HM. Pedometer-determined physical activity among obese and non-obese 8- to 12-year-old Saudi schoolboys. *J Physiol Anthropol* 2007;26(4):459–65.
 68. Al-Hazzaa HM, Al-Rasheedi AA. Adiposity and physical activity levels among preschool children in Jeddah, Saudi Arabia. *Saudi Med J* 2007;28(5):766–73.
 69. Cuddihy T, Michaud-Tomson L, Jones EK, Johnston AO. Exploring the relationship between daily steps, body mass index and physical activity self-esteem in female Australian adolescents. *J Exerc Sci Fitness* 2006;4:25–35.
 70. Hands B, Parker H. Pedometer-determined physical activity, BMI, and waist girth in 7- to 16-year-old children and adolescents. *J Phys Act Health* 2008;(5, 1S):S153–65.
 71. Schofield L, Mummery WK, Schofield G. Effects of a controlled pedometer-intervention trial for low-active adolescent girls. *Med Sci Sports Exerc* 2005;37(8):1414–20.
 72. Schofield L, Mummery WK, Schofield G, Hopkins W. The association of objectively determined physical activity behavior among adolescent female friends. *Res Q Exerc Sport* 2007;78(2):9–15.
 73. Duncan EK, Scott DJ, Schofield G. Pedometer-determined physical activity and active transport in girls. *Int J Behav Nutr Phys Act* 2008;5:2.
 74. Duncan JS, Hopkins WG, Schofield G, Duncan EK. Effects of weather on pedometer-determined physical activity in children. *Med Sci Sports Exerc* 2008;40(8):1432–8.
 75. Oliver M, Schofield G, McEvoy E. An integrated curriculum approach to increasing habitual physical activity in children: a feasibility study. *J Sch Health* 2006;76(2):74–9.
 76. Tomkinson GR, Olds TS. Secular changes in pediatric aerobic fitness test performance: the global picture. *Med Sport Sci* 2007;50:46–66.
 77. Sallis JF, Cervero RB, Ascher W, Henderson KA, Kraft MK, Kerr J. An ecological approach to creating active living communities. *Annu Rev Public Health* 2006;27:297–322.
 78. Sallis JF, Bauman A, Pratt M. Environmental and policy interventions to promote physical activity. *Am J Prev Med* 1998;15(4):379–97.
 79. Hu PS, Reusher TR. Summary of travel trends: 2001 National Household Travel Survey. Washington DC: U.S. Department of Transportation, Federal Highway Administration, 2004.
 80. Wang Y. Cross-national comparison of childhood obesity: the epidemic and the relationship between obesity and socioeconomic status. *Int J Epidemiol* 2001;30(5):1129–36.
 81. Prentice AM. The emerging epidemic of obesity in developing countries. *Int J Epidemiol* 2006;35(1):93–9.
 82. WHO. Global school-based student health survey. 2009. www.who.int/chp/gshs/en/.
 83. Basterfield L, Adamson AJ, Parkinson KN, Maute U, Li PX, Reilly JJ. Surveillance of physical activity in the UK is flawed: validation of the Health Survey for England Physical Activity Questionnaire. *Arch Dis Child* 2008;93(12):1054–8.
 84. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the U.S. measured by accelerometer. *Med Sci Sports Exerc* 2008;40(1):181–8.
 85. Troiano RP. Can there be a single best measure of reported physical activity? *Am J Clin Nutr* 2009;89(3):736–7.
 86. Matthews CE, Chen KY, Freedson PS, et al. Amount of time spent in sedentary behaviors in the U.S., 2003–2004. *Am J Epidemiol* 2008;167(7):875–81.
 87. Marshall SJ, Levy SS, Tudor-Locke CE, et al. Translating physical activity recommendations into a pedometer-based step goal: 3000 steps in 30 minutes. *Am J Prev Med* 2009;36(5):410–5.
 88. Nader PR, Bradley RH, Houts RM, McRitchie SL, O'Brien M. Moderate-to-vigorous physical activity from ages 9 to 15 years. *JAMA* 2008;300(3):295–305.

Appendix

Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.amepre.2009.09.045](https://doi.org/10.1016/j.amepre.2009.09.045).