

Local to Global: An Introduction to Sheaves

E. Thompson¹


¹Faculty of Science
University of Calgary

Math 511 Presentation

Motivation

Motivating Question


How can we study the relation between local and global properties of geometric spaces algebraically?



Motivation

Motivating Question

How can we study the relation between local and global properties of geometric spaces algebraically?



One Answer: Sheaves and sheaf cohomology!



UNIVERSITY OF
CALGARY

What is a sheaf?

- Throughout let $(X, \tau) \in \mathbf{Top}$.

Defⁿ: (Sheaves)

A **pre-sheaf** on X with values in \mathcal{C} is a functor

$$\mathcal{F} : \mathcal{O}(X)^{op} \rightarrow \mathcal{C}$$



What is a sheaf?

- Throughout let $(X, \tau) \in \mathbf{Top}$.

Defⁿ: (Sheaves)

A **pre-sheaf** on X with values in \mathcal{C} is a functor

$$\mathcal{F} : \mathcal{O}(X)^{op} \rightarrow \mathcal{C}$$



What is a sheaf?

- Throughout let $(X, \tau) \in \mathbf{Top}$.

Defⁿ: (Sheaves)

A **pre-sheaf** on X with values in \mathcal{C} is a functor

$$\mathcal{F} : \mathcal{O}(X)^{op} \rightarrow \mathcal{C}$$

If $\forall U \in \mathcal{O}(X)$ \mathcal{F} satisfies

- $\forall U = \bigcup_{i \in I} U_i, \forall s_i \in \mathcal{F}(U_i),$

$$\forall i, j \in I (s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j}) \implies \exists ! s \in \mathcal{F}(U), \forall i \in I (s|_{U_i} = s_i)$$

it is called a **sheaf**



Example: Smooth Manifolds


Eg: Smooth Manifolds

A smooth manifold is a pair (M, \mathcal{O}_M) , with $M \in \mathbf{Top}$ and $\forall U \in M$, $\mathcal{O}_M(U) = \text{smooth real-valued functions, satisfying}$

- $\forall p \in M, \exists U, p \in U$, such that

$$(U, \mathcal{O}_M|_U) \cong (\mathbb{R}^n, \mathcal{O}_{C^\infty})$$

for some $n \in \mathbb{N}$



Maps of sheaves

Defⁿ: (Sheaf Map)

A map between sheaves $\mathcal{F}, \mathcal{G} : \mathcal{O}(X)^{op} \rightarrow \mathcal{C}$ is a collection

$$(\eta_U \in \text{Hom}_{\mathcal{C}}(\mathcal{F}(U), \mathcal{G}(U)))_{U \in \mathcal{O}(X)}$$

such that the diagram commutes for any $U \subseteq V \in \mathcal{O}(X)$.

$$\begin{array}{ccc} \mathcal{F}(V) & \xrightarrow{|_U} & \mathcal{F}(U) \\ \eta_V \downarrow & & \downarrow \eta_U \\ \mathcal{G}(V) & \xrightarrow{|_U} & \mathcal{G}(U) \end{array}$$



Maps of sheaves


Defⁿ: (Sheaf Map)

A map between sheaves $\mathcal{F}, \mathcal{G} : \mathcal{O}(X)^{op} \rightarrow \mathcal{C}$ is a collection

$$(\eta_U \in \text{Hom}_{\mathcal{C}}(\mathcal{F}(U), \mathcal{G}(U)))_{U \in \mathcal{O}(X)}$$

such that the diagram commutes for any $U \subseteq V \in \mathcal{O}(X)$.

$$\begin{array}{ccc} \mathcal{F}(V) & \xrightarrow{|_U} & \mathcal{F}(U) \\ \eta_V \downarrow & & \downarrow \eta_U \\ \mathcal{G}(V) & \xrightarrow{|_U} & \mathcal{G}(U) \end{array}$$



Example: Smooth Manifolds


Eg: Smooth Manifolds

A smooth manifold is a pair (M, \mathcal{O}_M) , with $M \in \mathbf{Top}$ and $\forall U \in M, \mathcal{O}_M(U) = \text{smooth real-valued functions, satisfying}$

- $\forall p \in M, \exists U, p \in U$, such that

$$(U, \mathcal{O}_M|_U) \cong (\mathbb{R}^n, \mathcal{O}_{C^\infty})$$

for some $n \in \mathbb{N}$



Example: Smooth Manifolds

Eg: Smooth Manifolds


A smooth manifold is a pair (M, \mathcal{O}_M) , with $M \in \mathbf{Top}$ and $\forall U \in M, \mathcal{O}_M(U) = \text{smooth real-valued functions, satisfying}$

- $\forall p \in M, \exists U, p \in U$, such that

$$(U, \mathcal{O}_M|_U) \cong (\mathbb{R}^n, \mathcal{O}_{C^\infty})$$

for some $n \in \mathbb{N}$

Observation: Differentiation and other operations on functions depend only on local behaviour



Characterizing Locality Through Universality: Stalks

- Fix a sheaf $\mathcal{F} : \mathcal{O}(X)^{op} \rightarrow \mathcal{C}$

Defⁿ: (Stalks)

The **stalk** of \mathcal{F} at $x \in X$ is **colimit**

$$\mathcal{F}_x := \varinjlim_{x \in U} \mathcal{F}(U)$$




Important example: Stalks

- Fix a sheaf $\mathcal{F} : \mathcal{O}(X)^{op} \rightarrow \mathcal{C}$

Defⁿ: (Stalks)

The **stalk** of \mathcal{F} at $x \in X$ is **colimit**

$$\mathcal{F}_x := \varinjlim_{x \in U} \mathcal{F}(U)$$



Important example: Stalks

- Fix a sheaf $\mathcal{F} : \mathcal{O}(X)^{op} \rightarrow \mathcal{C}$

Defⁿ: (Stalks)

The **stalk** of \mathcal{F} at $x \in X$ is **colimit**

$$\mathcal{F}_x := \varinjlim_{x \in U} \mathcal{F}(U) \cong \coprod_{x \in U} \mathcal{F}(U) / \sim$$

A commutative diagram illustrating the relationship between sheaf sections and stalks. The diagram features three nodes: $\mathcal{F}(U)$ at the top left, $\mathcal{F}(V)$ at the bottom left, and \mathcal{F}_x in the center. A vertical arrow labeled $|_U$ points from $\mathcal{F}(V)$ to $\mathcal{F}(U)$. A diagonal arrow labeled $p_{U,x}$ points from $\mathcal{F}(U)$ to \mathcal{F}_x . Another diagonal arrow labeled $p_{V,x}$ points from $\mathcal{F}(V)$ to \mathcal{F}_x . A curved arrow labeled $\forall f_U$ points from $\mathcal{F}(U)$ to a node $\forall L$ on the right. A curved arrow labeled $\forall f_V$ points from $\mathcal{F}(V)$ to $\forall L$. A dashed arrow labeled $\exists ! \rightarrow$ points from \mathcal{F}_x to $\forall L$.




Important example: Stalks

- Fix a sheaf $\mathcal{F} : \mathcal{O}(X)^{op} \rightarrow \mathcal{C}$


Defⁿ: (Stalks)

The **stalk** of \mathcal{F} at $x \in X$ is **colimit**

$$\mathcal{F}_x := \varinjlim_{x \in U} \mathcal{F}(U) \cong \coprod_{x \in U} \mathcal{F}(U) / \sim$$



$$(s \in \mathcal{F}(V)) \sim (r \in \mathcal{F}(U)) \implies$$



Exact

A sequence of sheaves on X , $0 \rightarrow \mathcal{H} \xrightarrow{\eta} \mathcal{F} \xrightarrow{\mu} \mathcal{G} \rightarrow 0$, induces a sequence

$$\begin{array}{ccccccc}
 \mathcal{H}(U) & \xrightarrow{\eta_U} & \mathcal{F}(U) & \xrightarrow{\mu_U} & \mathcal{G}(U) & & \\
 \uparrow & \searrow & \uparrow & \searrow & \uparrow & \searrow & \\
 0 & \longrightarrow & \mathcal{H}_x & \xrightarrow{\eta_x} & \mathcal{F}_x & \xrightarrow{\mu_x} & \mathcal{G}_x \longrightarrow 0 \\
 \uparrow & \nearrow & \uparrow & \nearrow & \uparrow & \nearrow & \\
 \mathcal{H}(V) & \xrightarrow{\eta_V} & \mathcal{F}(V) & \xrightarrow{\mu_V} & \mathcal{G}(V) & &
 \end{array}$$



Exact

A sequence of sheaves on X , $0 \rightarrow \mathcal{H} \xrightarrow{\eta} \mathcal{F} \xrightarrow{\mu} \mathcal{G} \rightarrow 0$, induces a sequence

$$\begin{array}{ccccccc} \mathcal{H}(U) & \xrightarrow{\eta_U} & \mathcal{F}(U) & \xrightarrow{\mu_U} & \mathcal{G}(U) & & \\ \uparrow & \searrow & \uparrow & \searrow & \uparrow & \searrow & \\ 0 & \xrightarrow{\quad} & \mathcal{H}_x & \xrightarrow{\eta_x} & \mathcal{F}_x & \xrightarrow{\mu_x} & \mathcal{G}_x \longrightarrow 0 \\ \uparrow & \nearrow & \uparrow & \nearrow & \uparrow & \nearrow & \\ \mathcal{H}(V) & \xrightarrow{\eta_V} & \mathcal{F}(V) & \xrightarrow{\mu_V} & \mathcal{G}(V) & & \end{array}$$

Remark. The original sequence is exact if and only if

$$0 \rightarrow \mathcal{H}_x \rightarrow \mathcal{F}_x \rightarrow \mathcal{G}_x \rightarrow 0$$

is exact $\forall x \in X$.



Exact

A sequence of sheaves on X , $0 \rightarrow \mathcal{H} \xrightarrow{\eta} \mathcal{F} \xrightarrow{\mu} \mathcal{G} \rightarrow 0$, induces a sequence

$$\begin{array}{ccccccc} \mathcal{H}(U) & \xrightarrow{\eta_U} & \mathcal{F}(U) & \xrightarrow{\mu_U} & \mathcal{G}(U) & & \\ \uparrow & \searrow & \uparrow & \searrow & \uparrow & \searrow & \\ 0 & \longrightarrow & \mathcal{H}_x & \xrightarrow{\eta_x} & \mathcal{F}_x & \xrightarrow{\mu_x} & \mathcal{G}_x \longrightarrow 0 \\ \uparrow & \nearrow & \uparrow & \nearrow & \uparrow & \nearrow & \\ \mathcal{H}(V) & \xrightarrow{\eta_V} & \mathcal{F}(V) & \xrightarrow{\mu_V} & \mathcal{G}(V) & & \end{array}$$

Remark. The original sequence is exact if and only if

$$0 \rightarrow \mathcal{H}_x \rightarrow \mathcal{F}_x \rightarrow \mathcal{G}_x \rightarrow 0$$

is exact $\forall x \in X$.

Note

Surjectivity is local!

CALGARY

The Structure of Geometric Spaces: Category of Ringed Spaces

Defⁿ: (Ringed Space)

A **ringed space** is a pair (X, \mathcal{O}_X) of $X \in \mathbf{Top}$, and $\mathcal{O}_X : \mathcal{O}(X)^{op} \rightarrow \mathbf{Ring}$ a sheaf of rings



The Structure of Geometric Spaces: Category of Ringed Spaces

Defⁿ: (Ringed Space)

A **ringed space** is a pair (X, \mathcal{O}_X) of $X \in \mathbf{Top}$, and $\mathcal{O}_X : \mathcal{O}(X)^{op} \rightarrow \mathbf{Ring}$ a sheaf of rings

Defⁿ: (Maps of Ringed Spaces)

A map of ringed spaces $(X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ is a pair of maps $\varphi : X \rightarrow Y$ and $\varphi^\# : \mathcal{O}_Y \rightarrow \varphi_* \mathcal{O}_X$



Inducing Sheaves

- Fix a continuous map $f : X \rightarrow Y$ and a sheaf \mathcal{F} on X over \mathcal{C}

- Fix a continuous map $f : X \rightarrow Y$ and a sheaf \mathcal{F} on X over \mathcal{C}

Defⁿ: (Push-forward)

The push-forward of \mathcal{F} along f is the pre-sheaf

$$f_*\mathcal{F} : \mathcal{O}(Y)^{op} \rightarrow \mathcal{C}$$

given by $f_*\mathcal{F}(V) = \mathcal{F}(f^{-1}(V))$



The Structure of Geometric Spaces: Category of Ringed Spaces

Defⁿ: (Maps of Ringed Spaces)

A map of ringed spaces $(X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ is a pair of maps $\varphi : X \rightarrow Y$ and $\varphi^\# : \mathcal{O}_Y \rightarrow \varphi_* \mathcal{O}_X$



The Structure of Geometric Spaces: Category of Ringed Spaces

Defⁿ: (Maps of Ringed Spaces)

A map of ringed spaces $(X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ is a pair of maps $\varphi : X \rightarrow Y$ and $\varphi^\# : \mathcal{O}_Y \rightarrow \varphi_* \mathcal{O}_X$

Defⁿ: (Sheaf of \mathcal{O}_X -modules)

A sheaf of \mathcal{O}_X -**modules** on (X, \mathcal{O}_X) is a sheaf $\mathcal{F} : \mathcal{O}(X)^{op} \rightarrow \mathbf{Ab}$ for which $\forall U \in \mathcal{O}(X)$, $\mathcal{F}(U) \in \mathcal{O}_X(U)\text{-}\mathbf{Mod}$



The Structure of Geometric Spaces: Category of Ringed Spaces

Defⁿ: (Maps of Ringed Spaces)

A map of ringed spaces $(X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ is a pair of maps $\varphi : X \rightarrow Y$ and $\varphi^\# : \mathcal{O}_Y \rightarrow \varphi_* \mathcal{O}_X$

Defⁿ: (Sheaf of \mathcal{O}_X -modules)

A sheaf of \mathcal{O}_X -**modules** on (X, \mathcal{O}_X) is a sheaf $\mathcal{F} : \mathcal{O}(X)^{op} \rightarrow \mathbf{Ab}$ for which $\forall U \in \mathcal{O}(X)$, $\mathcal{F}(U) \in \mathcal{O}_X(U)\text{-Mod}$

$$\begin{array}{ccc} \mathcal{O}_X(V) \times \mathcal{F}(V) & \longrightarrow & \mathcal{F}(V) \\ |_{U \times U} \downarrow & & \downarrow |_U \\ \mathcal{O}_X(U) \times \mathcal{F}(U) & \longrightarrow & \mathcal{F}(U) \end{array}$$

Example: Smooth Manifolds Revisited


Eg: Smooth Manifolds Revisited

- Let (M, \mathcal{O}_M) be a smooth manifold
- Let $TM = \coprod_{p \in M} T_p M$ denote the tangent bundle
- The sheaf $V : \mathcal{O}(M)^{op} \rightarrow \mathbf{Ab}$ of smooth vector fields is a $\mathcal{O}_M\text{-Mod}$

Example: Smooth Manifolds Revisited

Eg: Smooth Manifolds Revisited

- Let (M, \mathcal{O}_M) be a smooth manifold
- Let $TM = \coprod_{p \in M} T_p M$ denote the tangent bundle
- The sheaf $V : \mathcal{O}(M)^{op} \rightarrow \mathbf{Ab}$ of smooth vector fields is a \mathcal{O}_M -**Mod**



Global sections functor

- The **global sections functor**, $\Gamma : \mathcal{O}_X\text{-Mod} \rightarrow \mathcal{O}_X(X)\text{-Mod}$, is given by $\Gamma(\mathcal{F}) = \mathcal{F}(X)$

Global sections functor

- The **global sections functor**, $\Gamma : \mathcal{O}_X\text{-Mod} \rightarrow \mathcal{O}_X(X)\text{-Mod}$, is given by $\Gamma(\mathcal{F}) = \mathcal{F}(X)$

Example of Surjectivity Failure:

- Let $X = \mathbb{C} \cup \{\infty\}$,

Global sections functor

- The **global sections functor**, $\Gamma : \mathcal{O}_X\text{-Mod} \rightarrow \mathcal{O}_X(X)\text{-Mod}$, is given by $\Gamma(\mathcal{F}) = \mathcal{F}(X)$

Example of Surjectivity Failure:

- Let $X = \mathbb{C} \cup \{\infty\}$,
- Let $\mathcal{A}_0, \mathcal{A}_\infty, \mathcal{A} : \mathcal{O}(X)^{op} \rightarrow \mathbf{Ab}$ be sheaves of analytic functions, with the ones in \mathcal{A}_0 vanishing at 0 and the ones in \mathcal{A}_∞ vanishing at ∞

Global sections functor

- The **global sections functor**, $\Gamma : \mathcal{O}_X\text{-Mod} \rightarrow \mathcal{O}_X(X)\text{-Mod}$, is given by $\Gamma(\mathcal{F}) = \mathcal{F}(X)$

Example of Surjectivity Failure:

- Let $X = \mathbb{C} \cup \{\infty\}$,
- Let $\mathcal{A}_0, \mathcal{A}_\infty, \mathcal{A} : \mathcal{O}(X)^{op} \rightarrow \mathbf{Ab}$ be sheaves of analytic functions, with the ones in \mathcal{A}_0 vanishing at 0 and the ones in \mathcal{A}_∞ vanishing at ∞
- Take the map $\mathcal{A}_0 \oplus \mathcal{A}_\infty \rightarrow \mathcal{A}$ given by addition



Global sections functor

- The **global sections functor**, $\Gamma : \mathcal{O}_X\text{-Mod} \rightarrow \mathcal{O}_X(X)\text{-Mod}$, is given by $\Gamma(\mathcal{F}) = \mathcal{F}(X)$

Example of Surjectivity Failure:

- Let $X = \mathbb{C} \cup \{\infty\}$,
- Let $\mathcal{A}_0, \mathcal{A}_\infty, \mathcal{A} : \mathcal{O}(X)^{op} \rightarrow \mathbf{Ab}$ be sheaves of analytic functions, with the ones in \mathcal{A}_0 vanishing at 0 and the ones in \mathcal{A}_∞ vanishing at ∞
- Take the map $\mathcal{A}_0 \oplus \mathcal{A}_\infty \rightarrow \mathcal{A}$ given by addition




Figure: Riemann Sphere



UNIVERSITY OF
CALGARY

Global sections functor

- The **global sections functor**, $\Gamma : \mathcal{O}_X\text{-Mod} \rightarrow \mathcal{O}_X(X)\text{-Mod}$, is given by $\Gamma(\mathcal{F}) = \mathcal{F}(X)$

Example of Surjectivity Failure:

- Let $X = \mathbb{C} \cup \{\infty\}$,
- Let $\mathcal{A}_0, \mathcal{A}_\infty, \mathcal{A} : \mathcal{O}(X)^{op} \rightarrow \mathbf{Ab}$ be sheaves of analytic functions, with the ones in \mathcal{A}_0 vanishing at 0 and the ones in \mathcal{A}_∞ vanishing at ∞
- Take the map $\mathcal{A}_0 \oplus \mathcal{A}_\infty \rightarrow \mathcal{A}$ given by addition
- By Liouville's Theorem $\mathcal{A}(X)$ consists of all constant functions




Figure: Riemann Sphere

Global sections functor

- The **global sections functor**, $\Gamma : \mathcal{O}_X\text{-Mod} \rightarrow \mathcal{O}_X(X)\text{-Mod}$, is given by $\Gamma(\mathcal{F}) = \mathcal{F}(X)$

Prop

Γ is a left-exact functor

Proof Idea: Let $0 \rightarrow \mathcal{H} \xrightarrow{\eta} \mathcal{F} \xrightarrow{\mu} \mathcal{G} \rightarrow 0$ be a SES. This induces a diagram

$$\begin{array}{ccccccc} \Gamma(\mathcal{H}) & \xrightarrow{\eta_X} & \Gamma(\mathcal{F}) & \xrightarrow{\mu_X} & \Gamma(\mathcal{G}) & & \\ \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{H}_x & \xrightarrow{\eta_x} & \mathcal{F}_x & \xrightarrow{\mu_x} & \mathcal{G}_x \longrightarrow 0 \end{array}$$



Remark. We want to measure the failure of Γ to be right-exact.

Remark. We want to measure the failure of Γ to be right-exact.

Construction. To extend Γ , for each $\mathcal{F} \in \mathcal{O}_X\text{-}\mathbf{Mod}$ we “take an injective resolution” $0 \rightarrow \mathcal{F} \rightarrow \mathcal{I}_\bullet$ and set

$$R^n\Gamma(\mathcal{F}) = H^n(\Gamma(\mathcal{I}_\bullet))$$

for $n \in \mathbb{Z}_{\geq 0}$.

Remark. We want to measure the failure of Γ to be right-exact.

Construction. To extend Γ , for each $\mathcal{F} \in \mathcal{O}_X\text{-Mod}$ we “take an injective resolution” $0 \rightarrow \mathcal{F} \rightarrow \mathcal{I}_\bullet$ and set

$$R^n\Gamma(\mathcal{F}) = H^n(\Gamma(\mathcal{I}_\bullet))$$

for $n \in \mathbb{Z}_{\geq 0}$.

Question. Does every $\mathcal{F} \in \mathcal{O}_X\text{-Mod}$ have an injective resolution?

Existence of Sheaf Cohomology

Thm

The category $\mathcal{O}_X\text{-}\mathbf{Mod}$ has enough injectives.



UNIVERSITY OF
CALGARY

Existence of Sheaf Cohomology

Thm

The category $\mathcal{O}_X\text{-Mod}$ has enough injectives.

Proof Sketch. Fix $\mathcal{F} \in \mathcal{O}_X\text{-Mod}$.



UNIVERSITY OF
CALGARY

Existence of Sheaf Cohomology

Thm

The category $\mathcal{O}_X\text{-Mod}$ has enough injectives.

Proof Sketch. Fix $\mathcal{F} \in \mathcal{O}_X\text{-Mod}$.

- $\forall x \in X$, $\mathcal{O}_{X,x}\text{-Mod}$ has enough injectives.

Existence of Sheaf Cohomology

Thm

The category $\mathcal{O}_X\text{-Mod}$ has enough injectives.

Proof Sketch. Fix $\mathcal{F} \in \mathcal{O}_X\text{-Mod}$.

- $\forall x \in X$, $\mathcal{O}_{X,x}\text{-Mod}$ has enough injectives.
- $\implies \forall x \in X$, $\exists \iota_x : \mathcal{F}_x \hookrightarrow \mathcal{I}(x)$ in $\mathcal{O}_{X,x}\text{-Mod}$



Existence of Sheaf Cohomology

Thm

The category $\mathcal{O}_X\text{-Mod}$ has enough injectives.

Proof Sketch. Fix $\mathcal{F} \in \mathcal{O}_X\text{-Mod}$.

- $\forall x \in X$, $\mathcal{O}_{X,x}\text{-Mod}$ has enough injectives.
- $\implies \forall x \in X$, $\exists \iota_x : \mathcal{F}_x \hookrightarrow \mathcal{I}(x)$ in $\mathcal{O}_{X,x}\text{-Mod}$
- Define $\mathcal{I} : \mathcal{O}(X)^{op} \rightarrow \mathbf{Ab}$ by $\mathcal{I}(U) = \prod_{x \in U} \mathcal{I}(x)$



Existence of Sheaf Cohomology

Thm

The category $\mathcal{O}_X\text{-Mod}$ has enough injectives.

Proof Sketch. Fix $\mathcal{F} \in \mathcal{O}_X\text{-Mod}$.

- $\forall x \in X$, $\mathcal{O}_{X,x}\text{-Mod}$ has enough injectives.
- $\implies \forall x \in X$, $\exists \iota_x : \mathcal{F}_x \hookrightarrow \mathcal{I}(x)$ in $\mathcal{O}_{X,x}\text{-Mod}$
- Define $\mathcal{I} : \mathcal{O}(X)^{op} \rightarrow \mathbf{Ab}$ by $\mathcal{I}(U) = \prod_{x \in U} \mathcal{I}(x)$
- It can be shown $\mathcal{I} \in \mathcal{O}_X\text{-Mod}$ is injective, and the induced map $\iota : \mathcal{F} \hookrightarrow \mathcal{I}$ is a monomorphism



Cor

A SES of \mathcal{O}_X -modules, $0 \rightarrow \mathcal{F} \rightarrow \mathcal{H} \rightarrow \mathcal{G} \rightarrow 0$, induces a long-exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Gamma(\mathcal{F}) & \longrightarrow & \Gamma(\mathcal{H}) & \longrightarrow & \Gamma(\mathcal{G}) \\ & & & & \delta^0 & & \\ & \nearrow & R^1\Gamma(\mathcal{F}) & \longrightarrow & R^1\Gamma(\mathcal{H}) & \longrightarrow & R^1\Gamma(\mathcal{G}) \\ & & & & & & \\ & \nearrow & R^n\Gamma(\mathcal{F}) & \longrightarrow & R^n\Gamma(\mathcal{H}) & \longrightarrow & R^n\Gamma(\mathcal{G}) \end{array}$$



Cor

A SES of \mathcal{O}_X -modules, $0 \rightarrow \mathcal{F} \rightarrow \mathcal{H} \rightarrow \mathcal{G} \rightarrow 0$, induces a long-exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Gamma(\mathcal{F}) & \longrightarrow & \Gamma(\mathcal{H}) & \longrightarrow & \Gamma(\mathcal{G}) \\ & & & & \delta^0 & & \\ & \nearrow & R^1\Gamma(\mathcal{F}) & \longrightarrow & R^1\Gamma(\mathcal{H}) & \longrightarrow & R^1\Gamma(\mathcal{G}) \\ & & & & & & \\ & \nearrow & R^n\Gamma(\mathcal{F}) & \longrightarrow & R^n\Gamma(\mathcal{H}) & \longrightarrow & R^n\Gamma(\mathcal{G}) \end{array}$$

Canonical Example:

- Studying global properties of the complex logarithm

Thank you for your time!
Any questions?

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Gamma(\mathcal{F}) & \longrightarrow & \Gamma(\mathcal{H}) & \longrightarrow & \Gamma(\mathcal{G}) \\ & & & & \delta^0 & & \\ & \searrow & & & & \nearrow & \\ & & R^1\Gamma(\mathcal{F}) & \longrightarrow & R^1\Gamma(\mathcal{H}) & \longrightarrow & R^1\Gamma(\mathcal{G}) \\ & & & & & & \\ & \searrow & & & & \nearrow & \\ & & R^n\Gamma(\mathcal{F}) & \longrightarrow & R^n\Gamma(\mathcal{H}) & \longrightarrow & R^n\Gamma(\mathcal{G}) \end{array}$$

References I