
The embryo continues to develop in the uterus and it is developed into body parts such as head, face, hands, legs, etc. Based on whether the embryo develops outside or inside the body, animals are classified into oviparous and viviparous respectively.




## ❖ Oviparous animals

Animals in which embryo develops outside the body are called oviparous animals. They produce their offspring by laying eggs. In the case of birds new ones are produced from the eggs. The egg shell protects the embryo from outer environment and the embryo receives its nutrients from the egg yolk. In these animals the new born one will have different developmental stages.



For example, in butterfly, there are different developmental stages like egg, larva, pupa and adult. Each stage is different. The process in which a butterfly becomes an adult is called metamorphosis. The life cycle process can take a month to year.

### Stage 1: Eggs

In the first stage a butterfly lays eggs on a leaf. These eggs are very small and round. About five days after the eggs are laid, a tiny worm-like creature will hatch from the egg.

### Stage 2: Caterpillar (Larva)

The second stage is the caterpillar. It is also called larva. It looks like a worm. The caterpillar starts to eat leaves and flowers once it has hatched. It grows very fast because it eats a lot. As it grows fast it sheds its old skin and gets new skin. A caterpillar shedding its outgrown skin is called molting.

### Stage 3: Chrysalis (Pupa)

The third stage is the pupa. It is mostly brown or green. This is the resting stage as well as the changing stage. The caterpillar turns into a butterfly.



### Do you know?

Amphibians are animals which have double life. The early part of an amphibian's life is spent in the water. As they get older, they spend time on land. Amphibians like frog lay thousands and sometimes millions of small, soft eggs in the water.

