Situation 2:

Patterns using Ice Candy Sticks



Pari and Manimegalai made some patterns with ice candy sticks.

To make one 'T', how many ice candy sticks are used by them? (Two sticks)

To make two 'T's, how many ice candy sticks are used by them? (Four sticks)

Continuing this, they prepared the following table to find the number of ice candy sticks used by them

Number of 'T's formed	1	2	3	4	. . .	k	. . .
Number of ice candy sticks used / required	2	4	6	8	. . .	2k	. . .
	1 × 2	2 × 2	3 × 2	4 × 2	. . .	k × 2	. . .

From the above table, it is clear that if the number of 'T's required by them is *'k'* then the number of ice candy sticks required by them will be $k \times 2 = 2k$. Here *'k'* is a variable.

TRY THESE

Use a variable to write the rule, which gives the number of ice candy sticks required to make the following patterns.

(i) a pattern of letter C as 

(ii) a pattern of letter M as 

2.4 Framing Algebraic Statements



Consider that there are *'n'* number of apples in a basket. If 5 more apples are added, what will be the total number of apples in the basket now?

The total number of apples can be easily framed into an algebraic statement as '$n + 5$'. This algebraic statement '$n + 5$' tells that, whatever be the number of apples you had earlier, there are 5 more apples now in the basket.

- Suppose there are unknown number of people in a bus, say *'x'* and if 2 more people get into the bus, then there will be '$x + 2$' people in the bus.

6th_Maths_EM_Term1_Chapter 2.indd 36 19/11/2022 15:39:16