there is no kink in the lab thermometer. A laboratory thermometer has only the Celsius scale ranging from −10°C to 110°C.



Precautions to be Followed While Using a Laboratory Thermometer

- Do not tilt the thermometer while measuring the temperature. Place it upright.
- Note the reading only when the bulb has been surrounded by the substance from all sides.



In humans, the average internal temperature is 37°C (98.6°F), though it varies among individuals. However, no person always has exactly the same temperature at every moment of the day. Temperatures cycle regularly up and down through the day according to activities and external factors.

ACTIVITY 3

Measure your body temperature

Wash the thermometer preferably with an antiseptic solution. Hold it firmly by the end and give it a few jerks. These jerks will bring the level of Mercury down. Ensure that it falls below 35°C (95°F). Now place the thermometer under your tongue or arm pit. After one minute, take the thermometer out and note the reading. It tells you your body temperature. **What did you record as your body temperature?_____**

ACTIVITY 4

Use of Laboratory thermometer

- Take some water in a beaker.
- Take a laboratory thermometer and immerse its bulb end in water; holding it vertically. Ensure to dip whole portion of bulb end. The bulb end should not touch the bottom or side of the beaker.
- Observe the movement of rise of mercury. When it becomes stable, take the reading of the thermometer.
- Repeat this with hot water and take the reading.

Difference between clinical and laboratory thermometer

Clinical Thermometer	Laboratory Thermometer
Clinical thermometer is scaled from 35°C to 42°C or from 94°F to 108°F.	Laboratory thermometer is generally scaled from —10°C to 110°C.
Mercury level does not fall on its own, as there is a kink near the bulb to prevent the fall of mercury level.	Mercury level falls on its own as no kink is present.
Temperature can be read after removing the thermometer from armpit or mouth.	Temperature is read while keeping the thermometer in the source of temperature, e.g. a liquid or any other thing.
To lower the mercury level jerks are given.	No need to give jerk to lower the mercury level.
It is used for taking the body temperature.	It is used to take temperature in laboratory.

VII Std Science Term-2 EM Unit 1.indd 5 11/11/2022 18:02:56