- There are unlimited numbers of rational numbers between two rational numbers.
- Subtracting two rational numbers is the same as adding the additive inverse of the second number to the first rational number.
- Multiplying two rational numbers is the same as multiplying their numerators and denominators separately and then writing the product in the standard form.
- Dividing a rational number by another rational number is the same as multiplying the first rational number by the reciprocal of the second rational number.
- The following table is about the properties of rational numbers($\mathbb{Q}$).

$\mathbb{Q}$	Closure	Commutative	Associative	Multiplication is distributive over +/−
+	✓	✓	✓	✓
−	✓	×	×	✓
×	✓	✓	✓	-
÷	×	×	×	-

- 0 and 1 are respectively the additive and the multiplicative identities of rational numbers.
- The additive inverse for $\frac{a}{b}$ is $\frac{-a}{b}$ and vice – versa.
- The reciprocal or the multiplicative inverse of a rational number $\frac{a}{b}$ is $\frac{b}{a}$ since $\frac{a}{b}\times\frac{b}{a}=1$.
- A natural number n is called a square number, if we can find another natural number m such that $n = m^2$.
- The square root of a number n, written as $\sqrt{n}$ (or) $n^{\frac{1}{2}}$, is the number that gives n when multiplied by itself.
- The number of times a prime factor occurs in the square is equal to twice the number of times it occurs in the prime factorization of the number.
- For any two positive numbers a and b. we have

  (i) $\sqrt{ab} = \sqrt{a}\times\sqrt{b}$ and (ii) $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}\ (b\neq 0)$
- If you multiply a number by itself and then by itself again, the result is a cube number.
- The cube root of a number is the value that when cubed gives the original number.
- An expression that represents repeated multiplication of the same factor is called a power.
- The exponent corresponds to the number of times the base is used as a factor.
- Laws of Exponents: (i) $a^m \times a^n = a^{m+n}$ (ii) $\frac{a^m}{a^n} = a^{m-n}$ (iii) $(a^m)^n = a^{mn}$
- Other results: (i) $a^0 = 1$ (ii) $a^{-m} = \frac{1}{a^m}$ (iii) $a^m \times b^m = (ab)^m$ (iv) $\frac{a^m}{b^m} = \left(\frac{a}{b}\right)^m$
- To write in scientific notation, we follow the form $S \times 10^a$ where S is a number (integer or integer with decimal) between 1 and 10, but not 10 itself, and a is a positive or negative integer.

01_8th_Maths_Numbers_Chap 1.indd 48 20/11/2022 12:48:40