Coordinate of a point on the axes:


- If y=0 then the coordinate (x, 0) lies on the 'x'-axis.
  For example (2, 0) (–5, 0) (7,0) are points on the 'x'-axis.
- If x=0 then the coordinate (0,y) lies on the 'y'-axis.
  For example (0, 3) (0,–4) (0, 9) are points on the 'y'-axis.

3.9.6 Plotting the given points on a graph

Consider the following points (4,3), (–4,5), (–3,–6), (5,–2), (6,0), (0,–5)

(i) To locate (4,3).

Start from origin O, move 4 units along OX and from 4, move 3 units parallel to OY to reach M(4,3).



(ii) To locate (–4,5)

From the origin, move 4 units along OX' and from –4, move 5 units parallel to OY to reach N(–4,5).

(iii) To locate (– 3, – 6)

From the origin move 3units along OX' and from –3, move 6 units parallel to OY' to reach P(-3,-6).

(iv) To locate (5,–2)

From the origin move 5 points along OX and from 5, move 2 units parallel to OY' to reach Q(5,–2).

(v) To locate (6,0) and (0, – 5)

In the given point (6, 0), X-coordinate is 6 and Y-coordinate is zero. So the point lies on the x-axis. Move 6 units on OX from the origin to reach R(6, 0).

In the given point (0,–5) X-coordinate is zero and Y- coordinate is (–5). So, the point lies on Y-axis. Move 5 units on OY' from the origin to reach S (0,–5).

8th_Maths_English Medium_Chapter 3.indd 110 23/11/2022 15:42:56