3.6 Division of Polynomials

Let us consider the numbers 13 and 5. When 13 is divided by 5 what is the quotient and remainder.?

Yes, of course, the quotient is 2 and the remainder is 3. We write $13 = (5\times2)+3$

Let us try.

Divide	Expressed as	Remainder	Divisor
11 by 4	$(4\times2)+3$	3	4
22 by 11	$(11\times2)+0$	0	11

Dividend = (Divisor × Quotient) + Remainder.

From the above examples, we observe that the remainder is less than the divisor.

3.6.1 Division Algorithm for Polynomials

Let $p(x)$ and $g(x)$ be two polynomials such that degree of $p(x)\geq$ degree of $g(x)$ and $g(x) \neq 0$. Then there exists unique polynomials $q(x)$ and $r(x)$ such that

$$p(x) = g(x)\times q(x) + r(x) \qquad \ldots (1)$$

where $r(x) = 0$ or degree of $r(x)$ < degree of $g(x)$.

The polynomial $p(x)$ is the Dividend, $g(x)$ is the Divisor, $q(x)$ is the Quotient and $r(x)$ is the Remainder. Now (1) can be written as

Dividend = (Divisor × Quotient) + Remainder.

If $r(x)$ is zero, then we say $p(x)$ is a multiple of $g(x)$. In other words, $g(x)$ divides $p(x)$.

If it looks complicated, don't worry! it is important to know how to divide polynomials, and that comes easily with practice. The examples below will help you.

Example 3.32

Divide $x^3 - 4x^2 + 6x$ by x, where , $x \neq 0$

Solution

We have

$$\frac{x^3 - 4x^2 + 6x}{x} = \frac{x^3}{x} - \frac{4x^2}{x} + \frac{6x}{x}, x \neq 0$$

$$= x^2 - 4x + 6$$

Example 3.33

Find the quotient and the remainder when $(5x^2 - 7x + 2) \div (x - 1)$

Solution

$(5x^2 - 7x + 2) \div (x - 1)$

3 Algebra_Term1.indd 108 02/12/2022 21:39:49