(iv) Let the assumed mean be 145. Then we can prepare the following table:

Height(x)	d = deviation from the assumed mean	Frequency (f)	fd
140	140 – 145 = – 5	4	–20
142	142– 145 = – 3	1	– 3
150	150 – 145 = + 5	2	+10
148	148 – 145 = + 3	1	+ 3
145 (Assumed)	145 – 145 = 0	2	0
147	147 – 145 = + 2	2	+ 4
Total		$\sum f = 12$	$\sum fd = -23 + 17 = -6$

Arithmetic mean = Assumed mean + Average of the sum of deviations

$$= A + \frac{\sum fd}{\sum f} = 145 + \left(\frac{-6}{12}\right) = 145.0 - 0.5 = 144.5$$

When large numbers are involved, this method could be useful.

8.4.3 Arithmetic Mean-Grouped Frequency Distribution

When data are grouped in class intervals and presented in the form of a frequency table, we get a frequency distribution like this one:

Age (in years)	10-20	20-30	30 – 40	40 – 50	50 - 60
Number of customers	80	120	50	22	8

The above table shows the number of customers in the various age groups. For example, there are 120 customers in the age group 20 – 30, but does not say anything about the age of any individual. (When we form a grouped frequency table the identity of the individual observations is lost). Hence we need a value that represents the particular class interval. Such a value is called mid value (mid-point or class mark) The mid-point or class mark can be found using the formula given below.

Mid Value = $\frac{UCL + LCL}{2}$, *UCL – Upper Class Limit, LCL – Lower Class Limit*

In grouped frequency distribution, arithmetic mean may be computed by applying any one of the following methods.

(i) Direct Method (ii) Assumed Mean Method (iii) Step Deviation Method

8-Stats.indd 272 02/12/2022 21:51:16