Example 3.5

If $p(x) = 4x^2 - 3x + 2x^3 + 5$ and $q(x) = x^2 + 2x + 4$, then find $p(x) - q(x)$


Solution


Given Polynomial	Standard form
$p(x) = 4x^2 - 3x + 2x^3 + 5$	$2x^3 + 4x^2 - 3x + 5$
$q(x) = x^2 + 2x + 4$	$x^2 + 2x + 4$
$p(x) - q(x) = 2x^3 + 3x^2 - 5x + 1$	

We see that $p(x)$ – $q(x)$ is also a polynomial. Hence the subtraction of any two polynomials is also a polynomial.


Multiplication of Two Polynomials

Divide a rectangle with 8 units of length and 7 units of breadth into 4 rectangles as shown below, and observe that the area is same, this motivates us to study the multiplication of polynomials.





For example, considering length as $(x+1)$ and breadth as $(3x+2)$ the area of the rectangle can be found by the following way.



Hence the area of the rectangle is

$= 3x^2 + 3x + 2x + 2$

$= 3x^2 + 5x + 2$

If x is a variable and m, n are positive integers then $x^m \times x^n = x^{m+n}$.

When two polynomials are multiplied the product will also be a polynomial.

Example 3.6

Find the product $(4x - 5)$ and $(2x^2 + 3x - 6)$.

Solution

To multiply $(4x - 5)$ and $(2x^2 + 3x - 6)$ distribute each term of the first polynomial to every term of the second polynomial. In this case, we need to distribute the terms $4x$ and -5. Then gather the like terms and combine them:

$$(4x - 5)(2x^2 + 3x - 6) = 4x(2x^2 + 3x - 6) - 5(2x^2 + 3x - 6)$$

$$= 8x^3 + 12x^2 - 24x - 10x^2 - 15x + 30$$

$$= 8x^3 + 2x^2 - 39x + 30$$

3 Algebra_Term1.indd 86 02/12/2022 21:38:48