Theorem 3

The opposite angles of a parallelogram are equal.

Now that we see congruence of triangles as a good "strategy", we can look for more triangles. Consider both diagonals AC and DB. We already know that ΔADC and ΔCBA are congruent. By a similar argument we can show that ΔDAB and ΔBCD are congruent as well. Are there more congruent triangles to be found in this figure ?




Fig. 4.32

Yes. The two diagonals intersect at point O. We now see 4 new ΔAOB, ΔBOC, ΔCOD and ΔDOA. Can you see any congruent pairs among them?

Since AB and CD are parallel and equal, one good guess is that ΔAOB and ΔCOD are congruent. We could again try the ASA crierion, in which case we want $\angle OAB = \angle OCD$ and $\angle ABO = \angle CDO$. But the first of these follows from the fact that $\angle CAB = \angle ACD$ (which we already established) and observing that $\angle CAB$ and $\angle OAB$ are the same (and so also $\angle OCD$ and $\angle ACD$). We now use the fact that BD is a transversal to get that $\angle ABD = \angle CDB$, but then $\angle ABD$ is the same as $\angle ABO$, $\angle CDB$ is the same as $\angle CDO$, and we are done.

Again, we need to write down the formal proof, and we have another theorem.

Theorem 4

The diagonals of a parallelogram bisect each other.

It is time now to reinforce our concepts on parallelograms. Consider each of the given statements, in the adjacent box, one by one. Identifiy the type of parallelogram which satisfies each of the statements. Support your answer with reason.

- Each pair of its opposite sides are parallel.
- Each pair of opposite sides is equal.
- All of its angles are right angles.
- Its diagonals bisect each other.
- The diagonals are equal.
- The diagonals are perpendicular and equal.
- The diagonals are perpendicular bisectors of each other.
- Each pair of its consecutive angles is supplementary.

Now we begin with lots of interesting properties of parallelograms. Can we try and prove some property relating to two or more parallelograms ? A simple case to try is when two parallelograms share the same base, as in Fig.4.33




Fig. 4.33

We see parallelograms $ABCD$ and $ABEF$ are on the common base AB. At once we can see a pair of triangles for being congruent ΔADF and ΔBCE. We already have that $AD = BC$ and $AF = BE$. But then since $AD \| BC$ and $AF \| BE$, the angle formed by AD and AF must be the same

4-Geometry_Term1.indd 154 02/12/2022 21:42:27