b. **Sublingual glands** are the smallest glands and lie beneath the tongue.

c. **Submaxillary** or **Submandibular glands** lie at the angles of the lower jaw.




Figure 20.3 Salivary glands

The salivary glands secrete a viscous fluid called saliva, approximately 1.5 liters per day. It digests starch by the action of the enzyme **ptyalin** (amylase) in the saliva which converts starch (polysaccharide) into maltose (disaccharide). Saliva also contain an antibacterial enzyme called **lysozyme**.

Tongue: The tongue is a muscular, sensory organ which helps in mixing the food with the saliva. The taste buds on the tongue help to recognize the taste of food. The masticated food in the buccal cavity becomes a bolus which is rolled by the tongue and passed through pharynx into the oesophagus by swallowing. During swallowing, the epiglottis (a muscular flap-like structure at the tip of the glottis, beginning of trachea) closes and prevents the food from entering into trachea (wind pipe).

Pharynx: The pharynx is a membrane lined cavity behind the nose and mouth, connecting them to the oesophagus. It serves as a pathway for the movement of food from mouth to oesophagus.

Oesophagus: Oesophagus or the food pipe is a muscular-membranous canal about 22 cm in length. It conducts food from pharynx to the stomach by peristalsis (wave-like movement) produced by the rhythmic contraction and relaxation of the muscular walls of alimentary canal.

Stomach: The stomach is a wide J-shaped muscular organ located between oesophagus and the small intestine. The gastric glands present in the inner walls of the stomach secrete gastric juice. The gastric juice is colourless, highly acidic, containing mucus, hydrochloric acid and enzymes rennin (in infants) and pepsin.

Inactive pepsinogen is converted to active **pepsin** which acts on the proteins in the ingested food. **Hydrochloric acid** kills the bacteria swallowed along with food and makes the medium acidic while the mucus protects the wall of the stomach. The action of the gastric juice and churning of food in the stomach convert the bolus into a semi-digested food called **chyme**. The chyme moves to the intestine slowly through the pylorus.

More to Know

- **Rennin:** Causes curdling of milk protein caesin and increases digestion of proteins.
- **Renin:** Converts angiotensinogen to angiotensin and regulate the absorption of water and Na^+ from glomerular filtrate.

Small intestine: The small intestine is the longest part of the alimentary canal, which is a long coiled tube measuring about 5 – 7 m. It comprises three parts- duodenum, jejunum and ileum.

a. **Duodenum** is C-shaped and receives the bile duct (from liver) and pancreatic duct (from pancreas).

b. **Jejunum** is the middle part of the small intestine. It is a short region of the small intestine. The secretion of the small intestine is intestinal juice which contains the enzymes like **sucrase**, **maltase**, **lactase** and **lipase**.

c. **Ileum** forms the lower part of the small intestine and opens into the large intestine. Ileum is the longest part of the small

IX_SCI_EM_Unit-20_BIO.indd 236 30/11/2022 20:18:18