medulla consists of multitubular conical masses called the medullary pyramids or renal pyramids whose bases are adjacent to cortex. On the inner concave side of each kidney, a notch called **hilum** is present through which blood vessels and nerves enter in and the urine leaves through the Ureter.

Ureters: Ureters are thin muscular tubes emerging out from the hilum. Urine enters the ureter from the renal pelvis and is conducted along the ureter by peristaltic movements of its walls. The ureters carry urine from kidney to urinary bladder.

Urinary bladder: Urinary bladder is a sac-like structure, which lies in the pelvic cavity of the abdomen. It stores urine temporarily.

Urethra: Urethra is a membranous tube, which conducts urine to the exterior. The urethral sphincters keep the urethra closed and opens only at the time of **micturition** (urination).




Figure 20.6 Longitudinal section of human kidney

Functions of kidney

1. Maintains the fluid and electrolytes balance in our body.
2. Regulates acid-base balance of blood.
3. Maintains the osmotic pressure in blood and tissues.
4. Helps to retain the important plasma constituents like glucose and amino acids.

20.2.3 Structure of Nephron

Each kidney consists of more than one million nephrons. **Nephrons or uriniferous tubules** are structural and functional units of the kidneys. Each nephron consists of **Renal corpuscle** or **Malphigian corpuscle** and **renal tubule**. The renal corpuscle consists of a cup-shaped structure called **Bowman's capsule** containing a bunch of capillaries called **glomerulus**. Blood enters the glomerular capillaries through **afferent** arterioles and leaves out through **efferent** arterioles. The Bowman's capsule continues as the renal tubule which consists of three regions **proximal convoluted tubule**, U-shaped hair pin loop, **the loop of Henle** and the **distal convoluted tubule**. The distal convoluted tubule opens into the **collecting tubule**. The nitrogenous wastes are drained into renal pelvis which leads to ureters and stored in the urinary bladder. Urine is expelled out through the urethra.

20.2.4 Mechanism of Urine Formation

The process of urine formation includes the following three stages.

- Glomerular filtration
- Tubular reabsorption
- Tubular secretion

Glomerular filtration: Urine formation begins with the filtration of blood through epithelial walls of the glomerulus and Bowman's capsule. The filtrate is called as the glomerular filtrate. Both essential and non-essential substances present in the blood are filtered.

Tubular reabsorption: The filtrate in the proximal tubule consists of essential substances such as glucose, amino acids, vitamins, sodium, potassium, bicarbonates and water that are reabsorbed into the blood by a process of **selective reabsorption**.

Tubular secretion: Substances such as H^+ or K^+ ions are secreted into the tubule. This tubular filtrate is finally known as urine, which is **hypertonic** in man. Finally the urine passes

IX_SCI_EM_Unit-20_BIO.indd 240 30/11/2022 20:18:20