22.6 Vector Borne Diseases

Vector is an agent that acts as an **intermediate carrier** of the pathogen. Many insects and animals act as vectors. Diseases transmitted by vectors are called vector borne diseases. These vectors can transfer infecting agents from an infected person to another healthy person. Some of the insect vector borne diseases are Malaria, Filaria, Chikungunya, Dengue, and the diseases which are transmitted through animals are Bird flu and Swine flu.

22.6.1 Malaria

Malaria continues to be one of the major health problems of developing countries. Malaria is caused by **protozoan** parasite *Plasmodium*. Four species of *Plasmodium* namely, *P.vivax, P.malariae, P.falciparum* and *P.ovale* cause malaria. Malaria caused by *Plasmodium falciparum* is malignant and fatal. Approximately 300 million people around the world get infected with Malaria every year.

It spreads through the bite of an insect vector, the female *Anopheles* mosquito which feeds on human blood and usually lasts less than 10 days. A person affected by malaria will show symptoms of headache, nausea, muscular pain, chillness and shivering, followed by rapid rise in temperature. Fever subsides with profuse sweating. Use of Quinine drugs kills the stages of malaria parasite.

Know your Scientist



Sir Ronald Ross, an Indian born British doctor, is famous for his work concerning malaria. He worked in the Indian Medical Service for 25 years. He identified the developing stages of malarial parasite in the gastrointestinal tract of mosquito and proved that malaria was transmitted by mosquito. In 1902, he received the Nobel Prize for Physiology or Medicine for his work on the transmission of malaria.

22.6.2 Chikungunya

Chikungunya is caused by virus. It is transmitted in humans by the bite of infected *Aedes aegypti* mosquito during the day time. It causes severe and persistent joint pain, body rashes, headache and fever. Joint pains can last for a very long time.

Incubation period of the virus is usually 2-12 days. Chillness, high fever, vomiting, nausea, headache, persistent joint pain and difficulty in walking are the common symptoms associated with this disease. The joints get inflamed and the person finds it difficult to walk. Paracetamol is given to relieve pain and reduce fever.

22.6.3 Dengue

Dengue is known as **break bone** fever. The name break bone fever was given due to the cause of intense joint and muscle pain. Dengue fever is caused by virus. It is transmitted by *Aedes aegypti* mosquito.

Incubation period of the virus is usually 5-6 days. Onset of high fever, severe headache, muscle and joint pain, rashes, haemorrhage, fall in blood platelet count are the symptoms associated with this disease. Vomiting and abdominal pain, difficulty in breathing, minute spots on the skin signifying bleeding within the skin are also associated with dengue fever. Paracetamol is given to reduce fever and body ache. Complete rest and increased intake of fluid is essential.




Figure 22.8 Dengue

IX_SCI_EM_Unit-22_BIO.indd 265 12/7/2022 5:31:30 PM