www.tntextbooks.in

Mass percentage of the solute = 13.79 %

(ii) Mass percentage of solvent = 100 – (Mass percentage of the solute)

= 100 – 13.79

= 86.21%

3) Find the amount of urea which is to be dissolved in water to get 500 g of 10% w/w aqueous solution?

$$\text{Mass percentage (w/w)} = \frac{\text{Mass of the solute}}{\text{Mass of the solution}} \times 100$$

$$10 = \frac{\text{Mass of the urea}}{500} \times 100$$

Mass of urea = 50g

(iii) **Problem based on volume – volume percentage.**

1) A solution is made from 35 ml of Methanol and 65 ml of water. Calculate the volume percentage.

Volume of the ethanol = 35 ml

Volume of the water = 65 ml

$$\text{Volume percentage} = \frac{\text{Volume of the solute}}{\text{Volume of the solution}} \times 100$$

$$\text{Volume percentage} = \frac{\text{Volume of the solute}}{\text{Volume of the solute} + \text{Volume of the solvent}} \times 100$$

$$\text{Volume percentage} = \frac{35}{35+65} \times 100$$

$$\text{Volume percentage} = \frac{35}{100} \times 100$$

$$= 35\%$$

2) Calculate the volume of ethanol in 200 ml solution of 20% v/v aqueous solution of ethanol.

Volume of aqueous solution = 200 ml

Volume percentage = 20%

$$\text{Volume percentage} = \frac{\text{Volume of solute}}{\text{Volume of solution}} \times 100$$

$$20 = \frac{\text{Volume of ethanol}}{200} \times 100$$

$$\text{Volume of ethanol} = \frac{20 \times 200}{100} = 40 \text{ ml}$$

## Points to Remember

- A solution is a homogeneous mixture of two or more substances.
- An aqueous solution is a solution in which the solvent is water.
- A non-aqueous solution is a solution in which the solvent is a liquid, other than water
- A solution in which no more solute can be dissolved in a definite amount of the solvent at a given temperature is called saturated solution.
- An unsaturated solution is one that contains less solute than the saturated solution at a given temperature.
- A supersaturated solution is one that contains more solute than the saturated solution at a given temperature.
- Polar compounds are soluble in polar solvents.
- Non-polar compounds are soluble in non-polar solvents.
- In endothermic process, solubility of solid solute increases with increase in temperature.
- In exothermic process, solubility of solid solute decreases with increase in temperature.



10th_Science_Unit-9.indd 133 06-12-2021 18:38:13