Points to Remember

- Light is a form of energy which travels along a straight line
- The deviation in the path of light ray is called refraction.
- The ratio of speed of light in vacuum to the speed of light in a medium is defined as refractive index 'μ' of that medium.
- Lens formula

$$\frac{1}{f} = \frac{1}{v} - \frac{1}{u}$$

- Magnification (m) $= \frac{h'}{h} = \frac{v}{u}$
- Power of lens. $P = \frac{1}{f}$
- The ability of the eye lens to focus nearby as well as the distant objects is called power of accommodation of the eye.
- A microscope is an optical instrument which helps us to see the objects which are very small in dimension.
- Telescope is an optical instrument used to see the distant objects clearly.

SOLVED PROBLEMS

Problem 1

Light rays travel from vacuum into a glass whose refractive index is 1.5. If the angle of incidence is 30°, calculate the angle of refraction inside the glass.

Solution:

accorting to Snell's law,

$$\frac{\sin i}{\sin r} = \frac{\mu_2}{\mu_1}$$

$$\mu_1 \sin i = \mu_2 \sin r$$

Here $\mu_1 = 1.0$, $\mu_2 = 1.5$, $i = 30°$

$$(1.0) \sin 30° = 1.5 \sin r$$

$$1 \times \frac{1}{2} = 1.5 \sin r$$

$$\sin r = \frac{1}{2 \times 1.5} = \frac{1}{3} = (0.333)$$

$$r = \sin^{-1}(0.333)$$

$$r = 19.45°$$

Problem-2

A beam of light passing through a diverging lens of focal length 0.3m appear to be focused at a distance 0.2m behind the lens. Find the position of the object.

Solution:

$f = -0.3$ m, $v = -0.2$ m

$$\frac{1}{f} = \frac{1}{v} - \frac{1}{u}$$

$$\frac{1}{u} = \frac{1}{v} - \frac{1}{f}$$

$$\frac{1}{u} = \frac{1}{-0.2} - \frac{1}{-0.3} = \frac{-10}{6}$$

$$u = \frac{-6}{10} = -0.6 \text{ m}$$

Problem-3

A person with myopia can see objects placed at a distance of 4m. If he wants to see objects at a distance of 20m, what should be the focal length and power of the concave lens he must wear?

Solution:

Given that $x = 4$m and $y = 20$m.

Focal length of the correction lens is

$$f = \frac{xy}{x-y} \quad \text{(Refer eqn.2.7)}$$

$$f = \frac{4 \times 20}{4 - 20} = \frac{80}{-16} = -5 \text{ m}$$

Power of the correction lens

$$= \frac{1}{f} = -\frac{1}{5} = -0.2 \text{ D}$$

Problem-4

For a person with hypermeteropia, the near point has moved to 1.5m. Calculate the focal length of the correction lens in order to make his eyes normal.

Solution:

Given that, d = 1.5m; D = 25cm = 0.25m (For a normal eye).

From equation (2.8), the focal length of the correction lens is

$$f = \frac{d \times D}{d - D} = \frac{1.5 \times 0.25}{1.5 - 0.25} = \frac{0.375}{1.25} = 0.3 \text{ m}$$

10th_Science_Unit-2.indd 28 12/2/2022 4:59:17 PM