
Table 11.6 IUPAC Name of various classes of compounds

No. of carbons atoms	IUPAC Name			
	Alcohols	Aldehydes	Ketones	Carboxylic acid
1	Methanol (CH_3OH)	Methanal ($HCHO$)	-	Methanoic acid ($HCOOH$)
2	Ethanol (CH_3CH_2OH)	Ethanal (CH_3CHO)	-	Ethanoic acid (CH_3COOH)
3	Propanol ($CH_3CH_2CH_2OH$)	Propanal (CH_3CH_2CHO)	Propanone (CH_3COCH_3)	Propanoic acid (CH_3CH_2COOH)
4	Butanol ($CH_3CH_2CH_2CH_2OH$)	Butanal ($CH_3CH_2CH_2CHO$)	Butanone ($CH_3COCH_2CH_3$)	Butanoic acid ($CH_3CH_2CH_2COOH$)
5	Pentanol ($CH_3CH_2CH_2CH_2CH_2OH$)	Pentanal ($CH_3CH_2CH_2CH_2CHO$)	Pentanone ($CH_3COCH_2CH_2CH_3$)	Pentanoic acid ($CH_3CH_2CH_2CH_2COOH$)

11.6 ETHANOL (CH_3CH_2OH)

Ethanol is commonly known as alcohol. All alcoholic beverages and some cough syrups contain ethanol. Its molecular formula is C_2H_5OH. Its structural formula is



Ethanol (C_2H_5OH)



11.6.1 Manufacture of ethanol

Ethanol is manufactured in industries by the fermentation of molasses, which is a by-product obtained during the manufacture of sugar from sugarcane. Molasses is a dark coloured syrupy liquid left after the crystallization of sugar from the concentrated sugarcane juice. Molasses contain about 30% of sucrose, which cannot be separated by crystallization. It is converted into ethanol by the following steps:

(i) Dilution of molasses

Molasses is first diluted with water to bring down the concentration of sugar to about 8 to 10 percent.

(ii) Addition of Nitrogen source

Molasses usually contains enough nitrogenous matter to act as food for yeast during the fermentation process. If the nitrogen content of the molasses is poor, it may be fortified by the addition of **ammonium sulphate** or **ammonium phosphate**.

(iii) Addition of Yeast

The solution obtained in step (ii) is collected in large 'fermentation tanks' and yeast is added to it. The mixture is kept at about 303K for a few days. During this period, the enzymes invertase and zymase present in yeast, bring about the conversion of sucrose into ethanol.

$$\underset{\text{Sugar}}{C_{12}H_{22}O_{11}} + H_2O \xrightarrow{\text{invertase}} \underset{\text{glucose}}{C_6H_{12}O_6} + \underset{\text{fructose}}{C_6H_{12}O_6}$$

$$\underset{\text{glucose or fructose}}{C_6H_{12}O_6} \xrightarrow{\text{zymase}} \underset{\text{ethanol}}{2C_2H_5OH} + 2\,CO_2$$

The fermented liquid is technically called **wash**.

(iv) Distillation of 'Wash'

The fermented liquid (i.e. wash), containing 15 to 18 percent alcohol, is now subjected to fractional distillation. The main fraction drawn is an aqueous solution of ethanol which contains

10th_Science_Unit-11.indd 163 12/3/2022 2:58:09 PM