daughter plant. It has only mitotic division, no gametic fusion and daughter plants are genetically similar to the parent plant.


Vegetative reproduction may take place through

(i) Leaves: In Bryophyllum small plants grow at the leaf notches



Figure 17.1 Vegetative reproduction by leaf

(ii) Stems: In strawberry aerial weak stems touch the ground and give off adventitious roots and buds. When the connections with the parent plant is broken, the offspring beomes independent.



Figuer 17.2 Vegetative reproduction by stem

(iii) Root: Tuberous roots (Asparagus and Sweet potato) can be used for vegetative propagation.

(iv) Bulbils: In some plants the flower buds modified into globose which are called as bulbils. When these falls on the ground they grow into new plants. e.g. Agave.

(v) Other types of Vegetative Reproduction

a. Fragmentation: In filamentous algae, breaking of the filament into many fragments is called fragmentation. Each fragment having atleast one cell, may give rise to a new filament of the algae by cell division e.g. Spirogyra.




Figure 17.3 Fragmentation in Spirogyra

b. Fission: In this type the parent cell divides into two daughter cells and each cell develops into a new adult organism e.g. Amoeba.




Figure 17.4 Fission in Amoeba

c. Budding: Formation of a daughter individual from a small projection, the bud, arising on the parent body is called budding. e.g. Yeast.




Figure 17.5 Budding in Yeast

d. Regeneration: The ability of the lost body parts of an individual organism to give rise to an whole new organism is called regeneration. It takes place by specialized mass of cells e.g *Hydra* and *Planaria.*

10th_Science Unit-17.indd 244 12/3/2022 3:06:35 PM