Unit Exercise - 5



1. PQRS is a rectangle formed by joining the points $P(-1,-1)$, $Q(-1,4)$, $R(5,4)$ and $S(5,-1)$. A, B, C and D are the mid-points of PQ, QR, RS and SP respectively. Is the quadrilateral $ABCD$ a square, a rectangle or a rhombus? Justify your answer.
2. The area of a triangle is 5 sq.units. Two of its vertices are $(2,1)$ and $(3, -2)$. The third vertex is (x, y) where $y = x + 3$. Find the coordinates of the third vertex.
3. Find the area of a triangle formed by the lines $3x + y - 2 = 0$, $5x + 2y - 3 = 0$ and $2x - y - 3 = 0$
4. If vertices of a quadrilateral are at $A(-5,7)$, $B(-4,k)$, $C(-1,-6)$ and $D(4,5)$ and its area is 72 sq.units. Find the value of k.
5. Without using distance formula, show that the points $(-2,-1)$, $(4,0)$, $(3,3)$ and $(-3,2)$ are vertices of a parallelogram.
6. Find the equations of the lines, whose sum and product of intercepts are 1 and -6 respectively.
7. The owner of a milk store finds that, he can sell 980 litres of milk each week at ₹14/litre and 1220 litres of milk each week at ₹16/litre. Assuming a linear relationship between selling price and demand, how many litres could he sell weekly at ₹17/litre?
8. Find the image of the point $(3,8)$ with respect to the line $x + 3y = 7$ assuming the line to be a plane mirror.
9. Find the equation of a line passing through the point of intersection of the lines $4x + 7y - 3 = 0$ and $2x - 3y + 1 = 0$ that has equal intercepts on the axes.
10. A person standing at a junction (crossing) of two straight paths represented by the equations $2x - 3y + 4 = 0$ and $3x + 4y - 5 = 0$ seek to reach the path whose equation is $6x - 7y + 8 = 0$ in the least time. Find the equation of the path that he should follow.

Points to Remember



- The area of a triangle formed by the points (x_1,y_1), (x_2,y_2) and (x_3,y_3) is $\frac{1}{2}\{(x_1y_2 + x_2y_3 + x_3y_1) - (x_2y_1 + x_3y_2 + x_1y_3)\}$ sq.units
- Three points $A(x_1,y_1)$, $B(x_2,y_2)$ and $C(x_3,y_3)$ are collinear if and only if
  (i) area of $\Delta ABC = 0$ or $x_1y_2 + x_2y_3 + x_3y_1 = x_2y_1 + x_3y_2 + x_1y_3$
  (ii) slope of AB=slope of BC or slope of AC
- The area of a quadrilateral formed by the four points (x_1,y_1), (x_2,y_2), (x_3,y_3) and (x_4,y_4) is $\frac{1}{2}\{(x_1y_2 + x_2y_3 + x_3y_4 + x_4y_1) - (x_2y_1 + x_3y_2 + x_4y_3 + x_1y_4)\}$ sq.units.
- If a line makes an angle θ with the positive direction of X axis, then its slope $m = \tan\theta$.
- If $A(x_1,y_1)$, $B(x_2,y_2)$ are two distinct points then the slope of AB is $\frac{y_2 - y_1}{x_2 - x_1}$.
- Slope of line $ax + by + c = 0$ is $m = \frac{-a}{b}$.

10th_Maths_Chapter 5_English.indd 237 12/27/2024 3:17:05 PM