Example 8.11 48 students were asked to write the total number of hours per week they spent on watching television. With this information find the standard deviation of hours spent for watching television.

x	6	7	8	9	10	11	12
f	3	6	9	13	8	5	4

Solution

x_i	f_i	$x_i f_i$	$d_i = x_i - \bar{x}$	d_i^2	$f_i d_i^2$
6	3	18	−3	9	27
7	6	42	−2	4	24
8	9	72	−1	1	9
9	13	117	0	0	0
10	8	80	1	1	8
11	5	55	2	4	20
12	4	48	3	9	36
	$N = 48$	$\Sigma x_i f_i = 432$	$\Sigma d_i = 0$		$\Sigma f_i d_i^2 = 124$

Mean

$$\bar{x} = \frac{\Sigma x_i f_i}{N} = \frac{432}{48} = 9 \quad (\because N = \Sigma f_i)$$

Standard deviation

$$\sigma = \sqrt{\frac{\Sigma f_i d_i^2}{N}} = \sqrt{\frac{124}{48}} = \sqrt{2.58}$$

$$\sigma \simeq 1.6$$

(ii) Assumed Mean method

Let $x_1, x_2, x_3, \ldots x_n$ be the given data with frequencies $f_1, f_2, f_3, \ldots f_n$ respectively. Let $\bar{x}$ be their mean and A be the assumed mean.

$$d_i = x_i - A$$

Standard deviation, $\sigma = \sqrt{\frac{\Sigma f_i d_i^2}{N} - \left(\frac{\Sigma f_i d_i}{N}\right)^2}$

Example 8.12 The marks scored by the students in a slip test are given below. Find the standard deviation of their marks.

x	4	6	8	10	12
f	7	3	5	9	5

Solution Let the assumed mean, $A = 8$

x_i	f_i	$d_i = x_i - A$	$f_i d_i$	$f_i d_i^2$
4	7	−4	−28	112
6	3	−2	−6	12
8	5	0	0	0
10	9	2	18	36
12	5	4	20	80
	$N = 29$		$\Sigma f_i d_i = 4$	$\Sigma f_i d_i^2 = 240$

Standard deviation

$$\sigma = \sqrt{\frac{\Sigma f_i d_i^2}{N} - \left(\frac{\Sigma f_i d_i}{N}\right)^2}$$

$$= \sqrt{\frac{240}{29} - \left(\frac{4}{29}\right)^2} = \sqrt{\frac{240 \times 29 - 16}{29 \times 29}}$$

$$\sigma = \sqrt{\frac{6944}{29 \times 29}} \Rightarrow \quad \sigma \simeq 2.87$$

Calculation of Standard deviation for continuous frequency distribution

(i) Mean method

Standard deviation $\sigma = \sqrt{\frac{\Sigma f_i \left(x_i - \bar{x}\right)^2}{N}}$ where, x_i = Middle value of the i th class.

f_i = Frequency of the i th class.

10th_Maths_Chapter 8_English.indd 311 12/16/2024 10:06:35 AM