www.tntextbooks.in



Figure 2.11 Example of Ctenophora- *Pleurobrachia*

help in locomotion, hence commonly called comb jellies or sea walnuts. Bioluminescence (the ability of a living organism to emit light) is well marked in ctenophores. They lack nematocysts but possess special cells called lasso cells or colloblasts which help in food capture. Digestion is both extracellular and intracellular. Sexes are not separate (monoecious). They reproduce only by sexual means. Fertilization is external and development is indirect and includes a larval stage called cydippid larva. e.g., *Pleurobrachia* (Figure 2.11).

Examples : *Pleurobrachia and Ctenoplana.*

2.3.4 Phylum: Platyhelminthes (Flatworms)

(G. *Platy* -broad or flat; *helmin*-worm)

They have a dorsoventrally flattened body and hence called flatworms. These animals are bilaterally symmetrical, triploblastic, acoelomate with organ system level of organisation. They show moderate cephalization and unidirectional movement. They are, mostly endoparasites of animals including human beings. Hooks and suckers are present in the parasitic forms and serve as organs of attachment. Their body is not segmented, but some exhibit pseudosegmentation. Some of the parasitic flatworms absorb nutrients directly from the host through their body surface. However, flatworms like liver fluke have an incomplete digestive system. Specialized excretory cells called flame cells help in osmoregulation and excretion. Sexes are not separate (monoecious); fertilisation is internal and development is through larval stages (miracidium, sporocyst, redia, cercaria). Polyembryony is common in some flatworms (Liver flukes). Some members like *Planaria* show high regeneration capacity (Figure 2.12).

Planaria *Liverfluke* *Tapeworm*

Figure 2.12 Examples of Platyhelminthes