www.tntextbooks.in

Aqueous humour supplies nutrients and oxygen to the lens, cornea and some retinal cells. The aqueous humor is produced and drained at the same rate, maintaining a constant intra ocular pressure of about 16 mmHg. Any block in the canal of schlemm increases the infra ocular pressure of aqueous humor and leads to 'Glaucoma' where the optic nerve and the retina are compressed due to pressure.

the cornea and lens. The aperture at the centre of the iris is the **pupil** through which the light enters the inner chamber. Iris is made of two types of muscles the **dilator papillae** (the radial muscle) and the **sphincter papillae** (the circular muscle).In the bright light, the circular muscle in the iris contract; so that the size of pupil decreases and less light enters the eye. In dim light, the radial muscle in the iris contract; so that the pupil size increases and more light enters the eye. Smooth muscle present in the ciliary body is called the **ciliary muscle** which alters the convexity of the lens for near and far vision. The ability of the eyes to focus objects at varying distances is called **accommodation** which is achieved by **suspensory ligament, ciliary muscle** and **ciliary body**. The suspensory ligament extends from the ciliary body and helps to hold the lens in its upright position. The ciliary body is provided with blood capillaries that secrete a watery fluid called **aqueous humor** that fills the anterior chamber.

Retina forms the inner most layer of the eye and it contains two regions: A sheet of **pigmented epithelium** (non visual part) and **neural visual regions.** The neural retina layer contains three types of cells: photoreceptor cells – **cones and rods** (Figure 10.16 and Table 10.6), **bipolar**



Figure 10.15 L.S. of the eye