www.tntextbooks.in

Figure. 11.14: Simple goitre

Figure. 11.15: Addison's disease

phosphate excretion level decreases. Generalized convulsion, locking of jaws increased heart beat rate, increased body temperature, muscular spasm are the major symptoms of tetany.

Hyperparathyroidism is caused due to excess PTH in blood. Demineralisation of bone, cyst formation, softening of bone, loss of muscle tone, general weakness, renal disorders are the symptoms of hyperparathyroidism.

Addison's disease is caused due to hyposecretion of glucocorticoids and mineralocorticoids from the adrenal cortex. Muscular weakness, low BP., loss of appetite, vomiting, hyper pigmentation of skin, low metabolic rate, subnormal temperature, reduced blood volume, weight loss are the symptoms that occur in Addison's disease (Figure 11.15). Reduced aldosterone secretion increases urinary excretion of Na Cl. and water and decreases potassium excretion leading to dehydration.

Cushing's syndrome is caused due to excess secretion of cortisol. Obesity of the face and trunk, redness of face, hand, feet, thin skin, excessive hair growth, loss of minerals from bone (osteoporosis) systolic hypertension are features of Cushing's syndrome. Suppression of sexual function like atrophy of gonads are the other symptoms of Cushing's syndrome. (Figure 11.16)

Hypoglycaemia is due to increased secretion of insulin thereby blood glucose level decreases. In this disorder blood

Normal blood glucose level:

Preprandial : 70 – 110 mg/dl (Before food) – (Fasting)

Postprandial : 110 – 140 mg/dl (About two hours after food)