shared Nobel Prize in Medicine in the year 1945 along with Ernest Boris chain and Sir Howard Walter Florey.

1.5.1 Milestones in Mycology

1729 P.A.Micheli conducted spore culture experiments

1767 Fontana proved that Fungi could cause disease in plants


1873 C.H. Blackley proved fungi could cause allergy in Human beings

1904 A.F.Blakeslee reported heterothallism in fungi

1952 Pontecorvo and Roper reported Parasexual cycle

The word 'fungus' is derived from Latin meaning 'mushroom'. Fungi are ubiquitous, eukaryotic, achlorophyllous heterotrophic organisms. They exist in unicellular or multicellular forms. The study of fungi is called mycology. (Gr. mykes – mushroom: logos – study). P.A. Micheli is considered as founder of Mycology. Few renowned mycologists include Arthur H.R. Buller, John Webster, D.L.Hawksworth, G.C.Ainsworth, B.B.Mundkur, K.C.Mehta, C.V. Subramanian and T.S. Sadasivan.

E.J. Butler is the Father of Indian Mycology. He established Imperial Agricultural Research Institute at Pusa, Bihar. It was later shifted to New Delhi and at present known as Indian Agricultural Research Insitute (IARI).



E.J. Butler (1874-1943)

He published a book, 'Fungi and Disease in Plants' on Indian plant diseases in the year 1918.

1.5.2 General characteristic features

- Majority of fungi are made up of thin, filamentous branched structures called hyphae. A number of hyphae get interwoven to form mycelium. The cell wall of fungi is made up of a polysaccharide called **chitin** (polymer of N-acetyl glucosamine) and fungal cellulose.
- The fungal mycelium is categorised into two types based on the presence or absence of septa (Figure 1.18). In lower fungi the hypha is aseptate, multinucleate and is known as coenocytic mycelium (Example: *Albugo*). In higher fungi a septum is present between the cells of the hyphae. Example: *Fusarium*.




Figure 1.18: Types of mycelium

- The mycelium is organised into loosely or compactly interwoven fungal tissues called **plectenchyma**. It is further divided into two types **prosenchyma** and **pseudoparenchyma**. In the former type the hyphae are arranged loosely but parallel to one another. In the latter hyphae are compactly arranged and loose their identity.
- In holocarpic forms the entire thallus is converted into reproductive structure whereas in Eucarpic some regions of the thallus are involved in the reproduction other regions remain vegetative. Fungi reproduce both by asexual and sexual methods. The asexual phase is called **Anamorph** and the sexual phase is called **Teleomorph**. Fungi having both phases are called **Holomorph**.

11th Botany_EM_Chapter 1.indd 21 06/01/2023 19:44:56