The sporophyte is not free-living but attached to the photosynthetic gametophyte and derives nutrition from it. Sporophyte is differentiated into foot, seta and capsule. The foot is bulbous and is embedded in the gametophyte. It derives nutrition from the gametophyte and transfers to the sporophyte. Seta is short and connects foot and capsule. The capsule consists of single layered jacket layer and encloses numerous haploid spores and elaters. The capsule is covered by protective covering called calyptra. On maturation the capsule dehisces and spores are released. Elaters helps in the dispersal of spores. The spores under favourable conditions germinate and develop into new gametophyte. The haploid gametophytic phase alternates with diploid sporophytic phase, thus the life cycle of *Marchantia* shows alternation of generation (Figure 2.14).




Figure 2.14: Life cycle of *Marchantia*

2.5 Pteridophytes

Seedless Vascular Cryptogams



From the previous section, we are aware of the salient features of amphibious plants called bryophytes. But there is a plant group called pteridophytes which are considered as first true land plants. Further, they were the first plants to acquire vascular tissue namely xylem and phloem, hence called vascular cryptogams. Club moss, horsetails, quill worts, water ferns and tree ferns belong to this group. This chapter deals with the characteristic features of Pteridophytes.

Pteridophytes are the vascular cryptogams and were abundant in the Devonian period of Palaeozoic era (400 million years ago). These plants are mostly small, herbaceous and grow well in moist, cool and shady places where water is available. The photographs for some pteridophytes are given in Figure 2.15.



a) *Lycopodium* (Club moss) b) *Equisetum* (Horse tail) c) *Azolla* (Water fern)

Figure 2.15: Pteridophytes

2.5.1 General characteristic features of Pteridophytes:

- Plant body is sporophyte (2n) and it is the dominant phase. It is differentiated into root, stem and leaves.
- Roots are adventitious.
- Stem shows monopodial or dichotomous branching.
- Leaves may be microphyllous or megaphyllous.
- Stele is protostele but in some forms siphonostele is present (*Marsilea*)
- Tracheids are the major water conducting elements but in *Selaginella* vessels are found.

11th Botany_EM_Chapter 2.indd 45 06/01/2023 19:41:29