Megasporophylls

The megasporophylls of *Cycas* are not organised into cones. They occur in close spirals around the tip of the stem of female plant. The megasporophylls are flat and measuring 15-30 cm in length. Each megasporophyll is differentiated into a basal stalk and an upper leaf like portion. The ovules are attached to the lateral side of the sporophyll. The ovules contain megaspore and it represent the female gametophyte.

Structure of Ovule




Figure 2.30: L.S. of Ovule

Cycas produces the largest ovule of the plant kingdom. The ovules are orthotropous, unitegmic and possess a short stalk. The single integument is very thick and covers the ovule leaving a small opening called *micropyle*. The integument consists of 3 layers, the outer and inner are fleshy (**sarcotesta**), the middle layer is stony called **sclerotesta**. The inner layer remains fused with the nucellus. The nucellus grows out into a beak-like structure and the upper part dissolves and forms a cavity-like structure called **pollen chamber**. A single megaspore mother cell undergoes meiosis to form four haploid megaspores. The lowermost becomes functional and others get degenerated. The nucellus gets reduced in the form of a thin papery layer in mature seeds and encloses the female gametophyte An enlarged megaspore or the embryo-sac is present within the nucellus. An archegonial chamber with 3-6 archegonia are present in the archegonial chamber below the pollen chamber (Figure 2.30).

Pollination and Fertilization

Pollination is carried out by wind and occurs at 3 celled stage(a prothallial cell, a large tube cell and a small generative cell). Pollen grains get lodged in the pollen chamber after pollination. The generative cell divides into a stalk and a body cell. The body cell divides to produce two large multiciliated antherozoids or sperms. During fertilization, one of the male gamete or multiciliated antherozoid fuses with the egg of the archegonium to form a diploid zygote (2n). The endosperm is haploid. The interval between pollination and fertilization is 4- 6 months. The zygote undergoes mitotic division and develops into embryo. The ovule is transformed into seed. The seed has two unequal cotyledons. Germination is hypogeal. The life cycle shows alternation of generations (Figure 2.31).




Figure 2.31: Life cycle of *Cycas*

11th Botany_EM_Chapter 2.indd 56 06/01/2023 19:41:39