leaves and root at nodes. Example: *Eichhornia, Pistia.*




Figure 3.9: (c) Sucker-*Chrysanthemum* **(d)** Offset-*Eichhornia*

III. Underground stem modifications

Perennial and some biennial herbs have underground stems, which are generally known as **root stocks**. Rootstock functions as a storage and protective organ. It remains alive below the ground during unfavourable conditions and resumes growth during the favourable conditions.

Underground stems are not roots because they possess nodes, internodes, scale-leaves and buds. Rootstock also lack root cap and root hairs but they possess terminal bud which is a characteristics of stem.

1. Bulb

It is a condensed conical or convex stem surrounded by fleshy scale leaves. Example: *Allium cepa, Allium sativum.*

2. Corm

This is a succulent underground stem with an erect growing tip. The corm is surrounded by scale leaves and exhibit nodes and internodes. Example: *Amorphophallus, Colocasia, Colchicum*



Bulb- *Allium cepa*

Rhizome *Zingiber officinale*

Corm- *Colocasia esculenta*

Tuber *Solanum tuberosum*

Figure 3.10: Underground Stem Modification

3. Rhizome

This is an underground stem growing horizontally with several lateral growing tips. Rhizome posses conspicuous nodes and internodes covered by scale leaves. Example: *Zingiber officinale, Canna, Curcuma longa, Musa.*

4. Tuber

This is a succulent underground spherical or globose stem with many embedded axillary buds called **"eyes".** Example: *Solanum tuberosum, Helianthus tuberosus.*

3.6.4 Stem Branching

The mode of arrangement of branches on a stem is known as **branching.** Branching pattern is determined by the relative activity of apical meristems. Based on growth pattern stems may show indeterminate or determinate growth.

1. **Indeterminate:** The terminal bud grows uninterrupted and produce several lateral branches. This type of growth is also known as **monopodial branching.** Example: *Polyalthia, Swietenia.*

11th Botany_EM_Chapter 3.indd 70 06-01-2023 12:12:54