2. To establish the evolution of a group of taxa by understanding the evolutionary and phylogenetic trends.
3. To involve any type of data gathering based on modern concepts and not only on morphology and anatomy.
4. To recognize the various groups as separate biosystematic categories such as ecotypes, ecospecies, cenospecies and comparium.

5.11.3 Karyotaxonomy

Chromosomes are the carriers of genetic information. Increased knowledge about the chromosomes have been used for extensive biosystematic studies and resolving many taxonomic problems. Utilization of the characters and phenomena of cytology for the explanation of taxonomic problem is known as **cytotaxonomy** or **karyotaxonomy**. The characters of chromosome such as number, size, morphology and behaviour during meiosis have proved to be of taxonomic value.

5.11.4 Serotaxonomy (Immunotaxonomy)

Systematic serology or serotaxonomy had its origin towards the end of twentieth century with the discovery of serological reactions and development of the discipline of immunology. The classification of very similar plants by means of differences in the proteins they contain, to solve taxonomic problems is called **serotaxonomy**. **Smith** (1976) defined it as "**the study of the origins and properties of antisera**."

Importance of serotaxonomy

It determines the degree of similarity between species, genera, families etc. by comparing the reactions of antigens from various plant taxa with antibodies raised against the antigen of a given taxon.

Example: 1. The assignment of *Phaseolus aureus* and *P. mungo* to the genus *Vigna* is strongly supported by serological evidence by **Chrispeels** and **Gartner**.

5.11.5 Molecular taxonomy (molecular systematics / molecular phylogenetics)

Molecular Taxonomy is the branch of phylogeny that analyses hereditary molecular differences, mainly in DNA sequences, to gain information and to establish genetic relationship between the members of different taxonomic categories. The advent of DNA cloning and sequencing methods have contributed immensely to the development of molecular taxonomy and population genetics over the years. These modern methods have revolutionised the field of molecular taxonomy and population genetics with improved analytical power and precision.

The results of a molecular phylogenetic analysis are expressed in the form of a tree called **phylogenetic tree.** Different molecular markers like allozymes, mitochondrial DNA, microsatellites, RFLP (Restriction Fragment Length Polymorphism), RAPD (Random amplified polymorphic DNA), AFLPs (Amplified Fragment Length Polymorphism), single nucleotide polymorphism- (SNP), microchips or arrays are used in analysis.

Uses of molecular taxonomy

1. Molecular taxonomy helps in establishing the relationship of different plant groups at DNA level.
2. It unlocks the treasure chest of information on evolutionary history of organisms.

RFLP (Restriction Fragment Length Polymorphism)

RFLPs is a molecular method of genetic analysis that allows identification of taxa based on unique patterns of restriction sites in specific regions of DNA. It refers to differences between taxa in restriction sites and therefore the lengths of fragments of DNA following cleavage with restriction enzymes.

11th Botany_EM_Chapter 5.indd 116 06/01/2023 19:43:39