
iii. **Polyphyletic group;** Taxa that includes members from two different lineages.



Need for cladistics

1. Cladistics is now the most commonly used and accepted method for creating phylogenetic system of classifications.
2. Cladistics produces a hypothesis about the relationship of organisms to predict the phylogeny
3. Cladistics helps to elucidate mechanism of evolution.

5.13 Selected Families of Angiosperms

Dicot Families

5.13.1 Family: Fabaceae (Pea family)



Systematic position

APG classification		Bentham and Hooker classification	
Kingdom	Plantae	Kingdom	Plantae
Clade	Angio-sperms	Class	Dicotyledonae
Clade	Eudicots	Sub-class	Polypetalae
Clade	Rosids	Series	Calyciflorae
Order	Fabales	Order	Rosales
Family	Fabaceae	Family	Fabaceae

General characters

Distribution: Fabaceae includes about 741 genera and more than 20,200 species. The members are cosmopolitan in distribution but abundant in tropical and subtropical regions.

Habit: All types of habits are represented in this family. Mostly herbs (*Crotalaria*), prostrate (*Indigofera enneaphylla)* erect (*Crotalaria verrucosa*), shrubs (*Cajanus cajan*), small trees (*Sesbania*), climbers (*Clitoria*), large tree (*Pongamia, Dalbergia*), woody climber (*Mucuna*), hydrophyte (*Aeschynomene aspera*) commonly called **pith plant.**

Root: Tap root system, roots are nodulated, have tubercles containing nitrogen – fixing bacteria (*Rhizobium leguminosarum)*


Root nodule

Stem: Aerial, herbaceous, woody (*Dalbergia*) twining or climbing (*Clitoria).*

Leaf: Leaf simple or unifoliate (*Desmodium gangeticum*) bifoliate (*Zornia diphylla,*), Trifoliate (*Lablab purpureus)*, alternate, stipulate, leaf base, **pulvinate**, reticulate venation terminal leaflet modifies into a **tendril** in *Pisum sativum.*

Inflorescence: Raceme (*Crotalaria verrucosa*), panicle (*Dalbergia latifolia*) axillary solitary (*Clitoria ternatea*)

Flowers: Bracteate, bracteolate, pedicellete, complete, bisexual, pentamerous, heterochlamydeous, zygomorphic hypogynous or sometimes perigynous.

Calyx: Sepals 5, green, synsepalous, more or less united in a tube and persistant, valvate or imbricate, odd sepal is anterior in position.

Corolla: Petals 5, apopetalous, unequal and **papilionaceous**, **vexillary** or **descendingly imbricate** aestivation, all petals have claw at the base. The outer most petal is large called **standard petal** or **vexillum**, Lateral 2 petals are lanceolate and curved. They are called **wing petals** or **alae**. Anterior two petals are partly fused and are called **keel petals** or **carina** which encloses the stamens and pistil.

Androecium: Stamens 10, diadelphous, usually 9+1 (*Clitoria ternatea*). The odd stamen is posterior

11th Botany_EM_Chapter 5.indd 119 06/01/2023 19:43:40