b. Secondary wall

Secondary wall is laid during maturation of the cell. It plays a key role in determining the shape of a cell. It is thick, inelastic and is made up of cellulose and lignin. The secondary wall is divided into three sublayers termed as S_1, S_2 and S_3 where the cellulose microfibrils are compactly arranged with different orientation forming a laminated structure and the cell wall strength is increased.

c. Middle lamellae

It is the outermost layer made up of calcium and magnesium pectate, deposited at the time of cytokinesis. It is a thin amorphous layer which cements two adjacent cells. It is optically inactive (isotropic).

Plasmodesmata and Pits

Plasmodesmata act as a channel between the protoplasm of adjacent cells through which many substances pass through. Moreover, at few regions, the secondary wall layer is laid unevenly whereas the primary wall and middle lamellae are laid continuously such regions are called **pits**. The Pits of adjacent cells are opposite to each other. Each pit has a pit chamber and a pit membrane. The pit membrane has many minute pores and thus they are permeable. The pits are of two types namely simple and bordered pit.

Functions of cell wall

The cell wall plays a vital role in holding several important functions given below

1. Offers definite shape and rigidity to the cell.
2. Serves as barrier for several molecules to enter the cells.
3. Provides protection to the internal protoplasm against mechanical injury.
4. Prevents the bursting of cells by maintaining the osmotic pressure.
5. Plays a major role by acting as a mechanism of defense for the cells.

6.5.4 Cell Membrane

The cell membrane is also called **cell surface** (or) **plasma membrane**. It is a thin structure which holds the cytoplasmic content called '**cytosol**'. It is extremely thin (less than 10nm).




Figure 6.11: Model of Cell membrane




Figure 6.10: Plant cell wall

11th Botany_EM_Chapter 6.indd 152 05-01-2023 18:25:10