Manganese is required for activity of enzyme needed for synthesis of oligosaccharides and glycoproteins. Molybdenum is necessary for fixation of nitrogen by enzyme nitrogenase.

Component	% of the total cellular mass
Water	70
Proteins	15
Carbohydrates	3
Lipids	2
Nucleic acids	6
Ions	4

8.1 Water

Water is the most abundant component in living organisms. Life on earth is inevitably linked to water. Water makes up 70% of human cell and upto 95% of mass of a plant cell (Figure 8.2).




Figure 8.2: Percentage of biomolecules in cell

8.1.1 Chemistry of Water

Water is a tiny polar molecule that can readily pass through membranes. Two electronegative atoms of oxygen share a hydrogen bonds of two water molecule. Thus, they can stick together by cohesion and results in lattice formation (Figure 8.3).




Figure 8.3: Water molecule

8.1.2 Properties of Water

- Adhesion and cohesion property
- High latent heat of vaporisation
- High melting and boiling point
- Universal solvent
- Specific heat capacity

8.2 Primary and Secondary Metabolites

Most plants, fungi and other microbes synthesizes a number of organic compounds called as **metabolites** which are intermediates and products of metabolism. The term metabolite is usually restricted to small molecules. It can be catergorized into two types namely primary and secondary metabolites based on their role in metabolic process (Figure 8.4).




Figure 8.4: Synthesis of metabolites during growth

Primary metabolites are those that are required for the basic metabolic processes like photosynthesis, respiration, protein and lipid metabolism of living organisms.

Secondary metabolites does not show any direct function in growth and development of organisms.



Morphine is the first alkaloid to be found. It comes from the plant Opium poppy (*Papaver somniferum*). It is used as a pain reliever in patients with severe pain levels and cough suppressant.



XIth_Botany_Chapter_8_EM.indd 177 05-01-2023 18:27:17