Glucose has a chemical formula of $C_6H_{12}O_6$. It is a six carbon molecule and hence is called as **hexose.**

All monosaccharides contain one or two functional groups. Some are aldehydes, Eg: glucose and are referred as **aldoses**; other are ketones, Eg: fructose and are referred as **ketoses**.



Glucose is one of the most well-known molecules due to its nature as an essential nutrient for human health. You ingest glucose in your food, and then your body uses blood to carry the glucose to the cells of every organ for the purpose of energy production.

8.3.2 Disaccharides

Disaccharides are formed when **two monosaccharides** join together. An example is **sucrose**. Sucrose is formed from a molecule of α-glucose and a molecule of fructose. This is a condensation reaction releasing water. The bond formed between the glucose and fructose molecule by removal of water is called **glycosidic bond**. This is another example of strong, covalent bond.




Figure 8.5: structure of carbohydrates

In the reverse process, a disaccharide is digested to the component monosaccharide in a hydrolysis reaction. This reaction involves addition of a water (hydro) molecule and splitting (lysis) of the glycosidic bond.

8.3.3 Polysaccharides

These are made of **hundreds of monosaccharide units**. Polysaccharides also called "**Glycans**". Long chain of branched or unbranched monosaccharides are held together by glycosidic bonds. Polysaccharide is an example of giant molecule, a macromolecule and consists of only one type of monomer. Polysaccharides are insoluble in water and are sweetless. **Cellulose** is an example built from repeated units of glucose monomer (Figure 8.6).




Figure 8.6: Branched and linear polysaccharides

Depending on the function, polysaccharides are of two types - **storage polysaccharide and structural polysaccharide.**

8.3.4 Starch

Starch is a storage polysaccharide made up of repeated units of **amylose and amylopectin**. Starch grains are made up of successive layers of amylose and amylopectin, which can be seen as growth rings. Amylose is a linear, unbranched polymer which makes up 80% of starch. Amylopectin is a polymer with some 1, 6 linkages that gives it a branched structure.

8.3.5 Test for Starch

We test the presence of starch by adding a solution of iodine in potassium iodide. Iodine molecules fit nearly into the starch helix, producing a **blue-black colour**.

XIth_Botany_Chapter_8_EM.indd 179 05-01-2023 18:27:18