Dicot Root (T.S)

Diagnostic features:

- Radial vascular bundle, exarch and tetrarch xylem.
- Parenchymatous conjuctive tissue is present.
- Pith is absent.

Dicot Stem (T.S)

Diagnostic features

- Cortex differentiated, hypodermis made up of collenchyma cells.
- Conjoint, Collateral and Open vascular bundle (Cambium present)
- Vascular bundle arranged like a ring, wedge shaped vascular bundle.
- Presence of pith and primary pith rays.

c. Dicot Leaf (T.S)

Diagnostic features

- Conjoint, Collateral and closed vascular bundle.
- Mesophyll tissue differentiated into upper palisade parenchyma and lower spongy parenchyma. (Dorsiventral leaf)
- Stomata are more in number on the lower epidermis.
- Stomata surrounded by bean shaped guard cells.




Figure 5c: T.S of Dicot leaf (Sun flower leaf)

d. Monocot Root (T.S)

Diagnostic features:

- Radial vascular bundle, exarch and Polyarch xylem.
- Pith is Present.

e. Monocot Stem (T.S)

Diagnostic features

- Conjoint, Collateral and Closed vascular bundle. (Cambium absent)
- Skull shaped and scattered vascular bundle.

11th Botany_EM_Chapter 15.indd 348 11/01/2023 15:00:43