6. The number of water molecules in a drop of water weighing 0.018 g is

(a) 6.022×10^{26} (b) 6.022×10^{23}

(c) 6.022×10^{20} (d) 9.9×10^{22}

7. 1 g of an impure sample of magnesium carbonate (containing no thermally decomposable impurities) on complete thermal decomposition gave 0.44 g of carbon dioxide gas. The percentage of impurity in the sample is

(a) 0 % (b) 4.4 % (c) 16 % (d) 8.4 %

8. When 6.3 g of sodium bicarbonate is added to 30 g of acetic acid solution, the residual solution is found to weigh 33 g. The number of moles of carbon dioxide released in the reaction is

(a) 3 (b) 0.75 (c) 0.075 (d) 0.3

9. When 22.4 litres of H_2 (g) is mixed with 11.2 litres of Cl_2 (g), each at 273 K at 1 atm the moles of HCl (g), formed is equal to

(a) 2 moles of HCl (g) (b) 0.5 moles of HCl (g)

(c) 1.5 moles of HCl (g) (d) 1 moles of HCl (g)

10. Hot concentrated sulphuric acid is a moderately strong oxidising agent. Which of the following reactions does not show oxidising behaviour?

(a) $Cu + 2H_2SO_4 \rightarrow CuSO_4 + SO_2 + 2H_2O$

(b) $C + 2H_2SO_4 \rightarrow CO_2 + 2SO_2 + 2H_2O$

(c) $BaCl_2 + H_2SO_4 \rightarrow BaSO_4 + 2HCl$

(d) none of the above

11. Choose the disproportionation reaction among the following redox reactions.

(a) $3Mg\,(s) + N_2\,(g) \rightarrow Mg_3N_2\,(s)$

(b) $P_4\,(s) + 3\,NaOH + 3H_2O \rightarrow PH_3(g) + 3NaH_2PO_2\,(aq)$

(c) $Cl_2\,(g) + 2KI(aq) \rightarrow 2KCl(aq) + I_2$

(d) $Cr_2O_3\,(s) + 2Al\,(s) \rightarrow Al_2O_3(s) + 2Cr(s)$

11th Std Chemistry 001-037.indd 30 23-12-2021 15:03:31