1.6.3 Error Analysis

i) Absolute Error

The magnitude of difference between the true value and the measured value of a quantity is called absolute error. If a_1, a_2, a_3, ...a_n are the measured values of any quantity 'a' in an experiment performed n times, then the arithmetic mean of these values is called the true value (a_m) of the quantity.

$$a_m = \frac{a_1 + a_2 + a_3 + ... + a_n}{n} \text{ or}$$

$$a_m = \frac{1}{n}\sum_{i=1}^{n} a_i$$

The absolute error in measured values is given by

$$|\Delta a_1| = |a_m - a_1|$$

$$|\Delta a_2| = |a_m - a_2|$$

.....

.....

$$|\Delta a_n| = |a_m - a_n|$$

ii) Mean Absolute error

The arithmetic mean of absolute errors in all the measurements is called the mean absolute error.

$$\Delta a_m = \frac{|\Delta a_1| + |\Delta a_2| + |\Delta a_3| + ... + |\Delta a_n|}{n}$$

$$\text{or} = \frac{1}{n}\sum_{i=1}^{n} |\Delta a_i|$$

If a_m is the true value and Δa_m is the mean absolute error then the magnitude of the quantity may lie between $a_m + \Delta a_m$ and $a_m - \Delta a_m$

iii) Relative error

The ratio of the mean absolute error to the mean value is called relative error. This is also called as fractional error. Thus

$$\text{Relative error} = \frac{\textit{Mean absolute error}}{\textit{Mean value}} = \frac{\Delta a_m}{a_m}$$

Relative error expresses how large the absolute error is compared to the total size of the object measured. For example, a driver's speedometer shows that his car is travelling at 60 km h^{-1} when it is actually moving at 62 km h^{-1}. Then absolute error of speedometer is 62-60 km h^{-1} = 2 km h^{-1} Relative error of the measurement is 2 km h^{-1} / 62 km h^{-1} = 0.032.

iv) Percentage error

The relative error expressed as a percentage is called percentage error.

$$\text{Percentage error} = \frac{\Delta a_m}{a_m} \times 100\%$$

A percentage error very close to zero means one is close to the targeted value, which is good and acceptable. It is always necessary to understand whether error is due to impression of equipment used or a mistake in the experimentation.

EXAMPLE 1.4

In a series of successive measurements in an experiment, the readings of the period of oscillation of a simple pendulum were found to be 2.63s, 2.56 s, 2.42s, 2.71s and

UNIT-1(XI-Physics_Vol-1).indd 22 05-01-2022 22:11:45