To find the resultant of the two vectors we apply the triangular law of addition as follows:

Represent the vectors $\vec{A}$ and $\vec{B}$ by the two adjacent sides of a triangle taken in the same order. Then the resultant is given by the third side of the triangle taken in the reverse order as shown in Figure 2.17.




Figure 2.17 Triangle law of addition

To explain further, the head of the first vector $\vec{A}$ is connected to the tail of the second vector $\vec{B}$. Let θ be the angle between $\vec{A}$ and $\vec{B}$. Then $\vec{R}$ is the resultant vector connecting the tail of the first vector $\vec{A}$ to the head of the second vector $\vec{B}$. The magnitude of $\vec{R}$ (resultant) is given geometrically by the length of $\vec{R}$ (OQ) and the direction of the resultant vector is the angle between $\vec{R}$ and $\vec{A}$. Thus we write $\vec{R} = \vec{A} + \vec{B}$.

$$\overrightarrow{OQ} = \overrightarrow{OP} + \overrightarrow{PQ}$$

(1) Magnitude of resultant vector

The magnitude and angle of the resultant vector are determined as follows.

From Figure 2.18, consider the triangle ABN, which is obtained by extending the side OA to ON. ABN is a right angled triangle.




Figure 2.18 Resultant vector and its direction by triangle law of addition.

From Figure 2.18

$$\cos\theta = \frac{AN}{B} \therefore AN = B\cos\theta \text{ and}$$

$$\sin\theta = \frac{BN}{B} \therefore BN = B\sin\theta$$

For ΔOBN, we have $OB^2 = ON^2 + BN^2$

$$\Rightarrow R^2 = (A + B\cos\theta)^2 + (B\ \sin\theta)^2$$

$$\Rightarrow R^2 = A^2 + B^2\cos^2\theta + 2AB\cos\theta + B^2\sin^2\theta$$

$$\Rightarrow R^2 = A^2 + B^2\left(cos^2\theta + sin^2\theta\right) + 2AB\cos\theta$$

$$\Rightarrow R = \sqrt{A^2 + B^2 + 2AB\cos\theta}$$

which is the magnitude of the resultant of $\vec{A}$ and $\vec{B}$

(2) Direction of resultant vectors: If θ is the angle between $\vec{A}$ and $\vec{B}$, then

$$\left|\vec{A} + \vec{B}\right| = \sqrt{A^2 + B^2 + 2AB\cos\theta} \quad (2.1)$$

If $\vec{R}$ makes an angle α with $\vec{A}$, then in ΔOBN,

$$\tan\alpha = \frac{BN}{ON} = \frac{BN}{OA + AN}$$

$$\tan\alpha = \frac{B\sin\theta}{A + B\cos\theta}$$

$$\Rightarrow \alpha = \tan^{-1}\left(\frac{B\sin\theta}{A + B\cos\theta}\right)$$

UNIT-2(XI-Physics_Vol-1).indd 48 05-01-2022 22:13:02