EXAMPLE 2.7

Given two vectors $\vec{A} = 2\hat{i} + 4\hat{j} + 5\hat{k}$ and $\vec{B} = \hat{i} + 3\hat{j} + 6\hat{k}$, Find the product $\vec{A}.\vec{B}$, and the magnitudes of $\vec{A}$ and $\vec{B}$. What is the angle between them?

Solution

$$\vec{A}.\vec{B} = 2 + 12 + 30 = 44$$

$$\text{Magnitude A} = \sqrt{4 + 16 + 25} = \sqrt{45} \text{ units}$$

$$\text{Magnitude B} = \sqrt{1 + 9 + 36} = \sqrt{46} \text{ units}$$

The angle between the two vectors is given by

$$\theta = \cos^{-1}\left(\frac{\vec{A}.\vec{B}}{AB}\right)$$

$$= \cos^{-1}\left(\frac{44}{\sqrt{45 \times 46}}\right) = \cos^{-1}\left(\frac{44}{45.49}\right)$$

$$= \cos^{-1}(0.967)$$

$$\therefore \theta \cong 15°$$

EXAMPLE 2.8

Check whether the following vectors are orthogonal.

i) $\vec{A} = 2\hat{i} + 3\hat{j}$ and $\vec{B} = 4\hat{i} - 5\hat{j}$

ii) $\vec{C} = 5\hat{i} + 2\hat{j}$ and $\vec{D} = 2\hat{i} - 5\hat{j}$

Solution


$$\vec{A}.\vec{B} = 8 - 15 = -7 \neq 0$$

Hence $\vec{A}$ and $\vec{B}$ are not orthogonal to each other.

$$\vec{C}.\vec{D} = 10 - 10 = 0$$

Hence, $\vec{C}$ and $\vec{D}$ are orthogonal to each other.

It is also possible to geometrically show that the vectors $\vec{C}$ *and* $\vec{D}$ are orthogonal to each other. This is shown in the following Figure.



In physics, the work done by a force $\vec{F}$ to move an object through a small displacement $d\vec{r}$ is defined as,

$$W = \vec{F}.d\vec{r}$$

$$W = F\ dr\ \cos\theta$$

The work done is basically a scalar product between the force vector and the displacement vector. Apart from work done, there are other physical quantities which are also defined through scalar products.

Note **In uniform circular motion, the centripetal force is perpendicular to the displacement. Hence, work done by this force is zero.**

UNIT-2(XI-Physics_Vol-1).indd 54 05-01-2022 22:13:24