



Solution

The baby and the chains are modeled as a particle hung by two strings as shown in the figure. There are three forces acting on the baby.

i) Downward gravitational force along negative y direction (mg)
ii) Tension (T) along the two strings

These three forces are coplanar as well as concurrent as shown in the following figure.



By using Lami's theorem

$$\frac{T}{\sin(180-\theta)} = \frac{T}{\sin(180-\theta)} = \frac{mg}{\sin(2\theta)}$$

Since $\sin(180-\theta) = \sin\theta$ and $\sin(2\theta) = 2\sin\theta\cos\theta$

We get $$\frac{T}{\sin\theta} = \frac{mg}{2\sin\theta\cos\theta}$$

From this, the tension on each string is $T = \frac{mg}{2\cos\theta}$.

Note

When $\theta = 0°$, the strings are vertical and the tension on each string is $T = \frac{mg}{2}$

3.5 LAW OF CONSERVATION OF TOTAL LINEAR MOMENTUM

In nature, conservation laws play a very important role. The dynamics of motion of bodies can be analysed very effectively using conservation laws. There are three conservation laws in mechanics. Conservation of total energy, conservation of total linear momentum, and conservation of angular momentum. By combining Newton's second and third laws, we can derive the law of conservation of total linear momentum.

When two particles interact with each other, they exert equal and opposite forces on each other. The particle 1 exerts force $\vec{F}_{21}$ on particle 2 and particle 2 exerts an exactly equal and opposite force $\vec{F}_{12}$ on particle 1, according to Newton's third law.

$$\vec{F}_{21} = -\vec{F}_{12} \qquad (3.20)$$

In terms of momentum of particles, the force on each particle (Newton's second law) can be written as

$$\vec{F}_{12} = \frac{d\vec{p}_1}{dt} \quad \text{and} \quad \vec{F}_{21} = \frac{d\vec{p}_2}{dt}. \qquad (3.21)$$

UNIT-3(XI-Physics_Vol-1).indd 130 05-01-2022 22:18:00