


(a) greater acceleration along the path AB


(b) greater acceleration along the path AC

(c) same acceleration in both the paths

(d) no acceleration in both the paths.

**8.** Two blocks of masses m and 2m are placed on a smooth horizontal surface as shown. In the first case only a force $F_1$ is applied from the left. Later only a force $F_2$ is applied from the right. If the force acting at the interface of the two blocks in the two cases is same, then $F_1 : F_2$ is

(Physics Olympiad 2016)




(a) 1:1 (b) 1:2

(c) 2:1 (d) 1:3

**9.** Force acting on the particle moving with constant speed is

(a) always zero

(b) need not be zero

(c) always non zero

(d) cannot be concluded

**10.** An object of mass m begins to move on the plane inclined at an angle $\theta$. The coefficient of static friction of inclined surface is $\mu_s$. The maximum static friction experienced by the mass is

(a) mg

(b) $\mu_s$mg

(c) $\mu_s$mg sin$\theta$

(d) $\mu_s$ mg cos $\theta$

**11.** When the object is moving at constant velocity on the rough surface,

(a) net force on the object is zero

(b) no force acts on the object

(c) only external force acts on the object

(d) only kinetic friction acts on the object

**12.** When an object is at rest on the inclined rough surface,

(a) static and kinetic frictions acting on the object is zero

(b) static friction is zero but kinetic friction is not zero

(c) static friction is not zero and kinetic friction is zero

(d) static and kinetic frictions are not zero

**13.** The centrifugal force appears to exist

(a) only in inertial frames

(b) only in rotating frames

(c) in any accelerated frame

(d) both in inertial and non-inertial frames



UNIT-3(XI-Physics_Vol-1).indd 161 05-01-2022 22:18:47