separated by an infinite distance and by distance r. $U(r) = U(r) - U(\infty)$. Here we choose $U(\infty) = 0$ as the reference point. The gravitational potential energy $U(r)$ is always negative because when two masses come together slowly from infinity, work is done by the system.

The unit of gravitational potential energy $U(r)$ is Joule and it is a scalar quantity. The gravitational potential energy depends upon the two masses and the distance between them.

6.2.4 Gravitational potential energy near the surface of the Earth

It is already discussed in chapter 4 that when an object of mass m is raised to a height h, the potential energy stored in the object is mgh (Figure 6.14). This can be derived using the general expression for gravitational potential energy.




Figure 6.14 Mass placed at a distance r from the center of the Earth

Consider the Earth and mass system, with *r*, the distance between the mass m and the Earth's centre. Then the gravitational potential energy,

$$U = -\frac{GM_e m}{r} \qquad (6.30)$$

Here r = R_e+h, where R_e is the radius of the Earth. h is the height above the Earth's surface

$$U = -G\frac{M_e m}{(R_e + h)} \qquad (6.31)$$

If h << R_e, equation (6.31) can be modified as

$$U = -G\frac{M_e m}{R_e(1 + h/R_e)}$$

$$U = -G\frac{M_e m}{R_e}(1 + h/R_e)^{-1} \qquad (6.32)$$

By using Binomial expansion and neglecting the higher order terms, we get

$$U = -G\frac{M_e m}{R_e}\left(1 - \frac{h}{R_e}\right). \qquad (6.33)$$

We know that, for a mass m on the Earth's surface,

$$G\frac{M_e m}{R_e} = mgR_e \qquad (6.34)$$

Substituting equation (6.34) in (6.33) we get,

$$U = -mgR_e + mgh \qquad (6.35)$$

It is clear that the first term in the above expression is independent of the height h. For example, if the object is taken from

UNIT-6(XI-Physics_Vol-2).indd 15 02-01-2023 14:28:57