$$m\vec{a} = -\frac{GmM_e}{R_e^2}\hat{r}$$

hence, acceleration is,

$$\vec{a} = -\frac{GM_e}{R_e^2}\hat{r} \qquad (6.43)$$

The acceleration experienced by the object near the surface of the Earth due to its gravity is called acceleration due to gravity. It is denoted by the symbol g. The magnitude of acceleration due to gravity is

$$|g| = g = \frac{GM_e}{R_e^2} \qquad (6.44)$$

It is to be noted that the acceleration experienced by any object is independent of its mass. The value of g depends only on the mass and radius of the Earth. Infact, Galileo arrived at the same conclusion 400 years ago that *all objects fall towards the Earth with the same acceleration* through various quantitative experiments. The acceleration due to gravity g is found to be 9.8 m s^{-2} on the surface of the Earth near the equator.

6.3.1 Variation of g with altitude, depth and latitude

Consider an object of mass m at a height h from the surface of the Earth. Acceleration experienced by the object due to Earth is

$$g' = \frac{GM}{(R_e + h)^2} \qquad (6.45).$$




Figure 6.17(a) Mass at a height h from the center of the Earth

$$g' = \frac{GM}{R_e^2\left(1+\frac{h}{R_e}\right)^2}$$

$$g' = \frac{GM}{R_e^2}\left(1+\frac{h}{R_e}\right)^{-2}$$

If $h \ll R_e$

We can use Binomial expansion. Taking the terms upto first order

$$g' = \frac{GM}{R_e^2}\left(1-2\frac{h}{R_e}\right)$$

$$g' = g\left(1-2\frac{h}{R_e}\right) \qquad (6.46)$$

We find that $g' < g$. This means that as altitude h increases the acceleration due to gravity g decreases.

UNIT-6(XI-Physics_Vol-2).indd 19 02-01-2023 14:29:10