7.3.4 Buoyancy

When a body is partially or fully immersed in a fluid, it displaces a certain amount of fluid. The displaced fluid exerts an upward force on the body. The upward force exerted by a fluid that opposes the weight of an immersed object in a fluid is called *upthrust* or *buoyant force* and the phenomenon is called *buoyancy*.

Archimedes principle:




Figure 7.13 Archimedes principle

It states that when a body is partially or wholly immersed in a fluid, it experiences an upward thrust equal to the weight of the fluid displaced by it and its upthrust acts through the centre of gravity of the liquid displaced.

upthrust or buoyant force = weight of liquid displaced.

Law of floatation

It is well-known that boats, ships, and some wooden objects move on the upper part of the water. We say that they float. Floatation can be defined as the tendency of an object to rise up to the upper levels of the fluid or to stay on the surface of the fluid.

The law of floatation states that a body will float in a liquid if the weight of the liquid displaced by the immersed part of the body equals the weight of the body. For example, a wooden object weighs 300 kg (about 3000 N) floats in water displaces 300 kg (about 3000 N) of water.

Note

If an object floats, the volume of fluid displaced is equal to the volume of the object submerged and the percentage of the volume of the object submerged is equal to the relative density of an object with respect to the density of the fluid in which it floats.

For example, if an ice cube of density 0.9 $g\,cm^{-3}$ floats in the fresh water of density 1.0 $g\,cm^{-3}$ then the percentage volume of an object submerged in fresh water is. $\frac{0.9\,g\,cm^{-3}}{1.0\,g\,cm^{-3}} \times 100\% = 90\%$.

Conversely, if the same ice cube floats in sea water of density 1.3 $g\,cm^{-3}$, then the percentage volume of the object submerged in seawater would be $\frac{0.9\,g\,cm^{-3}}{1.3\,g\,cm^{-3}} \times 100\% = 69.23\%$ only.

EXAMPLE 7.8

A cube of wood floating in water supports a 300 g mass at the centre of its top face. When the mass is removed, the cube rises by 3 cm. Determine the volume of the cube.

UNIT-7(XI-Physics_Vol-2).indd 64 02-01-2023 14:39:54