The Sun is approximately taken as a black body. Since any object above 0 K will emit radiation, Sun also emits radiation. Its surface temperature is about 5700 K. By substituting this value in the equation (8.12),

$$\lambda_m = \frac{b}{T} = \frac{2.898 \times 10^{-3}}{5700} \approx 508\, nm$$

It is the wavelength at which maximum intensity is 508*nm*. Since the Sun's temperature is around 5700K, the spectrum of radiations emitted by Sun lie between 400 nm to 700 nm which is the visible part of the spectrum. It is shown in Figure 8.14




Figure 8.14 Wien's law and Human's vision

The humans evolved under the Sun by receiving its radiations. The human eye is sensitive only in the visible not in infrared or X-ray ranges in the spectrum.

Suppose if humans had evolved in a planet near the star Sirius (9940K), then they would have had the ability to see the Ultraviolet rays!

EXAMPLE 8.9

The power radiated by a black body A is E_A and the maximum energy radiated was at the wavelength λ_A. The power radiated by another black body B is $E_B = N\, E_A$ and the radiated energy was at the maximum wavelength, $\frac{1}{2}\lambda_A$. What is the value of N?

According to Wien's displacement law

$\lambda_{max}\, T$ = constant for both object A and B

$\lambda_A T_A = \lambda_B T_B$. Here $\lambda_B = \frac{1}{2}\lambda_A$

$$\frac{T_B}{T_A} = \frac{\lambda_A}{\lambda_B} = \frac{\lambda_A}{\left(\frac{1}{2}\right)\lambda_A} = 2$$

$$T_B = 2T_A$$

From Stefan-Boltzmann law

$$\frac{E_B}{E_A} = \left(\frac{T_B}{T_A}\right)^4 = (2)^4 = 16 = N$$

Object B has emitted at lower wavelength compared to A. So the object B would have emitted more energetic radiation than A.

8.4 THERMODYNAMICS:

8.4.1 Introduction

In the previous sections we have studied about the heat, temperature and thermal properties of matter. Thermodynamics is

UNIT-8(XI-Physics_Vol-2).indd 112 02-01-2023 15:25:41